

HIV and Stage 3 (i.e., AIDS) Infections Among Adults and Adolescents, New Mexico 2019†

DEI ARTIMENT OF	Newly Diagnosed HIV Infections in 2019*			Persons Living with HIV or Stage 3 (i.e., AIDS)								
				HIV			Stage 3 (i.e., AIDS)			Total		
	N	%	Rate ^a	N	%	Rate ^a	N	%	Rate ^a	N	%	Rate ^a
Total	138	100.0	7.9	1908	47.3	108.6	2128	52.7	121.1	4,036	100.0	229.7
Gender												
Male	118	85.5	13.6	1625	85.2	188.0	1870	87.9	216.3	3,495	86.6	404.3
Female	19	13.8	2.1	253	13.3	28.3	235	11.0	26.3	488	12.1	54.7
Transgender Men	0	0.0	N/A	1	0.1	N/A	2	0.1	N/A	3	0.1	N/A
Transgender Women	1	0.7	N/A	29	1.5	N/A	21	1.0	N/A	50	1.2	N/A
Race/Ethnicity			22.7	450	7.0	205.2	440		200.0	252		505.0
African American	9	6.5	23.7	150	7.9	395.3	110	5.2	289.9	260	6.4	685.2
American Indian/Alaska Native Asian/Pacific Islander	23	16.7 0.7	15.0 3.1	192 18	10.1 0.9	124.9 56.1	167 11	7.8 0.5	108.6 34.3	359 29	8.9 0.7	233.5 90.4
Asian/Pacific Islander Hispanic	1 73	52.9	8.9	913	47.9	110.9	954	44.8	115.8	1,867	46.3	226.7
White	29	21.0	4.1	606	31.8	85.4	840	39.5	118.4	1,446	35.8	203.7
Multi-race	3	2.2	4.1 N/A	29	1.5	85.4 N/A	46	2.2	N/A	75	1.9	N/A
Age, years ^b		2.2	IN/A	23	1.5	IN/A	40	2.2	IN/A	/3	1.5	IN/A
	2.5	40.0		445		24.7	2.5	4.0		142		40.4
13–24 25–34	26 56	18.8 40.6	7.7 19.7	117 490	6.1 25.7	34.7 172.1	25 161	1.2 7.6	7.4 56.6	651	3.5 16.1	42.1 228.7
25–34 35–44	28	20.3	19.7	490	22.5	172.1	348	16.4	138.4	778	19.3	309.3
45–54 45–54	20	14.5	8.3	430	22.5	171.0	654	30.7	270.8	1,084	26.9	448.8
45-54 55+	8	5.8	1.2	441	23.1	68.7	940	44.2	146.4	1,381	34.2	215.0
Public Health Region ^c	·	5.0	1.2	771	23.1	00.7	340	77.2	140.4	1,501	34.2	215.0
	21	15.2	11.5	218	11.4	119.2	194	9.1	106.1	412	10.2	225.3
Northwest Northeast	13	9.4	5.1	218	15.5	115.9	401	18.8	157.0	697	17.3	272.8
Metro	62	44.9	8.0	959	50.3	124.1	1075	50.5	139.1	2,034	50.4	263.3
Southeast	14	10.1	5.9	127	6.7	53.6	127	6.0	53.6	2,034	6.3	107.2
Southwest	28	20.3	9.1	308	16.1	99.6	330	15.5	106.7	638	15.8	206.3
Missing	0	0.0	N/A	0	0.0	N/A	1	0.0	N/A	1	0.0	N/A
Transmission Category			.,,			.,,			.,,	_		.,,
Gender Identity	T T						ī					
Men												
Sex with Men (MSM)	73	61.9	N/A	1,152	70.9	N/A	1,297	69.4	N/A	2,449	70.1	N/A
Intravenous Drug Use (IDU)	5	4.2	N/A	76	4.7	N/A	122	6.5	N/A	198	5.7	N/A
MSM/IDU	14	11.9	N/A	152	9.4	N/A	212	11.3	N/A	364	10.4	N/A
Sex with Female	20	16.9	N/A	91	5.6	N/A	92	4.9	N/A	183	5.2	N/A
Sex with Female, High Risk	e 0	0.0	N/A	61	3.8	N/A	76	4.1	N/A	137	3.9	N/A
Perinatal Exposure	-		N/A	2	0.1	N/A	6	0.3	N/A	8	0.2	N/A
Received Blood Products or Tissue	0	0.0	N/A	1	0.1	N/A	6	0.3	N/A	7	0.2	N/A
No Identified or Reported Risk (NIR/NRR)	d 6	5.1	N/A	90	5.5	N/A	59	3.2	N/A	149	4.3	N/A
Transgender Men												
Sex with Women	0	0.0	N/A	0	0.0	N/A	0	0.0	N/A	0	0.0	N/A
Intravenous Drug Use (IDU)	0	0.0	N/A	0	0.0	N/A	1	50.0	N/A	1	33.3	N/A
No Identified or Reported Risk (NIR/NRR)	0	0.0	N/A	1	100.0	N/A	1	50.0	N/A	2	66.7	N/A
Women												
Sex with Men	13	68.4	N/A	104	41.1	N/A	77	32.8	N/A	181	37.1	N/A
Sex with Men, High Risk	e 0	0.0	N/A	69	27.3	N/A	82	34.9	N/A	151	30.9	N/A
Intravenous Drug Use (IDU)	5	26.3	N/A	43	17.0	N/A	64	27.2	N/A	107	21.9	N/A
Perinatal Exposure	-		N/A	3	1.2	N/A	3	1.3	N/A	6	1.2	N/A
Received Blood Products or Tissue	0	0.0	N/A	1	0.4	N/A	1	0.4	N/A	2	0.4	N/A
No Identified or Reported Risk (NIR/NRR)		5.3	N/A	33	13.0	N/A	8	3.4	N/A	41	8.4	N/A
Transgender Women			·			·						
Sex with Men	1	100.0	N/A	28	96.6	N/A	19	90.5	N/A	47	94.0	N/A
Intravenous Drug Use (IDU)	0	0.0	N/A	0	0.0	N/A	1	4.8	N/A	1	2.0	N/A
No Identified or Reported Risk (NIR/NRR)		0.0	N/A	1	3.4	N/A	1	4.8	N/A	2	4.0	N/A

NOTE: Percentages across characteristics within a section of this table may not add to 100.0% as a result of rounding.

NOTE: The data presented in this report provide a minimum estimate of New Mexico residents ≥ 13

years known to be infected with HIV .

N < 20: Rate calculations for small numbers may be statistically unreliable. Fluctuations in the associated rates may be due in part to statistical unreliability rather than true change. Region: New Mexico Department of Health constructed new health regions in 2012. Regional comparison to previous years is not recommended. A map depicting New Mexico's Health Regions is

available for download from the New Mexico Department of Health at

https://ibis.health.state.nm.us/view/docs/HealthRegions2012.pdf.

Persons imprisoned in a state and county correctional or federal detention facility, including ICE facilities, or who were housed in a residential facility (e.g., drug treatment facility), were included in the data presented unless otherwise noted.

Data Source: eHARS (Enhanced HIV/AIDS Reporting System) document and person data set 2020, 2nd quarter (June 26th, 2020).

^{*} All people ≥ 13 years who were newly diagnosed with Stages 1 through 3 HIV infection while residing in New Mexico during 2019.

^a Number of infections per 100,000 NM residents 13 years and older based on Bureau of Business and Economic Research Geospatial and Population Studies Program population estimates for 2018.

b For new diagnoses, age at time of diagnosis (2019). For persons living with HIV Stages 1 to 3, current age.

^c For new diagnoses, residence at time of diagnosis (2019). For persons living with HIV or AIDS, current residence.

d All HIV positive individuals that don't have identified or reported risk factors based on questions on the NM HIV Case Report Form and CDC HIV risk definitions.

^e High risk is defined as sexual contact with a person who injects drugs, bisexual males, or persons with a documented HIV diagnosis.

[†] The 2019 interim report does not include information about death or perinatal exposure that occured during 2019.