



Health Status of New Mexico 2018

December 16, 2017

New Mexico Health Regions




NM Urban-Rural County Classification



- Metropolitan Counties
- Small Metro Counties
- Mixed Urban/Rural Counties
- Rural Counties

Percentage of NM Adults Living in Households with Annual Income of Less Than \$20,000, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Age 65+ Living in Households with Annual Income of Less Than \$20,000, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Living in Households with Annual Income of Less Than \$20,000, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Living in Households with Annual Income of Less Than \$20,000, by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Less Than a High School Education, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Age 65+ with Less Than a High School Education, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Less Than a High School Education, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Less Than a High School Education, by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults without Health Care Coverage, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System


Percentage of NM Adults Age 65+ without Health Care Coverage, By Health Region

New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System


Percentage of NM Adults Age 18-64 without Health Care Coverage, by Health Region, New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults without Health Care Coverage, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults without Health Care Coverage, by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Fair or Poor General Health Status

New Mexico and U.S., 1996-2016




* Addition of Cellular Telephones & New Weighting Process

Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Fair or Poor General Health Status by Household Income


New Mexico, 2016



Source: Centers for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, with New Mexico Department of Health.

Percentage of Adults with Fair or Poor General Health Status by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults in Fair or Poor General Health Status, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Fair or Poor General Health Status by Race/Ethnicity

New Mexico, 2016




** Insufficient Sample Size

Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Fair or Poor General Health Status, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults in Fair or Poor General Health Status by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Current Smoking

New Mexico and U.S., 2001-2016




* Addition of Cellular Telephones & New Weighting Process

Source: NM Behavioral Risk Factor Surveillance System

Current Smoking by Household Income


New Mexico, 2016



Source: Centers for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, with New Mexico Department of Health.

Current Smoking by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Current Smoking, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Current Smoking by Race/Ethnicity


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Current Smoking By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Current Smoking by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Oral Health Visit within Past Year

New Mexico and U.S., 2002-2016 (Even Years Only)




Source: NM Behavioral Risk Factor Surveillance System

* Addition of Cellular Telephones & New Weighting Process

Percentage of Adults with an Oral Health Visit within the Past Year by Annual Household Income


New Mexico, 2016



Source: Centers for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, with New Mexico Department of Health.

Percentage of Adults with an Oral Health Visit within the Past Year by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with an Oral Health Visit within the Past Year, by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with an Oral Health Visit within the Past Year by Race/Ethnicity


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with an Oral Health Visit within the Past Year, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with an Oral Health Visit within the Past Year by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30

New Mexico and U.S., 1995-2016




* Addition of Cellular Telephones & New Weighting Process

Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30 by Household Income


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System Survey

Percentage of NM Adults Who Were Obese BMI \geq 30 by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30 by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30 by Race/Ethnicity


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30, By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults Who Were Obese BMI \geq 30, by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days


New Mexico and U.S., 2001-2016



Definition of Binge Drinking
 Males: 5+ drinks on an occasion, at least once in the past 30 days.
 Females: 4+ drinks on an occasion, at least once in the past 30 days.


Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days by Gender and Household Income

New Mexico, 2016




Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days by Gender and Educational Attainment

New Mexico, 2016



Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days by Gender and Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days by Gender and Race/Ethnicity

New Mexico, 2016




** Insufficient Sample Size

Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days By Gender and Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Binge Drinking at Least Once in Past 30 Days by Gender and Urban/Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes

New Mexico and U.S., 1995-2016




* Addition of Cellular Telephones & New Weighting Process

Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes by Household Income


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes by Race/Ethnicity


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults With Diagnosed Diabetes By Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of NM Adults with Diagnosed Diabetes by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress

New Mexico and U.S., 2003-2016




* Addition of Cellular Telephones & New Weighting Process

Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress by Household Income


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress by Educational Attainment


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress by Age Group


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System


Percentage of Adults with Mental Distress by Race/Ethnicity

New Mexico, 2016



Percentage of Adults with Mental Distress by Health Region


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress by Urban and Rural Counties


New Mexico, 2016



Source: NM Behavioral Risk Factor Surveillance System

Percentage of Adults with Mental Distress by Number of Chronic Health Conditions


New Mexico, 2016



Chronic Health Conditions include: history of MI, history of stroke, heart disease, asthma, COPD, cancer, chronic kidney disease, arthritis, or diabetes.

Alcohol and Drug Use, Grades 9 – 12

New Mexico and U.S., 2015




* Statistically significant difference


Source: NM Youth Risk and Resiliency Survey (YRRS).

Current Drinking by Health Region, Grades 9-12


New Mexico, 2015




Current Drinking by Urban and Rural Status, Grades 9-12 New Mexico, 2015



Binge Drinking by Health Region, Grades 9-12 New Mexico, 2015




Binge Drinking by Urban and Rural Status, Grades 9-12 New Mexico, 2015




Current Cigarette Smoking Grades 9-12

New Mexico and U.S., 2005 - 2015



Current Cigarette Smoking by Health Region, Grades 9-12


New Mexico, 2015



Any Tobacco Use*


by Health Region, Grades 9-12

New Mexico, 2015




Any Tobacco Use Grades 9-12


New Mexico, 2005 - 2015




Current Marijuana Use by Health Region, Grades 9-12 New Mexico, 2015




Current Marijuana Use by Urban and Rural Status, Grades 9-12 New Mexico, 2015



Current Cocaine Use by Health Region, Grades 9-12 New Mexico, 2015




Current Cocaine Use by Urban and Rural Counties, Grades 9-12 New Mexico, 2015




Suicide Attempts and Suicidal Ideation by Gender, Grades 9 – 12


New Mexico and U.S., 2015



Suicide Attempts by Health Region and Gender, Grades 9-12 New Mexico, 2015




Suicide Attempts by Urban and Rural Counties, Grades 9-12 New Mexico, 2015




Forced Sexual Intercourse Grades 9-12


New Mexico and U.S., 2005 - 2015




Forced Sexual Intercourse by Health Region, Grades 9-12 New Mexico, 2015




Forced Sexual Intercourse by Urban and Rural Counties, Grades 9-12 New Mexico, 2015



Selected Risk Behaviors by Sexual Orientation, Grades 9-12 New Mexico, 2015




Selected Risk Behaviors by Sexual Identity, Grades 9-12 New Mexico, 2015





Selected Risk Behaviors by Country of Birth, Grades 9-12

New Mexico, 2015




Selected Risk Behaviors by Family Relationships, Grades 9-12 New Mexico, 2015

In my home, a parent or other adult is interested in my school work.




Academic Success by Selected Risk Behaviors, Grades 9-12

New Mexico, 2015




Academic Success by Parent's Education, Grades 9-12

New Mexico, 2015




Used Painkillers to Get High by Parent's Education, Grades 9-12

New Mexico, 2015




Unstable Housing by Health Region, Grades 9-12

New Mexico, 2015




Unstable Housing by Urban and Rural Counties, Grades 9-12

New Mexico, 2015




Selected Risk Behaviors by Housing Stability, Grades 9-12

New Mexico, 2015



Pregnancy Intention among Women with Live Births


NM, 2008-2015



* 2012 – 2016 were measured differently from preceding years, since it included a third option, 'did not know'
Source: NM Pregnancy Risk Assessment Monitoring System, NMDOH

Preconception Contraceptive Use among Women with Live Births*

NM, 2008-2015




*Among women not trying to get pregnant

Source: NM Pregnancy Risk Assessment Monitoring System, NMDOH

Daily Preconception Multivitamin/Folic Acid Use among Women with Live Births


NM 2008-2015



Source: NM Pregnancy Risk Assessment Monitoring System, NMDOH

Diabetes among Women with Live Births


NM, 2009-2015



Source: NM Pregnancy Risk Assessment Monitoring System, NMDOH


Breastfeeding Initiation and Early Duration among Women with Live Births

NM, 2008-2015



Healthcare Provider Influenza Vaccination Rates


New Mexico, 2012 - 2017



Source: NMDOH Healthcare-associated Infections Program – 33 inpatient healthcare facilities in New Mexico; reporting is voluntary. Healthy People 2020 goal is 90%.

Measles


New Mexico, 1985-2016



Includes confirmed and probable cases, Source: NMDOH Infectious Disease Epidemiology Bureau.

Invasive Bacterial Infection Rates by Pathogen


New Mexico (2012-2016) and U.S (2011-2015)



Sources: New Mexico – NMDOH Infectious Disease Epidemiology Bureau, Active Bacterial Core Surveillance Program (ABCs); US Estimate – Centers for Disease Control and Prevention, Active Bacterial Core Surveillance Report, Emerging Infections Program Network.

Burden of Invasive Bacterial* Disease by Race/Ethnicity†


New Mexico, 2011–2016



**Streptococcus pneumoniae* (SPN), *Haemophilus influenzae* (HFLU), *Neisseria meningitidis* (NMEN), group A *Streptococcus* (GAS) and group B *Streptococcus* (GBS)
 †10% of cases missing race/ethnicity information;
 Source: New Mexico – NMDOH Infectious Disease Epidemiology Bureau, Active Bacterial Core Surveillance Program (ABCs) 2010-2014


Invasive *Streptococcus pneumoniae* Rates

New Mexico and U.S., 2011–2016



Sources: New Mexico – Sources: New Mexico – NMDOH Infectious Disease Epidemiology Bureau, Active Bacterial Core Surveillance Program (ABCs); New Mexico Indicator-Based Information System (NM-IBIS); US Estimate – Centers for Disease Control and Prevention, Active Bacterial Core Surveillance Report, Emerging Infections Program Network, *Streptococcus pneumoniae*, 2010-2014.


Invasive *Streptococcus pneumoniae* Rates by Urban-Rural New Mexico, 2011–2016



Sources: New Mexico – NMDOH Infectious Disease Epidemiology Bureau, Active Bacterial Core Surveillance Program (ABCs); New Mexico Indicator-Based Information System (NM-IBIS)

Hepatitis A Rates


New Mexico and United States, 1995-2016



Includes confirmed and probable cases, Sources: New Mexico – NMDOH Infectious Disease Epidemiology Bureau; United States – Centers for Disease Control & Prevention, Summary of Notifiable Diseases, United States (2014 data not available)

Pertussis Incidence


New Mexico and U.S., 2000-2016



Source: NM Electronic Disease Surveillance System (confirmed and probable cases) and United States – Centers for Disease Control & Prevention, Summary of Notifiable Diseases, United States.


Pertussis Rates in Infants <6 Months of Age

New Mexico, 2006-2016




Influenza-related Hospitalization Rates by Age

New Mexico, 2011-2016



Pneumonia and Influenza-related Death Rates by Year

New Mexico, 2012-2016*




*Data Source: NM Bureau of Vital Records & Statistics 2016-2017
October 1, 2016-April 30, 2017

Pneumonia and Influenza


Death Rates by Age

New Mexico, 2016-2017* Season




Influenza-like Illness (ILI) Visits

New Mexico* and United States**, 2016-2017 Season



Pneumonia and Influenza Death Rates


New Mexico, 2002-2016 and U.S., 2002-2015



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics; CDC WONDER

Pneumonia and Influenza Death Rates by Race/Ethnicity

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Pneumonia and Influenza Death Rates by Health Region


New Mexico, 2016



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics

Pneumonia and Influenza Death Rates by Urban and Rural Counties


New Mexico, 2014-2016



Human Plague Cases by State United States, 1970 - 2016


CDC confirmed cases

N = 478




Human Plague Cases

New Mexico, 1949-2016



Acute Hepatitis B Rates


New Mexico and US, 2004 – 2016



*New Mexico rates are based on less than 20 cases per year. Rates based on fewer than 20 events can be unstable and should be interpreted with caution. Includes confirmed cases only. Sources: New Mexico – NMDOH Infectious Disease Epidemiology Bureau; United States – Centers for Disease Control & Prevention, Division of Viral Hepatitis.

Hepatitis C Hospitalizations*

New Mexico, 2011-2016



*Data from HIDD using ICD9 & ICD 10 codes Acute Hepatitis C: (ICD-9-CM) 070.41, 070.51 – (ICD-10-CM) B17.1, B17.10, B17.11 Chronic Hepatitis C: (ICD-9-CM) 070.44, 070.54 – (ICD-10-CM) B18.2 Unspecified Hepatitis C: (ICD-9-CM) 070.7, 070.70, 070.71 – (ICD-10-CM) B19.2, B19.20, B19.21. Please note that the hepatitis C total hospitalizations does not add up to the total of Acute, Unspecified, and Chronic as there is the potential for a hospitalization to have more than one diagnosis code for these three types of hepatitis C. Non-federal hospitalizations for New Mexico residents only.

Reported Acute & Chronic Hepatitis C Cases

< 30 years old

New Mexico, 2011 - 2016




Source: New Mexico – NMDOH Infectious Disease Epidemiology Bureau. Includes probable and confirmed acute and chronic infections.

Limitations: Data does not differentiate between past and current infection. Reported cases includes only cases tested, most likely this underestimates true burden of disease. Age represents age at diagnosis, which is likely to be older than age of infection.


TB Rates

New Mexico and US, 2012 - 2016




TB Cases by Age Group

New Mexico, 2016



TB Rates Among US Born vs Foreign Born Persons


New Mexico, 2012 - 2016



	2012	2013	2014	2015	2016
US Born Rate	1.1	1.0	1.1	1.2	0.7
Foreign Born Rate	9	15.1	13.7	12.2	12.7


Chlamydia Rates

New Mexico and US, 1989 - 2016



Chlamydia Rates by Sex


New Mexico, 2000 - 2016



	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
■ NM Male CT Rate	110.9	135.3	158.1	158.5	226.7	220.9	243.7	245.7	228.5	248.5	292.6	297.2	307.5	309.3	300.6	327.4	363.7
◆ NM Female CT Rate	450.5	521.8	623.7	621.9	707.3	643.8	749.1	699.5	685.9	678.2	834.5	789.6	833.2	852.3	786.4	873.7	877.7


Chlamydia Rates by Age Group

New Mexico, 2015 - 2016




Chlamydia Rates by Race/Ethnicity

New Mexico, 2015 - 2016




Gonorrhea Rates

New Mexico and U.S., 1972 - 2016




Gonorrhea Rates by Sex

New Mexico, 2000-2016




Gonorrhea Rates by Age Group

New Mexico, 2015 - 2016



Gonorrhea Rates by Race/Ethnicity


New Mexico, 2015 - 2016



Similar to national level data, gonorrhea rates show health disparities: Gonorrhea disproportionately affects African American, American Indian, and Hispanic populations in New Mexico


Primary and Secondary Syphilis Rates

New Mexico and US, 1970 - 2016




Primary and Secondary Syphilis Cases by Sex

New Mexico, 2000-2016




Primary, Secondary, and Early Latent Syphilis Rates by Age Group, Males

New Mexico, 2015-2016




Primary, Secondary, and Early Latent Syphilis Rates by Race/Ethnicity

New Mexico 2015 – 2016




Percentage of Students Obese by Grade

New Mexico, 2010-2016




Percentage of Students Overweight and Obese by Gender

New Mexico, 2016



Percentage of Kindergarten Students Overweight and Obese by Race/Ethnicity

New Mexico, 2016




Source: New Mexico Statewide Childhood Obesity Surveillance System

*Due to small sample sizes, African American and Asian data have been aggregated to include 2015 and 2016 results. Some estimates may be unstable and comparisons cannot be made across groups.

Percentage of Third Grade Students Overweight and Obese by Race/Ethnicity

New Mexico, 2016




Source: New Mexico Statewide Childhood Obesity Surveillance System

*Due to small sample sizes, African American and Asian data have been aggregated to include 2015 and 2016 results. Some estimates may be unstable and comparisons cannot be made across groups.

Child Asthma Hospitalization Rates

Ages 0-17 Years, New Mexico, 1999-2016 and U.S., 2001-2010




NOTES: Includes asthma as first-listed ICD-9 diagnosis (493-493.92) and ICD-10 codes J45 starting September 2015. This series does not include Indian Health Service data.

Source: New Mexico Department of Health (data via <http://ibis.health.state.nm.us>).
 U.S.: CDC/NCHS, National Hospital Discharge Survey (discontinued in 2011).

Child Asthma Hospitalization Rates by Health Region

Ages 0-17 Years, New Mexico, 2012-2016




Source: New Mexico Department of Health (data via <http://ibis.health.state.nm.us>).

NOTES: Asthma as first-listed ICD-9 diagnosis (493-493.92) from 2012 to September 2015. ICD-10 codes J45 for September 2015 and after. This series does not include Indian Health Service data.

Child Asthma Hospitalization Rates by Urban and Rural Counties


Ages 0-17 Years, New Mexico, 2012-2016



Source: New Mexico Department of Health (data via <http://ibis.health.state.nm.us>).

NOTES: Includes asthma as first-listed ICD-9 diagnosis (493-493.92) until September 2015 and ICD-10 codes J45 starting September 2015. This series does not include Indian Health Service data.

Elevated Blood Lead Levels* in Children under Age 6 Years, New Mexico, 2006-2016



*Elevated blood lead levels: ≥ 10 $\mu\text{g}/\text{dL}$ (prior to 2016) & ≥ 5 $\mu\text{g}/\text{dL}$ (lowered in 2016) by venous test or 2 capillary tests within 12 weeks.

SOURCE: NM Environmental Health Epidemiology Bureau's Environmental Public Health Tracking and Childhood Lead Poisoning Prevention Program


Spina Bifida without Anencephaly, New Mexico, 2000-2016 vs U.S., 2000-2011



SOURCES: NM Environmental Health Epidemiology Bureau's Birth Defects Prevention and Surveillance System; National data: <http://www.cdc.gov/mmwr/pdf/wk/mm6401.pdf>, among programs with prenatal ascertainment. Data not available every year.

NOTE: Spina bifida is a neural tube defect, which can be prevented through a woman's intake of folic acid during reproductive years.


Cleft Lip with or without Cleft Palate, New Mexico, 2011-2016



SOURCE: NM Environmental Health Epidemiology Bureau's Birth Defects Prevention and Surveillance System

Occupational Injury Fatalities Per 100,000 Full-Time Equivalent Workers


New Mexico, 2001-2015 and U.S., 2006-2015



Source: U.S. Bureau of Labor Statistics Census of Fatal Occupational Injury

Ambulatory Care Sensitive Condition Hospitalization Rates by Region

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Enhanced Hospital Inpatient Discharge Data

Top Five Ambulatory Care Sensitive Condition Hospitalization Rates

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: Enhanced NM Hospital Inpatient Discharge Data

Diabetes Hospitalizations by Region

New Mexico 2000-2002 and 2014-2016




* The increase in rates, from earlier to later period, is statistically significant

Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Hospital Inpatient Discharge Data-Diabetes as primary diagnosis

Diabetes Hospitalizations by Urban and Rural Counties

New Mexico 2000-2002 and 2014-2016




* The increase in rates, from earlier to later period, is statistically significant

Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Hospital Inpatient Discharge Data -Diabetes as primary diagnosis

Ambulatory Care Sensitive Condition Emergency Department Visit Rates by Region


New Mexico, 2016



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Emergency Department Data


Top Five Ambulatory Care Sensitive Condition Emergency Department Visit Rates

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Emergency Department Data

DALYs: NM vs. US Age-Standardized Total DALYs 1990-2016




DALYs: NM vs. US Age 70+

Total DALYs 1990-2016




DALYs: Top 10 Conditions Age-Standardized

New Mexico vs. US, 2016




DALYs: Top 10 Conditions Age 70+

New Mexico vs. US, 2016




Teen Birth Rates

New Mexico, 2002-2016, and U.S., 2002-2015




Teen Birth Rates by Health Region

New Mexico, 2016




Teen Birth Rates by Urban and Rural Counties

New Mexico, 2014-2016




Percentage of Births to Single Women

New Mexico 2002-2016 and U.S., 2002-2015




Percentage of Births to Single Women by Health Region

New Mexico, 2016




Life Expectancy at Birth by Sex, New Mexico, 2016




Life Expectancy at Birth by Race/Ethnicity

New Mexico, 2016




Life Expectancy at Birth by Urban and Rural Counties

New Mexico, 2016



Life Expectancy at Birth


New Mexico, 2005-2016 and U.S., 2005-2015



Source: New Mexico Indicator-Based Information system (NM IBIS); and ; CDC


Life Expectancy at Birth by Health Region

New Mexico, 2016




Life Expectancy Trend by Sex

New Mexico, 2002-2016




Life Expectancy Trend by Race/Ethnicity

New Mexico, 2002-2016




Infant Mortality Rates

New Mexico, 1930 – 2016 and U.S., 1930-2014




Infant Mortality Rates

New Mexico, 2002-2016 and U.S., 2002-2014




Infant Mortality Rates by Health Region

New Mexico, 2016




Infant Mortality Rates by Urban and Rural Counties

New Mexico, 2014-2016




Infant Mortality Rates by Race/Ethnicity

New Mexico, 2016




Total Deaths Rates

New Mexico, 2005-2016 and U.S., 2005-2015



Total Death Rates by Health Region

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Total Death Rates by Urban and Rural Counties

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Total Death Rates by Race/Ethnicity and Sex


New Mexico, 2016

Female Male




Total Death Rates by Race/Ethnicity, Ages 0-24 Years

New Mexico, 2016




Total Death Rates by Race/Ethnicity Ages 25-64 Years

New Mexico, 2016



Total Death Rates by Race/Ethnicity Ages 65 and older Years

New Mexico, 2016




Top 8 Leading Causes of Death,

New Mexico, 2016 and U.S., 2015

New Mexico	Rank	United States
Heart disease	1	Heart disease
Cancer	2	Cancer
Unintentional Injuries	3	Chronic lower respiratory diseases
Chronic lower respiratory diseases	4	Accidents (unintentional injuries)
Cerebrovascular diseases	5	Cerebrovascular diseases
Diabetes mellitus	6	Alzheimer's disease
Chronic liver disease and cirrhosis	7	Diabetes mellitus
Alzheimer's disease	8	Influenza and Pneumonia

5 Leading Causes of Death

New Mexico, 2016, and U.S., 2015




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

5 Leading Causes of Death

Northwest Region, New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


5 Leading Causes of Death

Northeast Region, New Mexico, 2016



5 Leading Causes of Death

Metro Region, New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

5 Leading Causes of Death

Southeast Region, New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


5 Leading Causes of Death

Southwest Region, New Mexico, 2016




Heart Disease Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015



Heart Disease Death Rates by Race/Ethnicity

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Heart Disease Death Rates by Health Region

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Heart Disease Death Rates by Urban and Rural Counties

New Mexico, 2014-2016



Cancer Death Rates


New Mexico, 2002-2016 and U.S., 2002-2015



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics; CDC Wonder

Cancer Death Rates by Race/Ethnicity

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Cancer Death Rates by Health Region

New Mexico, 2016



Cancer Death Rates by Urban and Rural Counties

New Mexico, 2014-2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Chronic Lower Respiratory Disease Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015




Chronic Lower Respiratory Disease Death Rates by Health Region

New Mexico, 2016



Chronic Lower Respiratory Disease Death Rates by Urban and Rural Counties

New Mexico, 2014-2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Cerebrovascular Disease Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015



Cerebrovascular Disease Death Rates by Health Region

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Cerebrovascular Disease Death Rates by Urban and Rural Counties

New Mexico, 2014-2016



Diabetes Death Rates


New Mexico, 2002-2016 and U.S., 2002-2015



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics; CDC Wonder


Diabetes Death Rates by Health Region

New Mexico, 2016




Diabetes Death Rates by Urban and Rural Counties

New Mexico, 2014-2016




Injury Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015




Suicide Rates

New Mexico, 2002-2016 and U.S., 2002-2015




Suicide Rates by Age Group

New Mexico, 2016




Suicide Rates by Urban and Rural Counties

New Mexico, 2014-2016




Youth (15-19) Suicide Rates

New Mexico, 2005-2016 and U.S., 2005-2015




Youth (15-19) Suicide Rates by Health Region

New Mexico, 2012-2016




Unintentional Injury Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015



Unintentional Injury Death Rates by Race/Ethnicity

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics


Unintentional Injury Death Rates by Health Region

New Mexico, 2016




Unintentional Injury Death Rates by Urban and Rural Counties

New Mexico, 2014-2016




Fall-Related Death Rates Among Older Adults by Age Group

New Mexico, 2005-2016




Fall-Related Death Rates Among Older Adults 65+ by Race/Ethnicity

New Mexico, 2012-2016




Fall-Related Death Rates Among Older Adults 65+ by Health Region

New Mexico, 2012-2016




Fall-Related Death Rates Among Older Adults 65+ by Urban and Rural Counties

New Mexico, 2012-2016




Firearm Injury Death Rates

New Mexico, 2002-2016 and U.S., 2002-2015



Firearm Injury Death Rates by Race/Ethnicity

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Firearm Injury Death Rates by Health Region


New Mexico, 2016



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics


Firearm Injury Death Rates by Urban and Rural Counties

New Mexico, 2016



Chronic Liver Disease Death Rates


New Mexico, 2002-2016 and U.S., 2002-2015



Rates have been age-adjusted to the standard U.S. 2000 population
Source: NM Vital Records and Health Statistics; CDC WONDER


Chronic Liver Disease Death Rates by Race/Ethnicity

New Mexico, 2012-2016



Chronic Liver Disease Death Rates by Health Region

New Mexico, 2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics

Chronic Liver Disease Death Rates by Urban and Rural Counties

New Mexico, 2016



Smoking-attributable, Alcohol-attributable, and Drug Overdose Death Rates¹

New Mexico (2012-2016) and US (2015)

	New Mexico	U.S.	Rate Ratio
Smoking-attributable death rate ²	96.3	108.1	0.9
Alcohol-attributable death rate ³	59.7	32.2	1.9
Drug overdose death rate ⁴	24.6	16.3	1.5

¹Rates are deaths per 100,000 population, age-adjusted to the 2000 U.S. standard population


²Smoking-Attributable Mortality, Morbidity, and Economic Costs (SAMMEC), apps.nccd.cdc.gov/sammec/index.asp, 2015 for US and 2012-2016 for NM

³Alcohol-Related Disease Impact (ARDI), nccd.cdc.gov/DPH_ARDI/default/default.aspx, 2012-2016 for NM and 2015 for US

⁴Defined as International Classification of Diseases, tenth revision (ICD-10) codes X40-X44, X60-X64, X85, and Y10-Y14, 2012-2016 for NM and 2015 for US

Alcohol-Attributable Death Rates

New Mexico, 1990-2016, and U.S., 1990-2015




Source: NCHS (US); NMDOH BVRHS (NM); CDC ARDI v. 2

Data Sources: NCHS (US); NMDOH BVRHS; UNM-GPS; IBEB SAES; CDC ARDI
Rates are age-adjusted to the 2000 U.S. standard population
2015 for the US, and 2016 for NM

Alcohol-Attributable Death Rates by Region


New Mexico, 2012-2016



Data Sources: NMDOH BVRHS; UNM-GPS; IBEB SAES; CDC ARDI
Rates are age-adjusted to the 2000 U.S. standard population

Alcohol-Attributable Death Rates by Rural/Urban


New Mexico, 2012-2016



Data Sources: NMDOH BVRHS and IBEB SAE; CDC ARDI
Rates are age-adjusted to the 2000 U.S. standard population

Drug Overdose Death Rates

New Mexico, 1990-2016, and U.S., 1990-2015




Rates are age adjusted to the US 2000 standard population
Source: United States (CDC Wonder); New Mexico (NMDOH BVRHS/UNM GPS)

Rates are age-adjusted to the 2000 US Standard Population
ICD-10 codes include X40-X44, X60-X64, X85, and Y10-Y14
Source: CDC Wonder and NM Bureau of Vital Records and Health Statistics

Drug Overdose Death Rates by Health Region

New Mexico, 2012-2016




Rates have been age-adjusted to the standard U.S. 2000 population

Source: NM Vital Records and Health Statistics; UNM-GPS; IBEB SAES; CDC ARDI

Drug Overdose Death Rates by Rural/Urban


New Mexico, 2012-2016



Data Sources: NMDOH BVRHS and IBEB SAE;
Rates are age-adjusted to the 2000 U.S. standard population

Teen Birth Rates by Census Tract Poverty Level*


New Mexico, 2009-2013



*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty.
Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Percentage with Prenatal Care in the First Trimester by Census Tract Poverty Level*


New Mexico, 2011-2013



*Poverty level is the percentage of persons of all ages in the mother's census tract living at or below 100% of Poverty.
Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Percentage of Births that were Pre-Term (less than 37 weeks) by Census Tract Poverty Level*


New Mexico, 2011-2013



*Poverty level is the percentage of persons of all ages in the mother's census tract living at or below 100% of Poverty.
Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Total Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Infant Mortality by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Malignant Neoplasm Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Malignant neoplasms are defined by ICD 10 C00- C97

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Heart Disease Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Heart disease deaths are defined by ICD10: I00-I09, I11, I13, I20-I51

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Unintentional Injury Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Unintentional injury deaths are defined by ICD10: V01-X59, Y85-Y86

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

All Injury Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Injury deaths are defined by (ICD10: X85-Y09, Y87.1, *U01-*U02), Injury, Complications of medical and surgical care (ICD10: Y40-Y84, Y88), Injury, Intentional self-harm (suicide) (ICD10: X60-X84, Y87.0, *U03), Injury, Legal intervention (ICD10: Y35, Y89.0), Injury, Operations of war and their sequelae (ICD10: Y36, Y89.1), Injury, Unintentional injuries (ICD10: V01-X59, Y85-Y86) Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Cerebrovascular Disease Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Cerebrovascular disease deaths are defined by ICD10: I60-I69

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Chronic Lower Respiratory Disease Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Chronic lower respiratory disease deaths are defined by ICD10: J40-J47

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Chronic Liver Disease and Cirrhosis Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Chronic liver disease and cirrhosis deaths are defined by ICD10: K70, K73-K74

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Diabetes Mellitus Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Diabetes mellitus deaths are defined by ICD10: E10-E14

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Alcohol Related Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Alcohol related deaths are defined by ICD10: Alcohol-related deaths include deaths that are partially attributable to alcohol as well as 100% attributable alcohol-induced deaths. For more information, visit the CDC Alcohol-related Disease Impact (ARDI) software page: <http://www.cdc.gov/alcohol/ardi.htm>
Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Drug Overdose Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Drug Overdose deaths are defined by ICD 10: X40-X44, X60-X64, X85, Y10-Y14.

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Suicide Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Suicide deaths are defined by ICD 10: X60-X84, Y87.0, *U03.

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Youth (Ages 15-19) Suicide Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Suicide deaths are defined by ICD 10: X60-X84, Y87.0, *U03.

Rates have been age-adjusted to the standard U.S. 2000 population.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Older (Age 65+) Adult Fall Death Rates by Census Tract Poverty Level*


New Mexico, 2009-2013



*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Older adult fall deaths are defined by ICD10 external cause of injury codes retrieved from http://www.cdc.gov/nchs/data/ice/icd10_transcode.pdf
Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Infectious Disease Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty.

Rates have been age-adjusted to the standard U.S. 2000 population. Infectious Disease Include: Human immunodeficiency virus (HIV) disease (ICD10: B20-B24), Malaria (ICD10: B50-B54), Measles (ICD10: B05), Meningitis (ICD10: G00, G03), Meningococcal infection (ICD10: A39), Respiratory, Bronchitis and bronchiolitis, acute (ICD10: J20-J21), Respiratory, Influenza and pneumonia (ICD10: J09-J18), Scarlet fever and erysipelas (ICD10: A38, A46), Shigellosis and amebiasis (ICD10: A03, A06), Syphilis (ICD10: A50-A53), Tuberculosis (ICD10: A16-A19), Viral hepatitis (ICD10: B15-B19), Whooping cough (ICD10: A37)

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Septicemia Death Rates by Poverty Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Rates have been age-adjusted to the standard U.S. 2000 population.

Septicemia deaths are defined by ICD10 codes: A40-A41

Influenza and Pneumonia Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Rates have been age-adjusted to the standard U.S. 2000 population.

Influenza and Pneumonia deaths are defined by ICD10 codes: J09-J18

Firearm Injury Death Rates by Census Tract Poverty Level*

New Mexico, 2009-2013




*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty. Rates have been age-adjusted to the standard U.S. 2000 population.

Firearm deaths are defined by ICD10 external cause of injury codes retrieved from http://www.cdc.gov/nchs/data/ice/icd10_transcode.pdf
Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Life Expectancy from Birth by Census Tract Poverty Level*

New Mexico, 2009-2013



*Poverty level is the percentage of persons of all ages in the decedent's census tract living at or below 100% of Poverty.


Rates have been age-adjusted to the standard U.S. 2000 population.

Life expectancy calculated by methods retrieved from: <http://ibis.health.state.nm.us/resources/LifeExp.html>.

Source: NM Vital Records and Health Statistics, U.S. Bureau of the Census, American Community Survey

Years of Potential Life Lost (YPLL) Before Age 75

New Mexico, 2016



Conclusions

- Health status is strongly inversely associated with poverty at the community level
- Exceptions to this include suicide, older adult falls deaths, and Alzheimer's disease deaths
- Indicators with greater than twice the risk for persons living in high poverty areas include: teen birth, deaths due to chronic liver disease, diabetes, alcohol, infectious disease, and influenza and pneumonia

Conclusions

- In general, chronic disease mortality rates continue to decrease in New Mexico
- In general, injury and substance abuse death rates are stable or increasing
- Vaccination continues to reduce infectious disease rates and disparities
- The infant mortality rate reduction in NM is a major success but disparities remain
- Geographic, particularly unfavorable to rural areas, socioeconomic and racial/ethnic disparities persist in NM

NM Health Status Resources

- NM IBIS (Indicator Based Information System) – <http://ibis.health.state.nm.us>
- NM Environmental Public Health Tracking System – <https://nmtracking.unm.edu>
- NM DOH health status presentations - <https://nmhealth.org/about/erd/hdps/>
- NM DOH health data resources and publications - <https://nmhealth.org/data/>