

HIV

Surveillance and epidemiology Program

HIV Infection and AIDS among Adults and Adolescents, New Mexico 2014

	Newly Diagnosed in 2014*			Persons Living with HIV or AIDS in New Mexico								
	N	%	Rate†	HIV			AIDS			Total		
	N	%	Rate†	N	%	Rate†	N	%	Rate†	N	%	Rate†
Total	132	100.0	7.7	1375	100.0	80.0	1952	100.0	113.4	3327	100	193.3
Sex at birth												
Male	117	88.6	13.9	1175	85.4	139.2	1740	89.1	206.1	2915	87.6	345.3
Female	15	11.4	1.7	200	14.6	22.8	212	10.9	24.2	412	12.4	47.0
Race/Ethnicity												
American Indian/Alaska Native	28*	21.2	19.6	124	9.0	86.7	154	7.9	107.9	278	8.4	194.5
Asian/Pacific Islander	0	0.0	0.0	11	0.8	40.7	13	0.7	48.6	24	0.7	88.7
Black	8	6.1	22.8	108	7.9	308.5	105	5.4	302.6	213	6.4	608.5
Hispanic	66	50.0	8.7	651	47.3	86.2	850	43.6	112.7	1501	45.1	198.7
White	30	22.7	3.9	464	33.7	61.0	794	40.6	104.4	1258	37.8	165.4
Multi-race	0	0.0	0.0	17	1.2	CNC	36	1.8	CNC	53	1.6	CNC
Age‡												
13-24	23	17.4	6.5	65	4.7	18.5	15	0.8	4.3	80	2.4	22.8
25-34	49	37.1	18.1	318	23.3	117.5	134	6.8	49.5	452	13.6	167.0
35-44	27	20.5	10.8	336	24.4	134.2	341	17.5	136.2	677	20.4	270.4
45-54	22	16.7	7.6	371	27.0	128.6	761	39.0	263.9	1132	34.0	392.5
55+	11	8.3	2.0	285	20.6	50.9	701	36.0	125.1	986	29.6	175.9
Region§												
Northwest	26*	19.7	14.4	134	9.7	74.1	167	8.6	92.4	301	9.1	166.5
Northeast	17†	12.9	6.8	249	18.1	99.4	395	20.2	157.7	644	19.4	257.1
Metro	67	50.8	9.0	657	47.8	87.8	1004	51.3	134.2	1661	49.9	222.1
Southeast	7	5.3	3.0	74	5.4	31.5	120	6.2	51.2	194	5.8	82.7
Southwest	15	11.4	4.9	261	19.0	84.9	266	13.6	86.5	527	15.8	171.4
Transmission Category												
Men who have sex with men (MSM)	82	62.1	CNC	820	59.6	CNC	1226	62.9	CNC	2046	61.5	CNC
Intravenous Drug Use (IDU)	11	8.3	CNC	105	7.6	CNC	173	8.9	CNC	278	8.4	CNC
MSM/IDU	4	3.0	CNC	94	6.8	CNC	194	9.9	CNC	288	8.7	CNC
High-Risk-Hetrosexual	18	13.6	CNC	165	12.0	CNC	198	10.1	CNC	363	10.9	CNC
Other	0	0.0	CNC	2	0.1	CNC	11	0.6	CNC	13	0.4	CNC
Perinatal Exposure	0	0.0	CNC	5	0.4	CNC	8	0.4	CNC	13	0.4	CNC
No Risk Identified	17	12.9	CNC	184	13.4	CNC	142	7.3	CNC	326	9.8	CNC

Footnotes

NOTE: Distribution of prevalent HIV and AIDS cases amongst the total Persons Living with HIV or AIDS in New Mexico is 41.3% and 58.7%, respectively.

NOTE: CNC means Can Not Calculate - population data is unavailable.

* All people who were initially diagnosed while residing in New Mexico during 2013 - age 13 and older

† Rates per 100,000 based on Bureau of Business and Economic Research Geospatial and Population Studies Program population estimates for 2013.

‡ For new diagnoses, age at time of diagnosis. For persons living with HIV or AIDS, current age.

§ For new diagnoses, residence at time of diagnosis. For persons living with HIV or AIDS, current residence.

Cautions

N < 20: Rate calculations for small numbers may be statistically unreliable. Fluctuations in the associated rates may be due in part to statistical unreliability rather than true change.

Region: New Mexico Department of Health constructed new health regions in 2012. Regional comparison to previous years is not recommended. A map depicting New Mexico's Health Regions is available for download from the New Mexico Department of Health at <http://ibis.health.state.nm.us/image/HealthRegions2012.png>.

‡ - Indicates greater than 20% decrease from previous year for values > 20.

* - Indicates greater than 20% increase from previous year for values > 20.