

**NEW MEXICO
SELECTED HEALTH STATISTICS
ANNUAL REPORT
VOLUME 2**

**Mortality and Demographic/Socioeconomic Status
2007**



THE STATE CENTER FOR HEALTH STATISTICS
Bureau of Vital Records and Health Statistics

State of New Mexico
Bill Richardson, Governor

New Mexico
Department of Health

Alfredo Vigil, MD, Secretary

Karen Armitage, MD, Chief Medical Officer

Epidemiology and Response Division

C. Mack Sewell, DrPH, MS, Director and State
Epidemiologist

Michael G. Landen, MD, MPH, Deputy State Epidemiologist

The State Center for Health Statistics
Bureau of Vital Records and
Health Statistics

Larry Nielsen, Bureau Chief, and Director,
Designated State Center for Health Statistics

Mary Anne Escudero, Vital Records Manager, State Registrar

Renee Valencia, Registration Manager

Publication date: May 2010

Photos: Jim Orr (jamesorr.smugmug.com); Jon Branch, New Mexico Bureau of Vital Records and Health Statistics.

ACKNOWLEDGEMENTS

The *New Mexico Selected Health Statistics Annual Report, Volume 2, 2007* is produced by the designated State Center for Health Statistics within the Bureau of Vital Records and Health Statistics. The Epidemiology and Statistics Unit is responsible for analyzing data from the birth and death certificates and reports of fetal death and induced termination of pregnancy. Additionally, staff members are responsible for producing statistics, health surveillance reports and data files for researchers, public health professionals, other state and federal agencies and the public.

This report was prepared under the direction of Larry Nielsen, and would not be possible without the teamwork of Camille Clifford, Patricia Drypolcher, Irma Montoya, Larry Nielsen, Micah Nussbaum, Tony Ortiz, Rosemary Perez-Mendoza, and Anne Worthington.

The work of the Epidemiology and Statistics unit is dependent upon the efforts of the Records and Registration units, which register and issue certified copies of birth and death certificates. The collaboration of these units is essential for data quality and timeliness. The State Office is supported at the local level through the efforts of 26 Local Registrars from across the state. The Information Technology Services Division provides computer system support.

The Bureau is grateful to the Epidemiology and Response Division, the administration of the Department of Health, and various sections of the department for their support and assistance. Acknowledgement is extended to other state and federal agencies whose data aided in the preparation of this report. Among these agencies are: University of New Mexico's Bureau of Business and Economic Research (BBER), Bureau of the Census and the Centers for Disease Control's National Center for Health Statistics (NCHS). Special thanks are also extended to the providers (physicians, medical records staff, midwives, funeral directors, the Office of the Medical Investigator, tribal enrollment officers and others) who have made this publication possible through their conscientious data reporting efforts.

Mission Statement

The mission of the New Mexico Department of Health is to promote health and sound healthy policy, prevent diseases and disability, improve health services systems and assure that essential public health functions and safety net services are available to New Mexicans.

Executive Summary

The New Mexico Selected Health Statistics Annual Report has been produced since 1974. The 2007 report introduces a major change in that it has been divided into two volumes. This was done because of long delays in collecting data on deaths that occurred in some other states for 2007.

Volume 1 focused on natality. It also included data on fetal deaths and abortions.

Volume 2 focuses on mortality. It also includes a section on demographics and socioeconomic status.

2007 is also notable in that the Bureau of Business and Economic Research (BBER) at the University of New Mexico, the source of population data for the New Mexico Department of Health, adjusted its methodology for estimating intercensal populations. This adjustment has had a significant impact on age-adjusted, and some age-specific, mortality rates. See the Technical Appendix for additional information.

Demographic and Socioeconomic Highlights

New Mexico's estimated 2007 population reached 2,053,922, reflecting an increase of 12.5% since 2000. The state's population is projected to reach 2,865,000 by 2030.

The adjustments made by BBER in 2007 to its methodology for estimating population resulted in a change in how the population was distributed, especially among age groups. In 2006, 29,933 persons (1.5%) were age 85 and over. In 2007 the estimate for this age group dropped nearly in half to 15,718 (0.8%) while the estimate for ages 75-84 increased to 102,034 (5.0%), compared to 2006 when that age group was estimated at 82,916 (4.1%).

For race/ethnicity in 2007, 43.4% of New Mexicans were White (non-Hispanic), 41.2% were Hispanic (White), 11.0% were American Indian or Alaska Native, 2.7% were Black or African American and 1.7% were Asian or Pacific Islander.

Mortality Highlights

In 2007, there were 15,400 deaths among New Mexico residents, translating to an age-adjusted death rate of 902.2 deaths per 100,000. The state's 2007 death rate was higher than the 2006 national rate of 776.5. However, due to the changes in BBER's methodology for estimating population, it is also considerably higher than the 2006 New Mexico rate of 761.2. In number, there were only 169 or 1.1% more deaths among New Mexican residents in 2007 compared to the total of 15,231 in 2006.

The infant mortality rate was slightly higher in 2007 (6.1) than in 2006 (5.7). It is lower than the 2006 U.S. rate of 6.7. Of the 188 deaths of New Mexico infants under 1 year-old in 2007, 40 were caused by disorders related to length of gestation or birthweight, 39 were caused by congenital malformations (birth defects), and 21 were caused by maternal factors and complications of pregnancy or labor and delivery. There were 14 deaths attributed to sudden infant death syndrome (SIDS).

In 2007, the leading cause of death among New Mexico residents was Diseases of the Heart (3,263 deaths), with Malignant Neoplasms (cancer) next (3,210 deaths). Accidents accounted for 1,277 deaths.

Contact Information

Comments on the content and format of this report or requests for additional data or analyses should be directed to:

The State Center for Health Statistics
at the Bureau of Vital Records and Health Statistics
Epidemiology and Response Division
New Mexico Department of Health
1105 St. Francis Drive
P.O. Box 26110
Santa Fe, New Mexico 87502-6110
(866) 534-0051

Suggested Citation:

New Mexico Selected Health Statistics Annual Report, Volume 2, 2007.
Santa Fe, New Mexico: New Mexico Department of Health,
Bureau of Vital Records and Health Statistics. 2010.

Table of Contents

Demographic and Socioeconomic Status Section

Population and Components.....	12
Population Distribution.....	13
Age.....	13
Race and Ethnicity.....	13
Birth to Death Comparison.....	15
Economic Factors.....	15
Income	15
Poverty	17
Trends	17

Mortality Section

All Causes.....	32
Leading Causes of Death.....	34
Selected Causes.....	35
Injury Deaths and Alcohol- and Drug-Induced Deaths	35
Maternal Mortality.....	37
Infant Mortality.....	37
Disposition of Remains.....	39

Technical Appendix.....	69
--------------------------------	-----------

List of Tables and Figures

Demographic and Socioeconomic Status Section

Figure D-1	New Mexico Population: Counts and Projections	12
Figure D-2	Population Distribution by Age, New Mexico and United States	13
Figure D-3	Population Pyramids, New Mexico and United States	14
Figure D-4	New Mexico Population Distribution by Race/Ethnicity	15
Figure D-5	New Mexico Population Distribution by Race/Ethnicity, Age and Sex	16
Figure D-6	Per Capita Personal Income Ranked by County, New Mexico	16
Figure D-7	Children Under 18 at or Below Poverty Level, Ranked by County, New Mexico	17
Table D-1	Population by Age and County, New Mexico	18
Table D-2	Male Population by Age and County, New Mexico	20
Table D-3	Female Population by Age and County, New Mexico	22
Table D-4	Population by Race/Ethnicity, Sex and Age, New Mexico	24
Table D-5	Population by Race/Ethnicity, County and Health Region, New Mexico	26
Table D-6	Components of Population Change, New Mexico	27
Table D-7	Percent Distribution of New Mexico Population by Age Group	27
Table D-8	Population Estimates, Components of Change, New Mexico	28
Table D-9	Population, Births and Deaths by City, New Mexico	29
Table D-10	Summary of Trends in New Mexico Health Statistics	30

Mortality Section

Figure M-1	Percent of Deaths by Age Group and Sex, New Mexico	32
Figure M-2	Death Rates by Sex and Race/Ethnicity, New Mexico	33
Figure M-3	Death Rates, New Mexico and United States	33
Figure M-4	Leading Causes of Death, New Mexico and United States	34
Figure M-5	Death Rates by Cause, New Mexico	35
Figure M-6	Percent of Injury Deaths by Type, New Mexico	36
Figure M-7	Percent of Firearm Injury Deaths by Manner, New Mexico	36
Figure M-8	Infant Mortality Rates, New Mexico and United States	37
Figure M-9	Infant Mortality Rates by Race/Ethnicity, New Mexico	38
Figure M-10	Infant Mortality Rates, New Mexico and United States	38
Figure M-11	Percent Distribution by Method of Disposition, New Mexico	39
Table M-1	Numbers and Rates of Deaths by Age, Race/Ethnicity and Sex, New Mexico and United States	40
Table M-2	Number of Deaths by Age, Sex, County and Health Region, New Mexico and United States	42
Table M-3	Death Rates by Age, Sex, County and Health Region, New Mexico and United States	45
Table M-4	Numbers and Rates Ranked by Leading Causes of Death by County and Health Region New Mexico and United States - Both Sexes	48
Table M-5	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, All Ages New Mexico	50
Table M-6	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 1 to 4 New Mexico	50
Table M-7	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 5 to 14 New Mexico	51

List of Tables and Figures (continued)

Mortality Section (continued)

Table M-8	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 15 to 24 New Mexico	51
Table M-9	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 25 to 44 New Mexico	52
Table M-10	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 45 to 64 New Mexico	52
Table M-11	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 65 to 84 New Mexico	53
Table M-12	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, Ages 85 Plus New Mexico	53
Table M-13	Numbers and Rates of Deaths by Race/Ethnicity and Sex, New Mexico and United States	54
Table M-14	Rates for Leading Causes of Death, New Mexico	55
Table M-15	Rates and Numbers of Deaths for 39 Selected Causes by Age, New Mexico	56
Table M-16	Rates and Numbers of Deaths for 39 Selected Causes by Sex, New Mexico	57
Table M-17	Rates and Numbers of Deaths for 39 Selected Causes by Health Region, New Mexico	58
Table M-18	Death Rates for 39 Selected Causes by Race/Ethnicity, New Mexico	59
Table M-19	Number of Injury Deaths by Sex, County and Health Region, New Mexico and United States	60
Table M-20	Death Rates for Injury by Sex, County and Health Region, New Mexico and United States	61
Table M-21	Number of Deaths due to Firearms, Alcohol, Drugs and Injury at Work by Sex, County and Health Region, New Mexico and United States	62
Table M-22	Death Rates due to Firearms, Alcohol or Drugs, and Injury at Work by Sex, County and Health Region, New Mexico and United States	63
Table M-23	Numbers and Rates of Maternal Mortality by Age, New Mexico	64
Table M-24	Numbers and Average Rates of Maternal Mortality by Cause, New Mexico	65
Table M-25	Numbers and Rates of Infant Mortality by Sex and Race/Ethnicity, New Mexico	66
Table M-26	Numbers of Infant Deaths by Age and Cause, New Mexico and United States	67
Table M-27	Method of Final Disposition for Deaths Occurring in New Mexico	68

Demographic and Socioeconomic Status Section



DEMOGRAPHIC AND SOCIOECONOMIC STATUS SECTION

POPULATION AND COMPONENTS

The total United States population for 2007 was 301,621,157 (U.S. Census Bureau, released December 2007). This total represents a 0.7% increase in the nation's population since 2006. New Mexico's 2007 population was 2,053,922, a 2.1% increase from 2006. This is also an increase of 12.5% since 2000.

The Bureau of Business and Economic Research (BBER) at the University of New Mexico is the Department of Health's source for population data. Beginning with the 2007 population estimates, BBER adjusted its methodology for estimating intercensal populations. Please see the *Technical Appendix* for a brief discussion about this estimation change.

BBER's methodology change resulted in a change in how the population was distributed, especially among age groups. In 2006, 29,333 persons (1.5%) were age 85 and over. In 2007, the estimate for this age group dropped nearly in half to 15,718 (0.8%) while the estimate for ages 75-84 increased to 102,034 (5.0%), compared to 2006 when that age groups was estimated at 82,916 (4.1%).

**Figure D-1
New Mexico Population: Counts and Projections, 1850 to 2030**




Table D-6 lists the components of New Mexico's population change based on 10-year time periods and from 2000 to 2007. For the latter period, the state's population increased by 228,034 persons (12.5%).

From 1950-1990, natural increase exceeded net migration. During 1990-2000, these two components were almost identical. New Mexico experienced its highest percentage of growth in the 1950's due to a high rate of natural increase coupled with a high rate of in-migration. During the 1960's, the state's growth continued more slowly due to migration out of the state and a lower rate of natural increase. The 1970's were years of renewed growth for the state, including high in-migration. Steady population increases have occurred from 1980 to 2007 (Table D-6).

New Mexico's population is projected to reach 2,540,145 in 2020 and 2,864,796 in 2030. These are projected increases of 23.7% and 39.5%, respectively. (Figure D-1)

Sandoval County was the fastest growing county in the state for 2007, increasing by 7.6% between 2006 and 2007 (Table D-8). Valencia County, the second fastest growing county, had an increase of 5.5%. Dona Ana was third, with a 3.3% increase in population.

Table D-8 also details the 2006 to 2007 components of change at the national, state and county levels. Both the United States and New Mexico increased in population during this time period with increases of 0.7% and 2.1%, respectively. From 2000 to 2007, net migration exceeded natural increase in New Mexico by 10.1%.

Between 2000 and 2007, the counties with the largest natural increases were Bernalillo, Dona Ana, San Juan, McKinley and Santa Fe counties. Bernalillo, Sandoval, Dona Ana, Santa Fe and Valencia counties experienced the largest influx of people moving into the state. Eight New Mexico counties experienced a movement of people out of their areas between 2000 and 2007. McKinley County lost the largest number of inhabitants: 2,069 persons.

The percent contribution to the state's population increase by county for 2006 to 2007 is in Table D-8. Of the state's estimated population increase of 42,517 persons during 2007, Bernalillo County accounted for over 37% of that growth, while Sandoval County contributed 20.6%. Bernalillo, Sandoval and Dona Ana counties made up 73.4% of the state's growth from 2006 to 2007. Only one county, Torrance, decreased in population during the same time period.

POPULATION DENSITY

Table D-8 lists New Mexico's 2007 population densities by county and health region. The county densities ranged from a high of 552.3 persons per square mile in Bernalillo County to a low of 0.4 persons per square mile in Harding County. Eight of the state's counties exceeded the state population density of 16.9 persons per square mile. After Bernalillo County, Los Alamos County was the second most densely populated, followed by Santa Fe County. Twenty-five counties had lower population densities than that of the state.

POPULATION DISTRIBUTION

Age

For both the United States and New Mexico, the largest proportion of the 2007 population was between the ages of 20 and 44 (Figure D-2). The second largest proportion was between the ages of 45 and 64.

Figure D-2
Population Distribution by Age
New Mexico and United States, 2007




Figure D-3 illustrates changes in the age structure of New Mexico and the United States during the twentieth century. In 1900, both the United States and New Mexico were characterized by low life expectancy and high fertility rates resulting in a high proportion of young people and a low proportion of elderly. After World War II, the proportion of the population in the younger age groups declined to produce a more uniform pyramid. At the same time, the national figure becomes more rectangular in shape, indicating a more even distribution of ages in the population. New Mexico's older population was slower to increase than that of the United States. By 2000, the state population distribution more closely reflected that of the nation

Table D-1 shows the number and percent of the 2007 population in New Mexico by age and resident county. The largest proportion of New Mexico's population was between 45 and 54 (15.3%), followed by ages 5 to 14 (13.6%). Over 30% of the population of Bernalillo County was between the ages of 35 and 54. McKinley, Torrance and Valencia counties had much younger populations, with the largest proportions in ages 5 to 14. In McKinley County, only 6.7% of the population was 65 and older. Harding and Sierra counties had the oldest populations in the state, with approximately 33.2% and 27.5% of their populations over age 65, respectively.

Tables D-2 and D-3 show population distributions by sex and county in New Mexico. The same trends demonstrated in the entire population (Table D-1) were seen in the male and female groups. In all counties, there were more males than females in the younger age groups. However, starting at age 35 there were more females than males.







Table D-7, which contains the percent distribution of New Mexico's population by age groups, has been updated to include projections for 2020 and 2030. The percent population changes from 1990 to 2030 indicate that New Mexico's older populations will be increasing while the younger age groups will be decreasing. Specifically, the 65 plus age group may increase by over 80% while the under 5 years to under 35 years age groups may be decreasing from 13.0% to almost 35%.

Race and Ethnicity

The New Mexico Department of Health combines race and ethnicity for reporting purposes.

According to 2007 state population estimates, 43.4% of New Mexicans were White and 41.2% were Hispanic


Figure D-3
Population Pyramids
New Mexico and United States, 1900, 1950, and 2000



(Figure D-4). (The Hispanic category does not include Black, American Indian or Asian or Pacific Islander populations.) The American Indian or Alaska Native population comprised 11.0% of New Mexico's population; the Black or African American population made up 2.7%; and the Asian or Pacific Islander population constituted 1.7%.

The percentage of the population that was Hispanic was largest in Guadalupe, Mora and San Miguel counties (Table D-5). McKinley, Cibola and San Juan counties had the largest proportions of American Indian or Alaska Native residents. American Indian or Alaska Native residents comprised 78.5% of McKinley County and just under 42% of Cibola and San Juan counties. Curry County had the largest percentage of Black or African American residents, and Los Alamos County had the largest percentage of Asian or Pacific Islander residents.

Figure D-4
New Mexico Population Distribution by Race/Ethnicity
New Mexico, 2007



AI or AN = American Indian or Alaska Native, API = Asian or Pacific Islander, AA=African American.
 Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. Released 2009.
<http://www.unm.edu/~bber/>.

Figure D-5 illustrates the 2007 proportion of age groups, sexes, and race/ethnicity categories in New Mexico. The American Indian or Alaska Native, Black or

African American and Hispanic populations had higher proportions of individuals under age 25. The Asian or Pacific Islander and Black or African American populations also had higher proportions of 25-54 year-olds, while the White population had the highest proportion of those 55 and older (Table D-4).

Birth to Death Comparison

There were 2.0 births for every death in New Mexico in 2007 (Table D-8). Six counties exceeded this ratio: McKinley, Dona Ana, Lea, San Juan, Curry and Roosevelt. There were 2.8 births for every death in McKinley County, the county with the highest birth-to-death ratio. The birth-to-death ratio for 27 counties fell below the state ratio. In six of these counties, the number of deaths exceeded the number of births.

The number of births and deaths are shown for 98 New Mexico cities in Table D-9. Population estimates and percent changes for 2000 to 2007 are also shown.

ECONOMIC FACTORS

Income

Data on state ranking according to per capita personal income are gathered by the Bureau of Economic Analysis of the U. S. Department of Commerce. From 2000 to 2003, New Mexico ranked 47th among the states in per capita personal income. New Mexico ranked 46th and 45th in 2004 and 2005. In 2006 and 2007, New Mexico ranked 43rd. The 2007 per capita personal income in New Mexico was \$30,706, 20.5% lower than the United States per capita personal income of \$38,615.

Only two New Mexico counties, Los Alamos and Santa Fe, exceeded the 2007 national per capita personal income (Figure D-6). Los Alamos County's income was 42.7% higher than the nation's and 79.5% higher than the state's. The Los Alamos per capita personal income was 30.6% higher than second-ranking Santa Fe County and 57.5% higher than third-ranking Bernalillo County, the state's most populous county.

Twenty-five of the state's thirty-three counties registered per capita personal incomes less than 75% of the national figure. Four counties, Catron, Mora, Harding, and Guadalupe, had a per capita personal income less than half the national figure (Figure D-6).

Figure D-5
New Mexico Population distribution by
Race/Ethnicity, Age and Sex
New Mexico, 2007






Figure D-6
Per Capita Personal Income
Ranked by County
New Mexico and United States, 2007



For further explanation of Per Capita Personal Income, please see Technical Appendix.
 Ranking has not been tested for statistical significance.
 Source: U.S. Department of Commerce, Bureau of Economic Analysis.
 Table prepared by Bureau of Business and Economic Research (BBER), University of New Mexico, <http://www.unm.edu/~bber/>. Released April 2009.

Figure D-7
Children Under 18 at or Below Poverty
Level
Ranked by County
New Mexico and United States, 2007



For further explanation of poverty, please see Technical Appendix.
 Ranking has not been tested for statistical significance.
 Source: U.S. Census Bureau, Small Area Income and Poverty Estimates.
 www.census.gov. Current as of November 18, 2008.

Poverty

Data from the U.S. Census Bureau indicates that in 2007 New Mexico ranked 47th among states for children under 18 living in poverty with 25.2% at or below poverty level, compared to the national figure of 18.0% (Figure D-7). New Mexico is 40% higher than the nation for children under 18 living in poverty. Since 2000, the percent of children in poverty has increased by 11.1% for the United States, while New Mexico decreased by 1.2%. (See the *Technical Appendix* for information on poverty levels).

In 2007, Los Alamos and Sandoval counties were below the national figure for percent of children under age 18 at or below poverty level. Five counties were over twice the national percentage (Figure D-7).

TRENDS IN NEW MEXICO HEALTH STATISTICS

Table D-10 provides a summary of vital statistics information from 1960 through 2007. This information provides a historical perspective over the last four decades.

Demographic and Socioeconomic Status Section

**Table D-1
Population: Numbers and Percents by Age and County
New Mexico, 2007
Both Sexes**

Resident County	All Ages	Under 1 Year		1 to 4 Years		5 to 14 Years		15 to 17 Years		18 to 19 Years		20 to 24 Years	
	Number	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
New Mexico	2,053,922	26,388	1.3	99,193	4.8	280,227	13.6	99,784	4.9	67,576	3.3	150,915	7.3
Bernalillo	644,023	8,319	1.3	30,267	4.7	77,586	12.0	26,695	4.1	18,323	2.8	45,206	7.0
Catron	3,922	25	0.6	100	2.5	427	10.9	168	4.3	120	3.1	161	4.1
Chaves	63,480	756	1.2	2,829	4.5	9,292	14.6	3,403	5.4	2,480	3.9	4,687	7.4
Cibola	28,835	375	1.3	1,455	5.0	3,833	13.3	1,376	4.8	953	3.3	2,186	7.6
Colfax	14,619	137	0.9	538	3.7	1,853	12.7	662	4.5	463	3.2	821	5.6
Curry	47,261	748	1.6	2,694	5.7	6,568	13.9	2,314	4.9	1,592	3.4	4,073	8.6
De Baca	2,279	10	0.4	81	3.6	236	10.4	129	5.7	95	4.2	125	5.5
Doña Ana	205,247	3,302	1.6	11,488	5.6	28,270	13.8	9,940	4.8	7,007	3.4	20,425	10.0
Eddy	52,676	625	1.2	2,486	4.7	7,621	14.5	2,733	5.2	1,963	3.7	3,619	6.9
Grant	31,925	372	1.2	1,492	4.7	4,051	12.7	1,444	4.5	1,008	3.2	2,164	6.8
Guadalupe	4,833	47	1.0	172	3.6	514	10.6	246	5.1	164	3.4	370	7.7
Harding	817	8	1.0	9	1.1	45	5.5	23	2.8	11	1.3	43	5.3
Hidalgo	5,985	73	1.2	276	4.6	937	15.7	307	5.1	217	3.6	374	6.2
Lea	58,485	1,034	1.8	3,795	6.5	7,909	13.5	2,837	4.9	1,929	3.3	5,060	8.7
Lincoln	22,783	215	0.9	778	3.4	2,803	12.3	1,042	4.6	672	2.9	1,111	4.9
Los Alamos	20,005	182	0.9	633	3.2	2,559	12.8	938	4.7	543	2.7	713	3.6
Luna	28,189	467	1.7	1,630	5.8	4,658	16.5	1,503	5.3	1,006	3.6	1,847	6.6
McKinley	79,905	1,138	1.4	4,627	5.8	16,634	20.8	5,633	7.0	3,336	4.2	6,078	7.6
Mora	5,520	47	0.9	233	4.2	577	10.5	334	6.1	247	4.5	354	6.4
Otero	66,906	741	1.1	3,199	4.8	10,523	15.7	3,776	5.6	2,322	3.5	4,565	6.8
Quay	10,273	92	0.9	342	3.3	1,174	11.4	455	4.4	325	3.2	558	5.4
Rio Arriba	43,919	525	1.2	2,098	4.8	6,567	15.0	2,300	5.2	1,364	3.1	2,793	6.4
Roosevelt	19,093	215	1.1	952	5.0	2,698	14.1	958	5.0	781	4.1	2,054	10.8
San Juan	128,884	1,601	1.2	5,945	4.6	17,339	13.5	7,156	5.6	5,234	4.1	10,754	8.3
San Miguel	31,033	304	1.0	1,268	4.1	4,472	14.4	1,544	5.0	965	3.1	1,717	5.5
Sandoval	123,694	1,454	1.2	6,264	5.1	18,766	15.2	7,098	5.7	5,016	4.1	9,836	8.0
Santa Fe	146,295	1,616	1.1	5,870	4.0	17,363	11.9	6,143	4.2	3,960	2.7	8,649	5.9
Sierra	13,786	111	0.8	447	3.2	1,619	11.7	600	4.4	379	2.7	677	4.9
Socorro	18,788	216	1.1	921	4.9	2,739	14.6	1,010	5.4	756	4.0	1,862	9.9
Taos	32,331	364	1.1	1,310	4.1	4,025	12.4	1,410	4.4	901	2.8	1,574	4.9
Torrance	17,850	191	1.1	832	4.7	3,198	17.9	1,075	6.0	607	3.4	1,162	6.5
Union	4,474	61	1.4	213	4.8	609	13.6	243	5.4	176	3.9	186	4.2
Valencia	75,807	1,017	1.3	3,949	5.2	12,762	16.8	4,289	5.7	2,661	3.5	5,111	6.7

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

Table D-1 (continued)

25 to 34 Years		35 to 44 Years		45 to 54 Years		55 to 64 Years		65 to 74 Years		75 to 84 Years		85 Years and Older		Resident County
Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
237,926	11.6	300,860	14.6	314,296	15.3	210,988	10.3	148,017	7.2	102,034	5.0	15,718	0.8	New Mexico
78,826	12.2	100,072	15.5	102,867	16.0	66,218	10.3	46,802	7.3	37,270	5.8	5,572	0.9	Bernalillo
202	5.2	389	9.9	751	19.1	761	19.4	509	13.0	264	6.7	45	1.1	Catron
6,788	10.7	8,786	13.8	9,489	14.9	5,772	9.1	4,802	7.6	3,756	5.9	640	1.0	Chaves
3,221	11.2	4,307	14.9	4,254	14.8	2,957	10.3	2,462	8.5	1,293	4.5	163	0.6	Cibola
1,270	8.7	1,788	12.2	2,417	16.5	2,011	13.8	1,516	10.4	959	6.6	184	1.3	Colfax
6,397	13.5	6,780	14.3	6,260	13.2	4,025	8.5	3,272	6.9	2,168	4.6	370	0.8	Curry
128	5.6	302	13.3	295	12.9	272	11.9	323	14.2	247	10.8	36	1.6	De Baca
29,930	14.6	24,388	11.9	26,143	12.7	19,922	9.7	13,841	6.7	9,257	4.5	1,334	0.6	Doña Ana
4,925	9.3	7,220	13.7	8,673	16.5	5,130	9.7	3,976	7.5	3,158	6.0	547	1.0	Eddy
3,167	9.9	3,903	12.2	4,799	15.0	4,016	12.6	2,946	9.2	2,209	6.9	354	1.1	Grant
535	11.1	985	20.4	705	14.6	426	8.8	437	9.0	214	4.4	18	0.4	Guadalupe
46	5.6	78	9.5	175	21.4	108	13.2	166	20.3	88	10.8	17	2.1	Harding
540	9.0	820	13.7	849	14.2	655	10.9	502	8.4	377	6.3	58	1.0	Hidalgo
7,559	12.9	7,377	12.6	8,058	13.8	5,468	9.3	4,075	7.0	2,921	5.0	463	0.8	Lea
1,687	7.4	2,768	12.1	3,900	17.1	3,439	15.1	2,757	12.1	1,410	6.2	201	0.9	Lincoln
1,299	6.5	3,264	16.3	3,641	18.2	3,087	15.4	1,502	7.5	1,440	7.2	204	1.0	Los Alamos
2,842	10.1	3,621	12.8	3,130	11.1	2,691	9.5	2,881	10.2	1,624	5.8	289	1.0	Luna
9,266	11.6	11,846	14.8	10,164	12.7	5,814	7.3	3,543	4.4	1,575	2.0	251	0.3	McKinley
388	7.0	766	13.9	967	17.5	671	12.2	578	10.5	317	5.7	41	0.7	Mora
6,732	10.1	10,750	16.1	9,244	13.8	6,342	9.5	5,378	8.0	2,922	4.4	412	0.6	Otero
765	7.4	1,316	12.8	1,600	15.6	1,427	13.9	1,242	12.1	853	8.3	124	1.2	Quay
5,289	12.0	6,712	15.3	6,951	15.8	4,332	9.9	2,914	6.6	1,825	4.2	249	0.6	Rio Arriba
2,546	13.3	2,488	13.0	2,366	12.4	1,607	8.4	1,300	6.8	952	5.0	176	0.9	Roosevelt
13,308	10.3	19,750	15.3	19,402	15.1	13,778	10.7	8,505	6.6	5,203	4.0	909	0.7	San Juan
3,069	9.9	5,251	16.9	5,323	17.2	3,350	10.8	1,980	6.4	1,563	5.0	227	0.7	San Miguel
13,155	10.6	18,783	15.2	18,674	15.1	11,106	9.0	7,909	6.4	4,913	4.0	720	0.6	Sandoval
16,932	11.6	21,958	15.0	28,047	19.2	18,440	12.6	10,095	6.9	6,305	4.3	917	0.6	Santa Fe
925	6.7	1,680	12.2	1,806	13.1	1,748	12.7	1,905	13.8	1,653	12.0	236	1.7	Sierra
2,221	11.8	2,559	13.6	2,670	14.2	1,861	9.9	1,190	6.3	670	3.6	113	0.6	Socorro
2,961	9.2	4,484	13.9	6,243	19.3	4,425	13.7	2,867	8.9	1,519	4.7	248	0.8	Taos
2,001	11.2	2,861	16.0	2,784	15.6	1,721	9.6	919	5.1	412	2.3	87	0.5	Torrance
344	7.7	590	13.2	703	15.7	528	11.8	469	10.5	290	6.5	62	1.4	Union
8,662	11.4	12,218	16.1	10,946	14.4	6,880	9.1	4,454	5.9	2,407	3.2	451	0.6	Valencia

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

**Table D-2
Population: Numbers and Percents by Age and County
New Mexico, 2007
Males**

Resident County	All Ages	Under 1 Year		1 to 4 Years		5 to 14 Years		15 to 17 Years		18 to 19 Years		20 to 24 Years	
	Number	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
New Mexico	1,010,139	13,564	1.3	50,223	5.0	142,364	14.1	51,025	5.1	34,521	3.4	76,625	7.6
Bernalillo	315,101	4,248	1.3	15,263	4.8	39,623	12.6	13,592	4.3	9,244	2.9	22,688	7.2
Catron	1,979	11	0.6	47	2.4	225	11.4	98	5.0	74	3.7	85	4.3
Chaves	30,993	404	1.3	1,481	4.8	4,588	14.8	1,736	5.6	1,260	4.1	2,247	7.3
Cibola	14,067	188	1.3	765	5.4	1,883	13.4	709	5.0	493	3.5	1,099	7.8
Colfax	7,415	80	1.1	270	3.6	939	12.7	350	4.7	266	3.6	460	6.2
Curry	23,355	393	1.7	1,428	6.1	3,325	14.2	1,209	5.2	810	3.5	2,097	9.0
De Baca	1,118	6	0.5	38	3.4	107	9.6	68	6.1	57	5.1	74	6.6
Doña Ana	101,123	1,695	1.7	5,769	5.7	14,323	14.2	5,035	5.0	3,495	3.5	10,511	10.4
Eddy	25,726	315	1.2	1,249	4.9	3,760	14.6	1,400	5.4	1,039	4.0	1,852	7.2
Grant	15,433	206	1.3	746	4.8	2,067	13.4	749	4.9	524	3.4	1,054	6.8
Guadalupe	2,749	24	0.9	79	2.9	269	9.8	121	4.4	90	3.3	240	8.7
Harding	394	4	1.0	5	1.3	21	5.3	15	3.8	0	0.0	21	5.3
Hidalgo	2,979	38	1.3	141	4.7	480	16.1	161	5.4	116	3.9	182	6.1
Lea	29,216	533	1.8	1,937	6.6	4,034	13.8	1,431	4.9	955	3.3	2,621	9.0
Lincoln	11,186	102	0.9	384	3.4	1,452	13.0	553	4.9	362	3.2	590	5.3
Los Alamos	10,011	91	0.9	314	3.1	1,288	12.9	494	4.9	305	3.0	382	3.8
Luna	13,736	240	1.7	840	6.1	2,275	16.6	736	5.4	520	3.8	884	6.4
McKinley	38,597	584	1.5	2,338	6.1	8,304	21.5	2,902	7.5	1,737	4.5	3,010	7.8
Mora	2,799	26	0.9	117	4.2	316	11.3	169	6.0	126	4.5	191	6.8
Otero	33,029	376	1.1	1,593	4.8	5,556	16.8	1,934	5.9	1,157	3.5	2,214	6.7
Quay	5,024	41	0.8	180	3.6	617	12.3	236	4.7	166	3.3	286	5.7
Rio Arriba	21,855	281	1.3	1,006	4.6	3,254	14.9	1,171	5.4	705	3.2	1,456	6.7
Roosevelt	9,470	109	1.2	470	5.0	1,410	14.9	491	5.2	385	4.1	1,030	10.9
San Juan	62,980	850	1.3	2,936	4.7	8,707	13.8	3,556	5.6	2,584	4.1	5,216	8.3
San Miguel	15,133	160	1.1	636	4.2	2,267	15.0	794	5.2	499	3.3	882	5.8
Sandoval	61,471	723	1.2	3,243	5.3	9,714	15.8	3,702	6.0	2,614	4.3	5,070	8.2
Santa Fe	71,401	833	1.2	2,977	4.2	8,789	12.3	3,139	4.4	2,078	2.9	4,597	6.4
Sierra	6,892	59	0.9	225	3.3	826	12.0	329	4.8	203	2.9	351	5.1
Socorro	9,489	111	1.2	474	5.0	1,346	14.2	551	5.8	433	4.6	997	10.5
Taos	15,841	177	1.1	690	4.4	2,063	13.0	698	4.4	462	2.9	843	5.3
Torrance	9,255	101	1.1	432	4.7	1,652	17.8	545	5.9	301	3.3	644	7.0
Union	2,270	33	1.5	125	5.5	309	13.6	120	5.3	87	3.8	116	5.1
Valencia	38,052	522	1.4	2,025	5.3	6,575	17.3	2,231	5.9	1,374	3.6	2,635	6.9

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

Table D-2 (continued)

25 to 34 Years		35 to 44 Years		45 to 54 Years		55 to 64 Years		65 to 74 Years		75 to 84 Years		85 Years and Older		Resident County
Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
119,644	11.8	149,221	14.8	152,867	15.1	101,731	10.1	68,697	6.8	43,130	4.3	6,527	0.6	New Mexico
39,944	12.7	49,968	15.9	49,660	15.8	31,948	10.1	21,099	6.7	15,490	4.9	2,334	0.7	Bernalillo
97	4.9	158	8.0	384	19.4	393	19.9	282	14.2	112	5.7	13	0.7	Catron
3,470	11.2	4,570	14.7	4,577	14.8	2,697	8.7	2,106	6.8	1,595	5.1	262	0.8	Chaves
1,543	11.0	2,051	14.6	2,134	15.2	1,365	9.7	1,188	8.4	580	4.1	69	0.5	Cibola
648	8.7	925	12.5	1,240	16.7	1,032	13.9	740	10.0	396	5.3	69	0.9	Colfax
3,335	14.3	3,325	14.2	3,105	13.3	1,858	8.0	1,528	6.5	808	3.5	134	0.6	Curry
58	5.2	146	13.1	168	15.0	132	11.8	158	14.1	93	8.3	13	1.2	De Baca
15,451	15.3	11,758	11.6	12,586	12.4	9,563	9.5	6,295	6.2	4,077	4.0	565	0.6	Doña Ana
2,317	9.0	3,471	13.5	4,440	17.3	2,508	9.7	1,815	7.1	1,352	5.3	208	0.8	Eddy
1,514	9.8	1,829	11.9	2,230	14.4	1,985	12.9	1,415	9.2	970	6.3	144	0.9	Grant
388	14.1	591	21.5	433	15.8	203	7.4	215	7.8	88	3.2	8	0.3	Guadalupe
22	5.6	33	8.4	91	23.1	58	14.7	85	21.6	33	8.4	6	1.5	Harding
250	8.4	417	14.0	429	14.4	314	10.5	265	8.9	164	5.5	22	0.7	Hidalgo
3,878	13.3	3,781	12.9	4,111	14.1	2,695	9.2	1,851	6.3	1,207	4.1	182	0.6	Lea
812	7.3	1,316	11.8	1,862	16.6	1,629	14.6	1,366	12.2	673	6.0	85	0.8	Lincoln
661	6.6	1,588	15.9	1,810	18.1	1,610	16.1	722	7.2	649	6.5	97	1.0	Los Alamos
1,305	9.5	1,695	12.3	1,567	11.4	1,278	9.3	1,474	10.7	792	5.8	130	0.9	Luna
4,472	11.6	5,648	14.6	4,685	12.1	2,704	7.0	1,449	3.8	651	1.7	113	0.3	McKinley
181	6.5	393	14.0	465	16.6	356	12.7	284	10.1	154	5.5	21	0.8	Mora
3,333	10.1	5,368	16.3	4,445	13.5	3,026	9.2	2,637	8.0	1,216	3.7	174	0.5	Otero
367	7.3	624	12.4	780	15.5	704	14.0	610	12.1	371	7.4	42	0.8	Quay
2,683	12.3	3,394	15.5	3,433	15.7	2,238	10.2	1,355	6.2	782	3.6	97	0.4	Rio Arriba
1,298	13.7	1,267	13.4	1,102	11.6	804	8.5	608	6.4	416	4.4	80	0.8	Roosevelt
6,086	9.7	10,018	15.9	9,699	15.4	6,683	10.6	4,264	6.8	2,009	3.2	372	0.6	San Juan
1,514	10.0	2,541	16.8	2,587	17.1	1,618	10.7	875	5.8	668	4.4	92	0.6	San Miguel
6,561	10.7	9,224	15.0	9,211	15.0	5,373	8.7	3,628	5.9	2,114	3.4	294	0.5	Sandoval
8,589	12.0	10,887	15.2	13,173	18.4	8,597	12.0	4,663	6.5	2,694	3.8	385	0.5	Santa Fe
464	6.7	820	11.9	890	12.9	796	11.5	1,030	14.9	786	11.4	113	1.6	Sierra
1,150	12.1	1,238	13.0	1,324	14.0	947	10.0	578	6.1	292	3.1	48	0.5	Socorro
1,561	9.9	2,219	14.0	2,963	18.7	2,094	13.2	1,395	8.8	580	3.7	96	0.6	Taos
1,119	12.1	1,488	16.1	1,436	15.5	855	9.2	466	5.0	188	2.0	28	0.3	Torrance
193	8.5	283	12.5	370	16.3	268	11.8	211	9.3	132	5.8	23	1.0	Union
4,380	11.5	6,187	16.3	5,477	14.4	3,400	8.9	2,040	5.4	998	2.6	208	0.5	Valencia

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

**Table D-3
Population: Numbers and Percents by Age and County
New Mexico, 2007
Females**

Resident County	All Ages	Under 1 Year		1 to 4 Years		5 to 14 Years		15 to 17 Years		18 to 19 Years		20 to 24 Years	
	Number	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
New Mexico	1,043,783	12,824	1.2	48,970	4.7	137,863	13.2	48,759	4.7	33,055	3.2	74,290	7.1
Bernalillo	328,922	4,071	1.2	15,004	4.6	37,963	11.5	13,103	4.0	9,079	2.8	22,518	6.8
Catron	1,943	14	0.7	53	2.7	202	10.4	70	3.6	46	2.4	76	3.9
Chaves	32,487	352	1.1	1,348	4.1	4,704	14.5	1,667	5.1	1,220	3.8	2,440	7.5
Cibola	14,768	187	1.3	690	4.7	1,950	13.2	667	4.5	460	3.1	1,087	7.4
Colfax	7,204	57	0.8	268	3.7	914	12.7	312	4.3	197	2.7	361	5.0
Curry	23,906	355	1.5	1,266	5.3	3,243	13.6	1,105	4.6	782	3.3	1,976	8.3
De Baca	1,161	4	0.3	43	3.7	129	11.1	61	5.3	38	3.3	51	4.4
Doña Ana	104,124	1,607	1.5	5,719	5.5	13,947	13.4	4,905	4.7	3,512	3.4	9,914	9.5
Eddy	26,950	310	1.2	1,237	4.6	3,861	14.3	1,333	4.9	924	3.4	1,767	6.6
Grant	16,492	166	1.0	746	4.5	1,984	12.0	695	4.2	484	2.9	1,110	6.7
Guadalupe	2,084	23	1.1	93	4.5	245	11.8	125	6.0	74	3.6	130	6.2
Harding	423	4	0.9	4	0.9	24	5.7	8	1.9	11	2.6	22	5.2
Hidalgo	3,006	35	1.2	135	4.5	457	15.2	146	4.9	101	3.4	192	6.4
Lea	29,269	501	1.7	1,858	6.3	3,875	13.2	1,406	4.8	974	3.3	2,439	8.3
Lincoln	11,597	113	1.0	394	3.4	1,351	11.6	489	4.2	310	2.7	521	4.5
Los Alamos	9,994	91	0.9	319	3.2	1,271	12.7	444	4.4	238	2.4	331	3.3
Luna	14,453	227	1.6	790	5.5	2,383	16.5	767	5.3	486	3.4	963	6.7
McKinley	41,308	554	1.3	2,289	5.5	8,330	20.2	2,731	6.6	1,599	3.9	3,068	7.4
Mora	2,721	21	0.8	116	4.3	261	9.6	165	6.1	121	4.4	163	6.0
Otero	33,877	365	1.1	1,606	4.7	4,967	14.7	1,842	5.4	1,165	3.4	2,351	6.9
Quay	5,249	51	1.0	162	3.1	557	10.6	219	4.2	159	3.0	272	5.2
Rio Arriba	22,064	244	1.1	1,092	4.9	3,313	15.0	1,129	5.1	659	3.0	1,337	6.1
Roosevelt	9,623	106	1.1	482	5.0	1,288	13.4	467	4.9	396	4.1	1,024	10.6
San Juan	65,904	751	1.1	3,009	4.6	8,632	13.1	3,600	5.5	2,650	4.0	5,538	8.4
San Miguel	15,900	144	0.9	632	4.0	2,205	13.9	750	4.7	466	2.9	835	5.3
Sandoval	62,223	731	1.2	3,021	4.9	9,052	14.5	3,396	5.5	2,402	3.9	4,766	7.7
Santa Fe	74,894	783	1.0	2,893	3.9	8,574	11.4	3,004	4.0	1,882	2.5	4,052	5.4
Sierra	6,894	52	0.8	222	3.2	793	11.5	271	3.9	176	2.6	326	4.7
Socorro	9,299	105	1.1	447	4.8	1,393	15.0	459	4.9	323	3.5	865	9.3
Taos	16,490	187	1.1	620	3.8	1,962	11.9	712	4.3	439	2.7	731	4.4
Torrance	8,595	90	1.0	400	4.7	1,546	18.0	530	6.2	306	3.6	518	6.0
Union	2,204	28	1.3	88	4.0	300	13.6	123	5.6	89	4.0	70	3.2
Valencia	37,755	495	1.3	1,924	5.1	6,187	16.4	2,058	5.5	1,287	3.4	2,476	6.6

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

Table D-3 (continued)

25 to 34 Years		35 to 44 Years		45 to 54 Years		55 to 64 Years		65 to 74 Years		75 to 84 Years		85 Years and Older		Resident County
Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
118,282	11.3	151,639	14.5	161,429	15.5	109,257	10.5	79,320	7.6	58,904	5.6	9,191	0.9	New Mexico
38,882	11.8	50,104	15.2	53,207	16.2	34,270	10.4	25,703	7.8	21,780	6.6	3,238	1.0	Bernalillo
105	5.4	231	11.9	367	18.9	368	18.9	227	11.7	152	7.8	32	1.6	Catron
3,318	10.2	4,216	13.0	4,912	15.1	3,075	9.5	2,696	8.3	2,161	6.7	378	1.2	Chaves
1,678	11.4	2,256	15.3	2,120	14.4	1,592	10.8	1,274	8.6	713	4.8	94	0.6	Cibola
622	8.6	863	12.0	1,177	16.3	979	13.6	776	10.8	563	7.8	115	1.6	Colfax
3,062	12.8	3,455	14.5	3,155	13.2	2,167	9.1	1,744	7.3	1,360	5.7	236	1.0	Curry
70	6.0	156	13.4	127	10.9	140	12.1	165	14.2	154	13.3	23	2.0	De Baca
14,479	13.9	12,630	12.1	13,557	13.0	10,359	9.9	7,546	7.2	5,180	5.0	769	0.7	Doña Ana
2,608	9.7	3,749	13.9	4,233	15.7	2,622	9.7	2,161	8.0	1,806	6.7	339	1.3	Eddy
1,653	10.0	2,074	12.6	2,569	15.6	2,031	12.3	1,531	9.3	1,239	7.5	210	1.3	Grant
147	7.1	394	18.9	272	13.1	223	10.7	222	10.7	126	6.0	10	0.5	Guadalupe
24	5.7	45	10.6	84	19.9	50	11.8	81	19.1	55	13.0	11	2.6	Harding
290	9.6	403	13.4	420	14.0	341	11.3	237	7.9	213	7.1	36	1.2	Hidalgo
3,681	12.6	3,596	12.3	3,947	13.5	2,773	9.5	2,224	7.6	1,714	5.9	281	1.0	Lea
875	7.5	1,452	12.5	2,038	17.6	1,810	15.6	1,391	12.0	737	6.4	116	1.0	Lincoln
638	6.4	1,676	16.8	1,831	18.3	1,477	14.8	780	7.8	791	7.9	107	1.1	Los Alamos
1,537	10.6	1,926	13.3	1,563	10.8	1,413	9.8	1,407	9.7	832	5.8	159	1.1	Luna
4,794	11.6	6,198	15.0	5,479	13.3	3,110	7.5	2,094	5.1	924	2.2	138	0.3	McKinley
207	7.6	373	13.7	502	18.4	315	11.6	294	10.8	163	6.0	20	0.7	Mora
3,399	10.0	5,382	15.9	4,799	14.2	3,316	9.8	2,741	8.1	1,706	5.0	238	0.7	Otero
398	7.6	692	13.2	820	15.6	723	13.8	632	12.0	482	9.2	82	1.6	Quay
2,606	11.8	3,318	15.0	3,518	15.9	2,094	9.5	1,559	7.1	1,043	4.7	152	0.7	Rio Arriba
1,248	13.0	1,221	12.7	1,264	13.1	803	8.3	692	7.2	536	5.6	96	1.0	Roosevelt
7,222	11.0	9,732	14.8	9,703	14.7	7,095	10.8	4,241	6.4	3,194	4.8	537	0.8	San Juan
1,555	9.8	2,710	17.0	2,736	17.2	1,732	10.9	1,105	6.9	895	5.6	135	0.8	San Miguel
6,594	10.6	9,559	15.4	9,463	15.2	5,733	9.2	4,281	6.9	2,799	4.5	426	0.7	Sandoval
8,343	11.1	11,071	14.8	14,874	19.9	9,843	13.1	5,432	7.3	3,611	4.8	532	0.7	Santa Fe
461	6.7	860	12.5	916	13.3	952	13.8	875	12.7	867	12.6	123	1.8	Sierra
1,071	11.5	1,321	14.2	1,346	14.5	914	9.8	612	6.6	378	4.1	65	0.7	Socorro
1,400	8.5	2,265	13.7	3,280	19.9	2,331	14.1	1,472	8.9	939	5.7	152	0.9	Taos
882	10.3	1,373	16.0	1,348	15.7	866	10.1	453	5.3	224	2.6	59	0.7	Torrance
151	6.9	307	13.9	333	15.1	260	11.8	258	11.7	158	7.2	39	1.8	Union
4,282	11.3	6,031	16.0	5,469	14.5	3,480	9.2	2,414	6.4	1,409	3.7	243	0.6	Valencia

Due to rounding, totals may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Demographic and Socioeconomic Status Section

**Table D-4
Population: Numbers and Percents by Race/Ethnicity, Sex and Age
New Mexico And United States, 2007**

Race / Ethnicity and Sex	Under 1 Year		1 to 4 Years		5 to 14 Years		15 to 17 Years		18 to 19 Years		20 to 24 Years		25 to 34 Years	
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
New Mexico														
All Races														
Both Sexes	26,388	1.3	99,193	4.8	280,227	13.6	99,784	4.9	67,576	3.3	150,915	7.3	237,926	11.6
Male	13,564	1.3	50,223	5.0	142,364	14.1	51,025	5.1	34,521	3.4	76,625	7.6	119,644	11.8
Female	12,824	1.2	48,970	4.7	137,863	13.2	48,759	4.7	33,055	3.2	74,290	7.1	118,282	11.3
American Indian or Alaska Native														
Both Sexes	3,535	1.6	13,664	6.0	41,828	18.5	15,133	6.7	10,005	4.4	20,486	9.0	28,306	12.5
Male	1,805	1.7	6,798	6.2	20,830	19.1	7,662	7.0	5,035	4.6	9,645	8.8	13,662	12.5
Female	1,730	1.5	6,866	5.8	20,998	17.8	7,471	6.3	4,970	4.2	10,841	9.2	14,644	12.4
Asian or Pacific Islander														
Both Sexes	454	1.3	1,626	4.7	5,179	14.9	1,690	4.9	1,135	3.3	2,557	7.4	5,704	16.4
Male	241	1.5	841	5.1	2,562	15.6	890	5.4	597	3.6	1,411	8.6	2,781	16.9
Female	213	1.2	785	4.3	2,617	14.3	800	4.4	538	2.9	1,146	6.3	2,923	16.0
Black or African American														
Both Sexes	694	1.3	2,630	4.8	9,797	17.9	3,099	5.7	2,130	3.9	5,278	9.7	7,847	14.4
Male	349	1.2	1,306	4.6	4,932	17.5	1,562	5.5	1,124	4.0	2,772	9.8	4,093	14.5
Female	345	1.3	1,324	5.0	4,865	18.4	1,537	5.8	1,006	3.8	2,506	9.5	3,754	14.2
Hispanic														
Both Sexes	13,693	1.6	51,060	6.0	137,022	16.2	45,826	5.4	31,003	3.7	69,997	8.3	113,637	13.4
Male	7,026	1.7	25,962	6.2	69,612	16.6	23,291	5.5	15,789	3.8	35,253	8.4	57,942	13.8
Female	6,667	1.6	25,098	5.9	67,410	15.8	22,535	5.3	15,214	3.6	34,744	8.1	55,695	13.1
White														
Both Sexes	8,012	0.9	30,213	3.4	86,401	9.7	34,036	3.8	23,303	2.6	52,597	5.9	82,432	9.3
Male	4,143	1.0	15,316	3.5	44,428	10.2	17,620	4.0	11,976	2.7	27,544	6.3	41,166	9.4
Female	3,869	0.9	14,897	3.3	41,973	9.2	16,416	3.6	11,327	2.5	25,053	5.5	41,266	9.1
United States														
All Races														
Both Sexes	4,257,020	1.4	16,467,105	5.5	40,163,937	13.3	13,013,671	4.3	8,460,019	2.8	21,032,396	7.0	40,590,926	13.5
Male	2,178,808	1.5	8,424,049	5.7	20,548,505	13.8	6,666,273	4.5	4,340,596	2.9	10,852,937	7.3	20,682,550	13.9
Female	2,078,212	1.4	8,043,056	5.3	19,615,432	12.8	6,347,398	4.1	4,119,423	2.7	10,179,459	6.7	19,908,376	13.0

Notes: Due to rounding, percents may not add to 100.

Percents are by sex and reflect the age distribution within each Race/Ethnicity category.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

U.S. Data Source: U.S. Bureau of the Census, Population Division. www.census.gov.

Demographic and Socioeconomic Status Section

Table D-4 (continued)

35 to 44 Years		45 to 54 Years		55 to 64 Years		65 to 74 Years		75 to 84 Years		85 Plus Years		Race / Ethnicity and Sex
Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	
New Mexico												
All Races												
300,860	14.6	314,296	15.3	210,988	10.3	148,017	7.2	102,034	5.0	15,718	0.8	Both Sexes
149,221	14.8	152,867	15.1	101,731	10.1	68,697	6.8	43,130	4.3	6,527	0.6	Male
151,639	14.5	161,429	15.5	109,257	10.5	79,320	7.6	58,904	5.6	9,191	0.9	Female
American Indian or Alaska Native												
34,016	15.0	28,011	12.4	15,916	7.0	9,783	4.3	5,231	2.3	756	0.3	Both Sexes
16,546	15.2	12,997	11.9	7,334	6.7	4,248	3.9	2,114	1.9	309	0.3	Male
17,470	14.8	15,014	12.8	8,582	7.3	5,535	4.7	3,117	2.6	447	0.4	Female
Asian or Pacific Islander												
6,075	17.5	4,980	14.3	2,813	8.1	1,615	4.6	846	2.4	71	0.2	Both Sexes
2,812	17.1	2,158	13.1	1,140	6.9	644	3.9	318	1.9	29	0.2	Male
3,263	17.8	2,822	15.4	1,673	9.1	971	5.3	528	2.9	42	0.2	Female
Black or African American												
8,261	15.1	7,030	12.9	3,612	6.6	2,531	4.6	1,486	2.7	201	0.4	Both Sexes
4,477	15.9	3,775	13.4	1,894	6.7	1,230	4.4	575	2.0	76	0.3	Male
3,784	14.3	3,255	12.3	1,718	6.5	1,301	4.9	911	3.4	125	0.5	Female
Hispanic												
127,075	15.0	112,165	13.2	65,842	7.8	46,275	5.5	29,517	3.5	4,118	0.5	Both Sexes
63,371	15.1	55,045	13.1	31,782	7.6	21,001	5.0	12,673	3.0	1,771	0.4	Male
63,704	14.9	57,120	13.4	34,060	8.0	25,274	5.9	16,844	3.9	2,347	0.6	Female
White												
125,433	14.1	162,110	18.2	122,805	13.8	87,813	9.9	64,954	7.3	10,572	1.2	Both Sexes
62,015	14.2	78,892	18.1	59,581	13.7	41,574	9.5	27,450	6.3	4,342	1.0	Male
63,418	13.9	83,218	18.3	63,224	13.9	46,239	10.2	37,504	8.2	6,230	1.4	Female
United States												
All Races												
43,161,289	14.3	43,874,760	14.5	32,712,076	10.8	19,352,149	6.4	13,023,511	4.3	5,512,298	1.8	Both Sexes
21,618,734	14.5	21,594,913	14.5	15,775,088	10.6	8,886,973	6.0	5,312,673	3.6	1,776,799	1.2	Male
21,542,555	14.1	22,279,847	14.6	16,936,988	11.1	10,465,176	6.8	7,710,838	5.0	3,735,499	2.4	Female

Notes: Due to rounding, percents may not add to 100.

Percents are by sex and reflect the age distribution within each Race/Ethnicity category.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

U.S. Data Source: U.S. Bureau of the Census, Population Division. www.census.gov.

Demographic and Socioeconomic Status Section

**Table D-5
Population: Numbers and Percents by Race/Ethnicity, County and Health Region
New Mexico, 2007**

Resident County	All Races		American Indian or Alaska Native		Asian or Pacific Islander		Black or African American		Hispanic		White	
	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
New Mexico	2,053,922		226,670	11.0	34,745	1.7	54,596	2.7	847,230	41.2	890,681	43.4
Bernalillo	644,023		37,279	5.8	17,527	2.7	23,198	3.6	265,951	41.3	300,068	46.6
Catron	3,922		108	2.8	23	0.6	9	0.2	740	18.9	3,042	77.6
Chaves	63,480		1,249	2.0	705	1.1	1,802	2.8	28,470	44.8	31,254	49.2
Cibola	28,835		11,917	41.3	149	0.5	423	1.5	9,074	31.5	7,272	25.2
Colfax	14,619		362	2.5	54	0.4	74	0.5	6,763	46.3	7,366	50.4
Curry	47,261		1,082	2.3	1,478	3.1	3,911	8.3	14,804	31.3	25,986	55.0
De Baca	2,279		31	1.4	5	0.2	5	0.2	776	34.1	1,462	64.2
Doña Ana	205,247		4,771	2.3	2,781	1.4	5,324	2.6	127,105	61.9	65,266	31.8
Eddy	52,676		998	1.9	438	0.8	1,180	2.2	20,752	39.4	29,308	55.6
Grant	31,925		725	2.3	191	0.6	363	1.1	15,042	47.1	15,604	48.9
Guadalupe	4,833		59	1.2	37	0.8	84	1.7	3,792	78.5	861	17.8
Harding	817		9	1.1	0	0.0	2	0.2	373	45.7	433	53.0
Hidalgo	5,985		51	0.9	38	0.6	35	0.6	3,275	54.7	2,586	43.2
Lea	58,485		953	1.6	444	0.8	3,178	5.4	24,970	42.7	28,940	49.5
Lincoln	22,783		641	2.8	113	0.5	245	1.1	6,019	26.4	15,765	69.2
Los Alamos	20,005		310	1.5	1,129	5.6	146	0.7	2,495	12.5	15,925	79.6
Luna	28,189		453	1.6	156	0.6	608	2.2	16,590	58.9	10,382	36.8
McKinley	79,905		62,754	78.5	519	0.6	560	0.7	6,991	8.7	9,081	11.4
Mora	5,520		66	1.2	10	0.2	15	0.3	4,286	77.6	1,143	20.7
Otero	66,906		4,749	7.1	1,355	2.0	3,235	4.8	21,480	32.1	36,087	53.9
Quay	10,273		152	1.5	149	1.5	122	1.2	3,799	37.0	6,051	58.9
Rio Arriba	43,919		7,092	16.1	252	0.6	331	0.8	30,297	69.0	5,947	13.5
Roosevelt	19,093		321	1.7	162	0.8	458	2.4	6,566	34.4	11,586	60.7
San Juan	128,884		53,050	41.2	818	0.6	1,237	1.0	18,745	14.5	55,034	42.7
San Miguel	31,033		812	2.6	289	0.9	391	1.3	23,134	74.5	6,407	20.6
Sandoval	123,694		20,482	16.6	2,083	1.7	3,114	2.5	37,408	30.2	60,607	49.0
Santa Fe	146,295		6,348	4.3	2,238	1.5	1,965	1.3	69,096	47.2	66,648	45.6
Sierra	13,786		315	2.3	44	0.3	81	0.6	3,990	28.9	9,356	67.9
Socorro	18,788		2,597	13.8	343	1.8	200	1.1	8,523	45.4	7,125	37.9
Taos	32,331		2,432	7.5	273	0.8	250	0.8	17,229	53.3	12,147	37.6
Torrance	17,850		652	3.7	124	0.7	531	3.0	6,675	37.4	9,868	55.3
Union	4,474		50	1.1	21	0.5	0	0.0	1,584	35.4	2,819	63.0
Valencia	75,807		3,800	5.0	797	1.1	1,519	2.0	40,436	53.3	29,255	38.6
Health Region												
Northwest	437,125		152,003	34.8	4,366	1.0	6,853	1.6	112,654	25.8	161,249	36.9
Northeast	303,029		17,531	5.8	4,303	1.4	3,256	1.1	158,676	52.4	119,263	39.4
Bernalillo County	644,023		37,279	5.8	17,527	2.7	23,198	3.6	265,951	41.3	300,068	46.6
Southeast	254,364		4,795	1.9	3,381	1.3	10,658	4.2	100,510	39.5	135,020	53.1
Southwest	415,381		15,062	3.6	5,168	1.2	10,631	2.6	209,439	50.4	175,081	42.1

Due to rounding, percents may not add to 100. Percents based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. Differences in age-specific numbers or percentages between 2006 and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates.

Table D-6
Components of Population Change
New Mexico, 1950-2007

Net Change		Components of Change (Percent)		
Time Period	Number (Persons)*	Percent	Natural Increase	Net Migration
2000-2007	228,034	12.5	5.4	7.1
1990-2000	304,314	20.0	9.9	10.1
1980-1990	218,271	16.7	13.4	3.3
1970-1980	286,248	28.1	14.2	13.9
1960-1970	66,032	6.9	19.4	-12.4
1950-1960	269,836	39.6	29.7	9.9

Note: Natural Increase refers to the number of births minus the number of deaths. Net migration is the difference between migration in and out of an area.

*Population Source: Populations for 1960 and earlier are based on U.S. Census, April 1 enumeration counts.

Table D-7
Percent Distribution of New Mexico Population by Age Group
Population Counts and Estimates: 1990, 2000 and 2007
Population Projections: 2020 and 2030

Age Group	1990 ¹	2000 ¹	2007 ²	2020 ³	2030 ³	Percent Population Change 1990 to 2030
Under 5 Years	8.5	7.2	6.1	7.0	6.1	-28.2
5 to 14 Years	16.6	15.9	13.6	13.2	12.9	-22.3
15 to 24 Years	14.6	14.7	15.5	11.4	12.7	-13.0
25 to 34 Years	16.9	12.9	11.6	14.2	11.0	-34.9
35 to 44 Years	15.0	15.5	14.6	11.5	13.8	- 8.0
45 to 54 Years	9.7	13.5	15.3	12.7	11.9	22.7
55 to 64 Years	8.0	8.7	10.3	14.0	12.2	52.5
65 and Older	10.7	11.7	12.9	16.0	19.3	80.4

Sources: ¹1990 and 2000 population counts from the U.S. Census Bureau, www.census.gov.

²2007 population estimates from the Bureau of Business and Economic Research (BBER), University of New Mexico, www.unm.edu/~bber.

³2020 and 2030 population projections from the Bureau of Business and Economic Research (BBER), University of New Mexico, www.unm.edu/~bber.

**Table D-8
Population Estimates, Components of Change, Birth/Death Ratio, Percent Contribution to State's Population Increase and Population Density
New Mexico and United States
2006 and 2007 Estimates and Other Population Characteristics; 2000-2007 Components of Change**

Resident County	Population Estimates		% Net Change	Components of Change 2000-2007		Other Population Characteristics 2007		
	2006	2007	2006-2007	Natural Increase	Net Migration	Birth to Death Ratio***	% Contribution to State's Population Increase	Population Density ****
United States	299,398,484	301,621,157	0.7	N/A	N/A	1.8	N/A	85.3
New Mexico	2,011,406	2,053,923	2.1	111,813	123,063	2.0	100.0	16.9
Bernalillo	628,188	644,023	2.5	35,887	52,134	2.0	37.2	552.3
Catron	3,824	3,922	2.6	-89	468	0.4	0.2	0.6
Chaves	63,166	63,480	0.5	2,944	-846	1.8	0.7	10.5
Cibola	28,683	28,835	0.5	1,724	1,516	2.0	0.4	6.4
Colfax	14,540	14,619	0.5	110	320	1.2	0.2	3.9
Curry	46,666	47,261	1.3	3,964	-1,747	2.5	1.4	33.6
De Baca	2,271	2,279	0.4	-70	109	0.7	0.0	1.0
Doña Ana	198,625	205,247	3.3	16,416	14,149	2.7	15.6	53.9
Eddy	52,449	52,676	0.4	1,663	-645	1.6	0.5	12.6
Grant	31,733	31,925	0.6	489	434	1.0	0.5	8.0
Guadalupe	4,821	4,833	0.2	45	108	0.9	0.0	1.6
Harding	814	817	0.4	-35	42	0.7	0.0	0.4
Hidalgo	5,960	5,985	0.4	125	-72	1.2	0.1	1.7
Lea	58,175	58,485	0.5	4,174	-1,200	2.6	0.7	13.3
Lincoln	22,523	22,783	1.2	440	2,932	1.3	0.6	4.7
Los Alamos	19,906	20,005	0.5	656	1,006	1.6	0.2	183.5
Luna	27,844	28,189	1.2	1,202	1,971	1.7	0.8	9.5
McKinley	79,781	79,905	0.2	7,176	-2,069	2.8	0.3	14.7
Mora	5,472	5,520	0.9	18	322	0.9	0.1	2.9
Otero	66,027	66,906	1.3	3,051	1,557	1.8	2.1	10.1
Quay	10,208	10,273	0.6	-77	195	1.0	0.2	3.6
Rio Arriba	43,530	43,919	0.9	2,642	86	1.5	0.9	7.5
Roosevelt	18,858	19,094	1.3	1,318	-243	2.1	0.6	7.8
San Juan	127,618	128,884	1.0	10,090	4,993	2.6	3.0	23.4
San Miguel	30,817	31,033	0.7	872	35	1.3	0.5	6.6
Sandoval*	114,929	123,694	7.6	5,491	27,619	2.0	20.6	33.3
Santa Fe**	144,809	146,295	1.0	6,448	10,560	2.0	3.5	76.6
Sierra	13,726	13,786	0.4	-875	1,391	0.5	0.1	3.3
Socorro	18,656	18,788	0.7	838	-128	1.6	0.3	2.8
Taos	32,127	32,331	0.6	1,040	1,312	1.4	0.5	14.7
Torrance	18,302	17,850	-2.5	483	452	1.6	-1.1	5.3
Union	4,470	4,474	0.1	-45	345	1.4	0.0	1.2
Valencia	71,888	75,807	5.5	3,703	5,952	1.8	9.2	71.0
Health Region								
Northwest	422,899	437,125	3.4	28,184	38,011	2.3	33.5	21.6
Northeast	300,492	303,029	0.8	11,786	14,094	1.6	6.0	11.1
Bernalillo County	628,188	644,023	2.5	35,887	52,134	2.0	37.2	552.3
Southeast	252,607	254,365	0.7	13,881	-4,335	2.0	4.1	9.8
Southwest	407,220	415,381	2.0	22,080	23,154	1.9	19.2	8.9

Natural increase is the excess of births minus deaths; net migration is the difference between immigration (in-migration) and emigration (out-migration). For counties included in state health regions, please see Technical Appendix.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>.

*Note from BBER: In light of revised data on housing unit stock in Sandoval County, the estimate for 2006 was up-adjusted from 111,505 to 114,929. A similar adjustment was made for 2005, which is not included in this table. The 2006 total New Mexico population increased accordingly.

**Note from BBER: Recent review of economic and housing-unit-occupancy data sources has indicated a need to down-adjust earlier BBER Santa Fe County estimates for 2006 and 2007. Therefore, the 2006 estimate for Santa Fe County was revised from 147,413 to 144,809.

Birth to death ratio is calculated by dividing the number of births by the number of deaths.

****Population density is persons per square mile based on land area in square miles.

U.S. Data Source: U.S. Bureau of the Census, Population Division. www.census.gov. 2007 U.S. births and deaths not available at time of publication.

Table D-9
Population, Births and Deaths by City
Population, 2000 and 2007, Births and Deaths, 2007

City	Population			Births	Deaths	City	Population			Births	Deaths
	2000 Census Counts	2007 Estimate	Percent Change 2000-2007				2000 Census Counts	2007 Estimate	Percent Change 2000-2007		
Alamogordo	35,582	35,607	0.1	566	328	Lake Arthur	432	437	1.2	9	3
Albuquerque	448,607	518,271	15.5	9,568	4,681	Las Cruces	74,267	89,722	20.8	2,017	964
Artesia	10,692	10,485	-1.9	267	139	Las Vegas	14,565	13,539	-7.0	262	192
Aztec	6,378	6,810	6.8	268	112	Logan	1,094	952	-13.0	7	19
Bayard	2,534	2,376	-6.2	47	29	Lordsburg	3,379	2,665	-21.1	47	43
Belen	6,901	7,159	3.7	282	217	Los Lunas	10,034	12,115	20.7	668	278
Bernalillo	6,611	7,121	7.7	130	63	Los Ranchos De Albuquerque	5,092	5,480	7.6	3	2
Bloomfield	6,417	7,129	11.1	257	89	Loving	1,326	1,295	-2.3	32	14
Bosque Farms	3,931	4,002	1.8	48	36	Lovington	9,471	9,793	3.4	299	110
Capitan	1,443	1,445	0.1	24	17	Magdalena	913	859	-5.9	33	7
Carlsbad	25,625	25,033	-2.3	480	340	Maxwell	274	248	-9.5	4	2
Carrizozo	1,036	1,029	-0.7	7	16	Melrose	736	711	-3.4	10	11
Causey	52	60	15.4	2	0	Mesilla	2,180	2,195	0.7	43	15
Chama	1,199	1,167	-2.7	22	17	Milan	1,891	2,481	31.2	46	22
Cimarron	917	832	-9.3	10	6	Moriarty	1,765	1,769	0.2	102	51
Clayton	2,524	2,100	-16.8	27	21	Mosquero	120	87	-27.5	1	3
Cloudcroft	749	762	1.7	12	9	Mountainair	1,116	1,054	-5.6	23	33
Clovis	32,667	33,182	1.6	809	320	Pecos	1,441	1,354	-6.0	45	24
Columbus	1,765	1,846	4.6	43	11	Portales	11,131	11,992	7.7	319	144
Corona	165	164	-0.6	6	3	Questa	1,864	1,870	0.3	38	17
Corrales	7,334	7,865	7.2	62	73	Raton	7,282	6,584	-9.6	108	110
Cuba	590	632	7.1	73	16	Red River	484	485	0.2	5	4
Deming	14,116	15,277	8.2	408	258	Reserve	387	326	-15.8	6	9
Des Moines	177	147	-16.9	1	1	Rio Rancho	51,765	75,978	46.8	1,128	525
Dexter	1,235	1,244	0.7	81	20	Roswell	45,293	45,569	0.6	967	552
Dora	130	129	-0.8	2	3	Roy	304	218	-28.3	2	5
Eagle Nest	306	278	-9.2	6	2	Ruidoso	7,698	8,899	15.6	45	85
Edgewood	1,893	1,994	5.3	127	65	Ruidoso Downs	1,824	2,651	45.3	109	21
Elida	183	180	-1.6	2	6	San Jon	306	239	-21.9	1	3
Encino	94	89	-5.3	10	0	Santa Clara	1,944	1,826	-6.1	20	24
Espanola	9,688	9,549	-1.4	296	117	Santa Fe	62,203	73,199	17.7	1,494	803
Estancia	1,584	1,529	-3.5	31	12	Santa Rosa	2,744	2,534	-7.7	33	28
Eunice	2,562	2,654	3.6	53	24	San Ysidro	238	254	6.7	2	1
Farmington	37,844	42,425	12.1	1,021	398	Silver City	10,545	9,977	-5.4	230	196
Floyd	78	77	-1.3	0	0	Socorro city	8,877	8,518	-4.0	147	92
Folsom	75	62	-17.3	2	2	Springer	1,285	1,164	-9.4	23	12
Fort Sumner	1,249	983	-21.3	16	23	Sunland Park	13,309	14,225	6.9	346	37
Gallup	20,209	18,802	-7.0	448	166	Taos	4,700	5,265	12.0	164	101
Grady	98	95	-3.1	1	3	Tatum	683	722	5.7	19	11
Grants	8,806	8,876	0.8	182	92	Texico	1,065	1,035	-2.8	25	8
Grenville	25	21	-16.0	0	0	Tijeras	474	495	4.4	44	54
Hagerman	1,168	1,174	0.5	33	18	Truth or Consequences	7,289	6,689	-8.2	54	152
Hatch	1,673	1,644	-1.7	69	16	Tucumcari	5,989	5,123	-14.5	85	72
Hobbs	28,657	29,602	3.3	823	302	Tularosa	2,864	2,865	0.0	57	43
Hope	107	105	-1.9	5	1	Vaughn	539	463	-14.1	2	5
House	72	61	-15.3	1	0	Virden	143	112	-21.7	1	3
Hurley	1,464	1,366	-6.7	19	21	Wagon Mound	369	345	-6.5	5	8
Jal	1,996	2,062	3.3	29	22	Willard	240	247	2.9	5	1
Jemez Springs	375	396	5.6	3	9	Williamsburg	527	472	-10.4	4	14

Census Population Source: U.S. Census Bureau

Population Estimates Source: Bureau of Business and Economic Research (BBER), University of New Mexico. <http://www.unm.edu/~bber/>.

Demographic and Socioeconomic Status Section

Table D-10
Summary of Trends in New Mexico Health Statistics
New Mexico Residents, 1960-2007

Year	Population	Births	Deaths	Maternal Mortality		Infant Mortality				Fetal Mortality	
				Number	Rate	All Infants Deaths	Mortality Rate	Neonatal Rate	Post-neonatal Rate	Number	Rate
2007	2,053,922	30,605	15,400	4	13.1	188	6.1	3.8	2.4	82	2.7
2006	2,010,787	29,918	15,231	1	3.3	170	5.7	3.6	2.1	69	2.3
2005	1,968,347	28,822	14,866	2	6.9	175	6.1	3.6	2.5	84	2.9
2004	1,929,713	28,355	14,197	4	14.1	178	6.3	3.2	2.2	78	2.7
2003	1,899,846	27,799	14,493	4	14.4	150	5.4	3.2	2.2	89	3.2
2002	1,876,287	27,708	14,114	6	21.7	168	6.1	4.2	1.9	89	3.2
2001	1,852,740	27,101	14,016	8	29.5	174	6.4	4.0	2.4	64	2.4
2000	1,825,888	27,206	13,384	8	29.4	180	6.6	3.7	2.9	104	3.8
1999	1,808,082	27,133	13,433	5	18.4	185	6.8	3.9	2.9	90	3.3
1998	1,793,484	27,294	12,858	3	11.0	194	7.1	4.3	2.8	84	3.1
1997	1,774,839	26,844	12,613	1	3.7	165	6.1	3.3	2.9	103	3.8
1996	1,752,326	27,216	12,456	3	11.0	169	6.2	3.9	2.4	84	3.1
1995	1,720,394	26,914	12,500	3	11.1	162	6.0	4.0	2.0	106	3.9
1994	1,682,398	27,585	12,106	5	18.1	228	8.3	4.9	3.4	108	3.9
1993	1,636,453	27,831	11,689	2	7.2	233	8.4	4.6	3.8	108	3.9
1992	1,595,442	27,910	11,130	2	7.2	209	7.5	3.9	3.6	104	3.7
1991	1,555,305	27,783	11,225	3	10.8	224	8.1	4.9	3.1	100	3.6
1990	1,521,574	27,318	10,549	9	32.9	243	8.9	5.2	3.7	123	4.5
1989	1,503,901	27,265	10,473	5	18.3	232	8.5	5.4	3.1	111	4.1
1988	1,490,336	26,935	10,381	2	7.4	268	9.9	6.1	3.9	147	5.4
1987	1,478,519	27,246	10,324	1	3.7	218	8.0	5.0	3.0	121	4.4
1986	1,462,728	27,281	10,007	3	11.0	254	9.3	5.4	4.0	111	4.1
1985	1,438,360	27,449	9,637	3	10.9	291	10.6	6.2	4.4	157	5.7
1984	1,416,719	27,350	9,504	5	18.3	263	9.6	6.0	3.6	186	6.8
1983	1,394,362	27,508	9,138	3	10.9	274	10.0	5.5	4.5	192	6.9
1982	1,363,822	27,630	9,186	4	14.5	316	11.4	6.7	4.7	202	7.3
1981	1,332,747	26,565	8,668	2	7.5	256	9.6	5.7	3.9	180	6.7
1980	1,303,303	26,589	9,032	3	11.3	293	11.0	7.0	4.0	204	7.6
1979	1,283,000	24,821	8,617	10	40.3	352	14.2	8.8	5.4	206	8.2
1978	1,254,000	23,907	8,331	2	8.4	330	13.8	8.7	5.1	199	8.3
1977	1,227,000	23,100	8,223	2	8.7	333	14.4	9.8	4.6	201	8.6
1976	1,196,000	22,180	8,204	3	13.5	347	15.6	10.0	5.6	196	8.8
1975	1,164,000	21,078	8,003	2	9.5	357	16.9	11.0	6.0	169	8.0
1974	1,130,000	21,339	8,029	10	46.9	391	18.3	12.2	6.1	184	8.5
1973	1,104,000	20,852	8,139	2	9.6	421	20.2	12.9	7.3	211	10.0
1972	1,078,000	20,813	7,877	5	24.0	403	19.4	14.4	5.0	248	11.8
1971	1,053,000	22,205	7,638	5	22.5	460	20.7	15.0	5.8	254	11.3
1970	1,017,055	22,004	7,411	6	27.3	463	21.0	15.0	6.0	245	11.0
1969	1,011,000	21,543	7,180	5	23.2	508	23.6	16.1	7.5	266	12.2
1968	994,000	20,346	7,128	7	34.4	487	23.9	16.8	7.2	264	12.8
1967	1,000,000	21,243	6,897	4	18.8	527	24.8	16.1	8.7	249	11.6
1966	1,007,000	22,363	6,971	9	40.2	601	26.9	16.7	10.2	278	12.3
1965	1,012,000	24,352	6,801	10	41.1	657	27.0	17.8	9.2	326	13.2
1964	1,006,000	26,862	6,902	14	52.1	781	29.1	19.2	9.9	349	12.8
1963	989,000	27,820	6,837	10	35.9	848	30.5	20.0	10.5	371	13.2
1962	979,000	29,226	6,507	16	54.7	877	30.0	19.5	10.5	409	13.8
1961	965,000	30,009	6,344	19	63.3	880	29.3	20.0	9.3	370	12.2
1960	951,023	30,747	6,503	13	42.3	1,022	33.2	20.2	13.0	414	13.3

Maternal mortality rates per 100,000 live births, infant mortality rates per 1,000 live births and fetal mortality rates per 1,000 live births + fetal deaths.
 Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 Population Source: 2000-2007 Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. Population Source 1960 to 1999: U.S. Census Bureau. Population is based on April 1 Census enumeration for Census years. Years prior to 2000 are based on revised estimates. Estimates prior to 1980 are rounded to thousands.

Mortality Section



MORTALITY SECTION

ALL CAUSES


In 2007, 15,400 New Mexico resident deaths were reported. The age-adjusted death rate was 902.2 deaths per 100,000 standard population (Table M-1). (**Important Note:** The University of New Mexico's Bureau of Business and Economic Research (BBER) updated its methodology for estimating intercensal populations beginning with the 2007 populations. For **death rates** calculated from BBER population estimates, any rate differences from previous years and 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators. For more information, please see the *Technical Appendix.*)

Figure M-1 presents the distribution of deaths by age group and sex. The majority (67.8%) of the 2007 deaths were to individuals age 65 or older (60.9% of male deaths and 75.9% of female deaths). A larger percentage of males died between ages 15 and 64 (37.2%) than did females (22.5%). Females were much more likely to have died at ages 85 or older (33.6%) compared to males (18.7%). For residents under age 15, the distribution of deaths for both males and females was similar.

Figure M-2 shows the age-adjusted death rates by sex for each of New Mexico's racial/ethnic groups. Age-adjusted death rates allow for comparisons between time periods and geographic regions with different population age distributions. (See the *Technical Appendix.*)

In 2007, the overall age-adjusted death rate for New Mexico males was 1,014.6 per 100,000 population compared to 798.2 for females. The age-adjusted death rate of American Indian or Alaska Native males was higher than any other group in the state (1,074.7). The rate for Hispanic males (1,035.5) was higher than the overall age-adjusted death rate for New Mexico males. The rates for White males (972.5), Black males (1,000.5) and Asian or Pacific Islander males (659.3) were lower than the overall age-adjusted death rate for New Mexico males.

Figure M-1
Percent Deaths by Age Group and Sex
New Mexico, 2007




Among New Mexican females, the death rates for Hispanic females (800.0) and White females (801.3) were almost identical. Black females (601.8) had the lowest age-adjusted death rate among all groups in the state.

Table M-1 presents the 2007 death numbers and rates by age group and sex for each racial/ethnic group in New Mexico. 47.5% of American Indian or Alaska Native deaths were to residents ages 65 or older compared to 61.1% and 74.7% for Hispanics and Whites, respectively.

Table M-2 shows the numbers of deaths by sex and age group for each county in 2007. Union (83.3%) and Los Alamos (81.7%) counties had the highest percentages of deaths of persons aged 65 and older. For the young adult age group, which is 15-24 years, Guadalupe (6.8%) and Rio Arriba (5.4%) counties had the largest percentages of youth deaths.

Age-specific death rates by county, health region and sex in 2007 are presented in Table M-3.

Figure M-2
Death Rates by Sex and Race/Ethnicity
New Mexico, 2007




Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

AI or AN = American Indian or Alaska Native, API = Asian or Pacific Islander, AA=African American. The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, if any, are included in the "All Races" category.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. Released 2009. <http://www.unm.edu/~bber/>.

Table M-13 also shows that males in 2007 had a 12.5% higher age-adjusted death rate than that for both sexes together while females had an age-adjusted death rate that was 11.5% lower than the age-adjusted death rate for both sexes.

Figure M-3
Death Rates
New Mexico, 1998-2007,
United States, 1998-2006



Age-adjusted death rates are the number of deaths per 100,000 U.S. standard population.

Population Source: Bureau of Business and Economic Research (BBER), Population Estimates, University of New Mexico. Released 2009.

<http://www.unm.edu/~bber/>.

U.S. Data Source: CDC, National Center for Health Statistics. NVRS, Vol 57, No. 14, April 17, 2009. * U.S. 2007 data is not available at time of publication.

American Indian and Alaska Native and Hispanic males had the highest age-adjusted death rates in 2007 (1,074.4 and 1,035.5, respectively). The Asian or Pacific Islander population had the lowest death rates. Due to the small number of deaths and the small population for Asian or Pacific Islanders, death rates for this race/ethnicity fluctuate greatly from year-to-year and should be interpreted with caution.

Table M-13 shows that for the period 2003-2007, the number of New Mexico deaths ranged from 14,197 in 2004 to a high of 15,400 in 2007. In 2007, there were 8,233 deaths to males, accounting for 53.5% of all deaths; 7,167 deaths were to females, 46.5% of the total. In 2007, New Mexico's greatest difference in percents of deaths by sex was among the Asian/Pacific Islander population with 60.0% of deaths to females and 40.0% to males.

LEADING CAUSES OF DEATH

For New Mexico, the 2007 leading causes of death were:

- 1) Heart Disease
- 2) Malignant Neoplasms (Cancer)
- 3) Accidents (Unintentional Injuries)
- 4) Chronic Lower Respiratory Diseases
- 5) Cerebrovascular Diseases (Stroke)
- 6) Diabetes Mellitus
- 7) Intentional Self-harm (Suicide)
- 8) Chronic Liver Disease and Cirrhosis
- 9) Influenza and Pneumonia
- 10) Alzheimer's Disease

For the United States, the 2006 leading causes of death were:

- 1) Heart Disease
- 2) Malignant Neoplasms (Cancer)
- 3) Cerebrovascular Diseases (Stroke)
- 4) Chronic Lower Respiratory Diseases
- 5) Accidents (Unintentional Injuries)
- 6) Diabetes Mellitus
- 7) Alzheimer's Disease
- 8) Influenza and Pneumonia
- 9) Nephritis, Nephritic Syndrome and Nephrosis (Kidney Disease)
- 10) Septicemia

New Mexico's leading causes of death in 2007, along with the corresponding percents of total deaths, are presented in rank order in Figure M-4. Heart Disease, the leading cause of death, accounted for 21.2% of all deaths in the state, followed by Malignant Neoplasms (Cancer) at 20.8% and Accidents (Unintentional Injuries) at 8.3%.


Among New Mexico counties shown in Table M-4, Guadalupe, McKinley and Torrance counties had the highest age-adjusted death rates for all causes. Los Alamos, Union and Lincoln counties had the lowest age-adjusted death rates for all causes.

Figure M-5 illustrates changes in age-adjusted death rates, by cause, from 1998 to 2007. With the exception of the rates for 2007, this figure shows that the death rates for Heart Disease and Malignant

Neoplasms (Cancer) declined, while death rates for Accidents (Unintentional Injuries) increased.


For children ages 1-4 (Table M-6), the three leading causes of death were Accidents (Unintentional Injuries), Congenital Malformations and Assault (Homicide). For the 5-14 age group (Table M-7), the three leading causes were Accidents (Unintentional Injuries), Intentional Self-harm (Suicide) and Malignant Neoplasms (Cancer). Suicide was the second leading cause of death, after Unintentional Injuries, for both the 15-24 (Table M-8) and the 25-44 (Table M-9) age groups, overall.

Figure M-4
Leading Causes of Death
New Mexico, 2007 and United States, 2006



A historical pattern of the leading causes of death can be derived from Table M-14. With the exception of the 2007 rates that may be influenced by the population denominators, all of the age-adjusted death rates for the leading causes of death except for Accidents (Unintentional Injuries) showed declines from 1999-2006. During this period, rates of Heart Disease fell 21.0%, Cancer fell by 10.0% and Cerebrovascular Diseases (Stroke) fell by 65.2%. Compared to 1999, the rate for Accidents (Unintentional Injuries) increased 19.7%. From 2006 to 2007, Suicide and Chronic Liver Disease replaced Alzheimer's Disease and Influenza/Pneumonia as the seventh and eighth ranked causes of death.

Figure M-5
Death Rates by Cause
New Mexico, 1998-2007



Age-adjusted death rates are the number of deaths per 100,000 U.S. standard population. Age-adjusting allows for comparison across time periods. To adjust for differences in coding causes of death prior to 1999, comparability ratios have been applied to 1997-1998 rates. For further explanation of comparability ratios, please see Technical Appendix.

For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Population Source: 2000-2007: Bureau of Business and Economic Research (BBER), University of New Mexico. <http://www.unm.edu/~bber/>; 1998-1999: U.S. Census Bureau in collaboration with NCHS.

SELECTED CAUSES

Tables M-15 through M-18 contain the New Mexico death numbers and rates for 39-selected causes of death by age group, sex, health region and race/ethnicity for 2007.

Causes of death associated with chronic diseases in the 65+ years of age showed the largest number of deaths and included Malignant Neoplasms (Cancer), Major Cardiovascular Disease and Chronic Lower Respiratory Diseases.

The 25-64 age group had the largest numbers of deaths from Motor Vehicle Accidents, Intentional Self-harm (Suicide) and Assault (Homicide) (Table M-15).

Death rates for males exceeded those for females for Major Cardiovascular Diseases (16.3%) and Cancer (40.3%). The male death rates were higher than those for females for the following causes: Suicide (370% higher); Homicide (237.8% higher); and Motor Vehicle Accidents (146.6% higher). For Alzheimer's Disease, the death rate for males was 35.0% lower than the female rate.

INJURY DEATHS AND ACOHOL- AND DRUG-INDUCED DEATHS

Unintentional Injuries (Accidents) accounted for 5.0% of all deaths in the United States in 2006 and 8.3% of New Mexico deaths in 2007. When all injury (both intentional and unintentional) deaths were combined (see Figure M-6), Motor Vehicle Accident deaths (18.9%) and Intentional Self-harm (Suicide) deaths (20.6%) constituted similar percents of deaths. The bulk of injury deaths (46.8%) were caused by non-Motor Vehicle Accidents. Assault (Homicide) accounted for 8.4% of injury deaths. The remaining 5.4% of injury deaths were due to Other Injuries.

For counties with more than 20 injury deaths (and therefore with more statistically reliable rates), Rio Arriba, Taos and McKinley Counties had the highest death rates due to total injuries (170.8, 168.2 and 151.2, respectively). These rates were all more than twice the 2006 national rate of 58.8 deaths per 100,000 standard population. In contrast, Los Alamos County had the lowest age-adjusted injury

Mortality Section

death rate at 59.6, which was slightly higher than the United States 2006 rate (Table M-20).

Firearm injury deaths, which are defined in the *Technical Appendix*, accounted for 15.0% of all New Mexico injury deaths in 2007 (Tables M-19 and M-21). Figure M-7 shows the breakdown of firearm injury deaths. Most firearm injury deaths (63.2%) were Suicides, 27.5% were Homicides, 2.7% were Legal Intervention and 3.1% were Accidents (Unintentional Injuries).

Of those counties with over 7 firearm injury deaths during 2007, Taos and Rio Arriba counties had the highest firearm injury death rates, 31.3 and 20.8, respectively, per 100,000 standard population (Tables M-21 and M-22).

Only Bernalillo and Dona Ana Counties had more than 20 firearm injury deaths in 2007, giving the rates for these counties adequate levels of statistical reliability.

Figure M-6
Percent of Injury Deaths by Type
New Mexico, 2007





Figure M-7
Percent of Firearm Injury Deaths by Manner
New Mexico, 2007



New Mexico's alcohol-induced death rate was 19.0 deaths per 100,000 standard population. Of those counties with more than 20 alcohol-induced deaths, McKinley and Rio Arriba Counties had the highest rates at 60.8 and 56.1 deaths per 100,000 standard population, respectively (Table M-22).

Approximately 2.0% of all injury deaths in the state occurred at work, compared to 2.9% for the United States (Table M-21). The total number of injury at work deaths in New Mexico was 39 in 2007, with 36 of these occurring among males.

MATERNAL MORTALITY

Maternal mortality includes deaths that were the result of, or aggravated by, pregnancy or pregnancy management, and occurred within 42 days of pregnancy and excludes all external injury deaths. (See the *Technical Appendix*.)

For the period 2000-2003, the maternal death rate was 21.9 in New Mexico. The maternal mortality rate for 2003-2007 was 9.3.

During both time periods, mothers ages 35-49 had the highest maternal mortality rates.

The numbers of maternal deaths are so few each year that even with reporting 4-year totals for the death counts, the rates presented in Table M-23 should be interpreted with caution.

INFANT MORTALITY


From Table M-25, in 2007, 188 infant deaths to New Mexico residents were reported. Infant deaths are defined as deaths before the age of one. The 2007 infant mortality rate was 6.1 infant deaths per 1,000 live births. This rate was an increase of 7.0% from the 2006 New Mexico rate, which was 5.7. The 2007 New Mexico rate was lower than the 2006 U.S. rate of 6.7.

A historical perspective is provided in Figure M-8. In 1930, New Mexico's infant mortality rate (145.4) was 125.1% higher than the national rate. By the early 1970s, however, New Mexico had achieved parity with the United States and by the 1980s the rate had dropped below the United States rate. With the exception of one year (1994), the state's infant mortality rate has remained below the national rate every year since 1980. The historical decline in infant mortality is due to the role of public health outreach, improved education about pregnancy and childbirth, and expanded medical services.

Infant deaths may be broken down into two age categories: neonatal deaths (less than 28 days old) and postneonatal deaths (at least 28 days but less than one year old). 61.7% of infant deaths in 2007 occurred during the neonatal period and 38.3% occurred during the postneonatal period.

Figure M-9 shows the 2007 neonatal and postneonatal mortality rates by race/ethnicity in New Mexico. Black infants had the highest neonatal and postneonatal mortality rates in the state (9.4 and 4.7, respectively), followed by American Indian or Alaska Native infants with rates of 3.3 and 4.3 for the two age categories. Asian or Pacific Islander, Hispanic and White infants had lower rates for both the neonatal and postneonatal periods. Due to small numbers, the Asian or Pacific Islander rates may not be statistically reliable and should be interpreted with caution.


Figure M-8
Infant Mortality Rates
New Mexico, 1930-2007
United States, 1930-2006



Males had a higher infant mortality rate than females in 2007: 7.1 versus 5.1 (Table M-25).



Figure M-10 compares the state and national neonatal and postneonatal death rates from 1991 onward. During this time period, the state's neonatal infant mortality rate, like the overall infant mortality rate, remained below the United States rate, while state and national postneonatal rates fluctuated but remained close to one another.

Figure M-9
Infant Mortality Rates by Race/ Ethnicity
New Mexico, 2007



Of the 188 New Mexico infant deaths in 2007, 40 were caused by congenital malformations (birth defects); 39 were caused by disorders related to length of gestation or birth weight; and 21 were caused by maternal factors and complications of pregnancy, labor and delivery. There were 14 deaths attributed to Sudden Infant Death Syndrome (SIDS) (Table M-26).

Figure M-10
Infant Mortality Rates,
New Mexico, 1991-2007
United States 1991-2006




Postneonatal mortality rates are the numbers of infant deaths 28 days of age to under 1 year per 1,000 births. Neonatal mortality rates are the numbers of infant deaths under 28 days of age per 1,000 births. For further information, please see Technical Appendix. U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol. 57, No. 14. 2007 U.S. data not available at time of publication.

DISPOSITION OF REMAINS

Figure M-11 shows the distribution of the method of disposition. Table M-27 shows disposition of remains over a ten-year period. 2007 was the second year (2006 was the first) for which a greater percentage of individuals who died in the state were cremated (49.2%) than were buried (45.0%). The percent of burials has declined from 49.2% in 1998 to 45.0% in 2007; whereas, cremations have increased from 32.2% in 1998 to 49.2% in 2007.

Figure M-11
Percent Distribution by Method of
Disposition
New Mexico Occurrence, 2007



Mortality Section

Table M-1
Numbers and Rates of Deaths by Age, Race/Ethnicity and Sex
New Mexico, 2007, and United States, 2006

Race / Ethnicity and Sex	Total		Under 1 Year		1 to 4 Years		5 to 14 Years		15 to 24 Years		25 to 34 Years	
	Number	Age-adjusted Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
New Mexico												
All Races												
Both Sexes	15,400	902.2	188	712.4	47	47.4	42	15.0	323	101.5	410	172.3
Male	8,233	1,014.6	112	825.7	22	43.8	24	16.9	249	153.5	281	234.9
Female	7,167	798.2	76	592.6	25	51.1	18	13.1	74	47.4	129	109.1
American Indian or Alaska Native												
Both Sexes	1,142	917.8	30	848.7	12	87.8	10	23.9	69	151.2	94	332.1
Male	657	1,074.7	18	997.2	6	88.3	7	33.6	50	223.8	71	519.7
Female	485	781.9	12	693.6	6	87.4	3	14.3	19	81.6	23	157.1
Asian or Pacific Islander												
Both Sexes	90	662.3	1	220.3	0	0.0	0	0.0	6	111.5	1	17.5
Male	36	659.3	1	414.9	0	0.0	0	0.0	4	138.0	0	0.0
Female	54	658.0	0	0.0	0	0.0	0	0.0	2	80.5	1	34.2
Black or African American												
Both Sexes	248	794.3	9	1,296.8	2	76.0	1	10.2	13	123.7	6	76.5
Male	163	1,000.5	6	1,719.2	0	0.0	1	20.3	11	201.5	5	122.2
Female	85	601.8	3	869.6	2	151.1	0	0.0	2	39.6	1	26.6
Hispanic												
Both Sexes	4,924	914.0	105	766.8	23	45.0	21	15.3	150	102.2	197	173.4
Male	2,788	1,035.5	59	839.7	10	38.5	11	15.8	122	164.1	141	243.3
Female	2,136	800.0	46	690.0	13	51.8	10	14.8	28	38.6	56	100.5
White												
Both Sexes	8,954	883.5	42	524.2	10	33.1	10	11.6	84	76.4	111	134.7
Male	4,553	972.5	28	675.8	6	39.2	5	11.3	61	106.8	63	153.0
Female	4,401	801.3	14	361.9	4	26.9	5	11.9	23	43.6	48	116.3
United States												
All Races												
Both Sexes	2,426,264	776.5	28,527	690.7	4,631	28.4	6,149	15.2	34,887	82.2	42,952	106.3
Male	1,201,942	924.8	15,980	756.3	2,541	30.5	3,625	17.6	26,070	119.3	30,188	146.8
Female	1,224,322	657.8	12,547	622.0	2,090	26.3	2,524	12.8	8,817	42.8	12,764	64.3

Age-specific death rates are the numbers of deaths per 100,000 population in specified age group.

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

Notes: The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races or ages, if any, are included in the "All Races" and "All Ages" categories. Numbers may not add to total due to unknown.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-1 (continued)

35 to 44 Years		45 to 54 Years		55 to 64 Years		65 to 74 Years		75 to 84 Years		85 Plus Years		Race / Ethnicity and Sex
Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	
New Mexico												
All Races												
652	216.7	1,357	431.8	1,934	916.6	2,521	1,703.2	3,976	3,896.7	3,948	25,117.7	Both Sexes
462	309.6	887	580.2	1,186	1,165.8	1,456	2,119.5	2,015	4,671.9	1,538	23,563.7	Male
190	125.3	470	291.1	748	684.6	1,065	1,342.7	1,961	3,329.1	2,410	26,221.3	Female
American Indian or Alaska Native												
115	338.1	140	499.8	129	810.5	190	1,942.1	179	3,421.9	174	23,015.9	Both Sexes
86	519.8	89	684.8	83	1,131.7	93	2,189.3	84	3,973.5	70	22,653.7	Male
29	166.0	51	339.7	46	536.0	97	1,752.5	95	3,047.8	104	23,266.2	Female
Asian or Pacific Islander												
3	49.4	10	200.8	15	533.2	20	1,238.4	16	1,891.3	18	25,352.1	Both Sexes
2	71.1	5	231.7	6	526.3	6	931.7	4	1,257.9	8	27,586.2	Male
1	30.6	5	177.2	9	538.0	14	1,441.8	12	2,272.7	10	23,809.5	Female
Black or African American												
14	169.5	21	298.7	45	1,245.8	39	1,540.9	62	4,172.3	36	17,910.4	Both Sexes
11	245.7	17	450.3	33	1,742.3	27	2,195.1	42	7,304.3	10	13,157.9	Male
3	79.3	4	122.9	12	698.5	12	922.4	20	2,195.4	26	20,800.0	Female
Hispanic												
279	219.6	525	468.1	616	935.6	818	1,767.7	1,161	3,933.3	1,029	24,987.9	Both Sexes
204	321.9	360	654.0	406	1,277.5	479	2,280.8	596	4,702.9	400	22,586.1	Male
75	117.7	165	288.9	210	616.6	339	1,341.3	565	3,354.3	629	26,800.2	Female
White												
238	189.7	649	400.3	1,117	909.6	1,447	1,647.8	2,555	3,933.6	2,689	25,435.1	Both Sexes
156	251.6	406	514.6	649	1,089.3	844	2,030.1	1,286	4,684.9	1,048	24,136.3	Male
82	129.3	243	292.0	468	740.2	603	1,304.1	1,269	3,383.6	1,641	26,340.3	Female
United States												
All Races												
83,043	190.2	185,031	427.5	281,401	890.9	390,093	2,062.1	667,338	5,115.0	701,992	13,253.1	Both Sexes
52,150	238.7	115,172	541.0	168,987	1,110.0	218,165	2,516.2	327,315	6,177.7	241,578	14,309.1	Male
30,893	141.6	69,859	317.7	112,414	687.0	171,928	1,677.9	340,023	4,388.3	460,414	12,759.0	Female

Age-specific death rates are the numbers of deaths per 100,000 population in specified age group.

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

Notes: The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races or ages, if any, are included in the "All Races" and "All Ages" categories. Numbers may not add to total due to unknown.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Mortality Section

Table M-2
Number of Deaths by Age, Sex, County and Health Region
New Mexico, 2007, and United States, 2006

Resident County	All Ages			Under 1 Year			1 to 4 Years			5 to 14 Years		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	2,426,264	1,201,942	1,224,322	28,527	15,980	12,547	4,631	2,541	2,090	6,149	3,625	2,524
New Mexico	15,400	8,233	7,167	188	112	76	47	22	25	42	24	18
Bernalillo	4,829	2,586	2,243	60	36	24	17	10	7	12	5	7
Catron	43	28	15	0	0	0	0	0	0	0	0	0
Chaves	594	294	300	4	3	1	0	0	0	0	0	0
Cibola	228	132	96	7	3	4	1	1	0	1	0	1
Colfax	142	74	68	0	0	0	0	0	0	0	0	0
Curry	344	169	175	7	3	4	0	0	0	1	0	1
De Baca	25	11	14	0	0	0	0	0	0	1	0	1
Doña Ana	1,287	678	609	10	6	4	4	1	3	5	3	2
Eddy	498	245	253	8	4	4	1	1	0	0	0	0
Grant	351	186	165	2	0	2	0	0	0	1	1	0
Guadalupe	44	21	23	0	0	0	0	0	0	0	0	0
Harding	9	6	3	0	0	0	0	0	0	0	0	0
Hidalgo	55	26	29	0	0	0	0	0	0	0	0	0
Lea	471	243	228	9	6	3	2	1	1	2	2	0
Lincoln	181	108	73	3	1	2	2	1	1	0	0	0
Los Alamos	115	51	64	0	0	0	1	1	0	0	0	0
Luna	267	163	104	2	1	1	0	0	0	2	2	0
McKinley	495	291	204	12	10	2	4	2	2	4	3	1
Mora	47	25	22	1	1	0	0	0	0	0	0	0
Otero	481	229	252	5	3	2	0	0	0	1	1	0
Quay	98	55	43	0	0	0	0	0	0	0	0	0
Rio Arriba	373	223	150	3	1	2	1	0	1	1	1	0
Roosevelt	158	76	82	4	2	2	0	0	0	1	1	0
San Juan	867	485	382	18	12	6	5	1	4	5	2	3
San Miguel	266	152	114	2	2	0	0	0	0	0	0	0
Sandoval	809	434	375	8	6	2	2	1	1	0	0	0
Santa Fe	943	489	454	13	7	6	3	1	2	2	1	1
Sierra	219	130	89	0	0	0	1	0	1	0	0	0
Socorro	151	81	70	4	3	1	0	0	0	0	0	0
Taos	269	139	130	2	1	1	0	0	0	1	1	0
Torrance	127	69	58	0	0	0	0	0	0	0	0	0
Union	30	14	16	0	0	0	0	0	0	0	0	0
Valencia	578	317	261	4	1	3	2	1	1	2	1	1
Health Region												
Northwest	2,977	1,659	1,318	49	32	17	14	6	8	12	6	6
Northeast	2,229	1,188	1,041	21	12	9	5	2	3	4	3	1
Bernalillo County	4,829	2,586	2,243	60	36	24	17	10	7	12	5	7
Southeast	2,197	1,099	1,098	32	18	14	3	2	1	5	3	2
Southwest	3,162	1,698	1,464	26	14	12	7	2	5	9	7	2

Unknown ages, sexes or counties are included in the total categories. Numbers may not add to total due to unknowns.

For counties included in state health regions, please see Technical Appendix.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-2 (continued)

15 to 24 Years			25 to 34 Years			35 to 44 Years			45 to 54 Years			Resident County
Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	
34,887	26,070	8,817	42,952	30,188	12,764	83,043	52,150	30,893	185,031	115,172	69,859	United States
323	249	74	410	281	129	652	462	190	1,357	887	470	New Mexico
79	66	13	107	72	35	212	146	66	438	296	142	Bernalillo
2	2	0	0	0	0	1	1	0	3	2	1	Catron
12	8	4	16	10	6	19	10	9	52	30	22	Chaves
6	6	0	7	6	1	9	6	3	26	17	9	Cibola
3	2	1	3	2	1	3	2	1	9	7	2	Colfax
1	1	0	11	9	2	16	11	5	23	14	9	Curry
0	0	0	0	0	0	3	3	0	1	1	0	De Baca
16	13	3	29	16	13	39	29	10	94	60	34	Doña Ana
6	6	0	7	5	2	14	10	4	36	26	10	Eddy
7	7	0	7	3	4	7	4	3	24	16	8	Grant
3	1	2	0	0	0	1	1	0	3	1	2	Guadalupe
0	0	0	1	1	0	0	0	0	0	0	0	Harding
1	1	0	1	1	0	4	3	1	2	1	1	Hidalgo
15	12	3	12	9	3	15	7	8	36	20	16	Lea
5	4	1	3	3	0	7	5	2	20	11	9	Lincoln
3	3	0	0	0	0	3	3	0	6	1	5	Los Alamos
6	3	3	4	3	1	4	2	2	24	14	10	Luna
23	18	5	44	36	8	39	29	10	49	28	21	McKinley
1	0	1	0	0	0	0	0	0	0	0	0	Mora
8	5	3	5	5	0	19	14	5	41	25	16	Otero
2	1	1	1	1	0	3	1	2	14	10	4	Quay
20	17	3	17	12	5	28	21	7	47	32	15	Rio Arriba
1	0	1	6	6	0	4	1	3	10	6	4	Roosevelt
39	28	11	40	25	15	49	39	10	95	63	32	San Juan
8	8	0	7	4	3	10	10	0	28	19	9	San Miguel
11	7	4	15	11	4	35	27	8	76	49	27	Sandoval
18	13	5	17	10	7	44	33	11	74	50	24	Santa Fe
1	0	1	0	0	0	3	3	0	19	12	7	Sierra
2	2	0	8	4	4	8	4	4	15	12	3	Socorro
5	4	1	11	9	2	17	11	6	26	17	9	Taos
1	1	0	1	1	0	6	4	2	11	8	3	Torrance
1	1	0	1	1	0	0	0	0	2	1	1	Union
17	9	8	27	15	12	29	21	8	52	37	15	Valencia
												Health Region
96	68	28	133	93	40	161	122	39	298	194	104	Northwest
62	49	13	56	38	18	106	81	25	195	128	67	Northeast
79	66	13	107	72	35	212	146	66	438	296	142	Bernalillo County
37	28	9	54	41	13	74	43	31	172	107	65	Southeast
49	38	11	58	36	22	98	69	29	253	161	92	Southwest

Unknown ages, sexes or counties are included in the total categories. Numbers may not add to total due to unknowns.

For counties included in state health regions, please see Technical Appendix.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Mortality Section

Table M-2 (continued)

Resident County	55 to 64 Years			65 to 74 Years			75 to 84 Years			85 Plus Years		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	281,401	168,987	112,414	390,093	218,165	171,928	667,338	327,315	340,023	701,992	241,578	460,414
New Mexico	1,934	1,186	748	2,521	1,456	1,065	3,976	2,015	1,961	3,948	1,538	2,410
Bernalillo	624	405	219	671	390	281	1,252	623	629	1,357	537	820
Catron	7	4	3	10	8	2	13	6	7	7	5	2
Chaves	54	27	27	80	52	28	173	86	87	184	68	116
Cibola	24	12	12	56	38	18	48	22	26	43	21	22
Colfax	17	11	6	26	14	12	36	19	17	45	17	28
Curry	48	26	22	63	35	28	76	34	42	98	36	62
De Baca	3	1	2	2	1	1	4	4	0	11	1	10
Doña Ana	163	91	72	229	126	103	377	202	175	321	131	190
Eddy	52	32	20	88	48	40	147	70	77	139	43	96
Grant	37	19	18	59	37	22	100	57	43	107	42	65
Guadalupe	8	5	3	4	1	3	12	9	3	13	3	10
Harding	2	2	0	1	1	0	3	1	2	2	1	1
Hidalgo	5	3	2	14	8	6	15	7	8	13	2	11
Lea	62	37	25	84	52	32	128	62	66	106	35	71
Lincoln	28	18	10	45	29	16	39	20	19	29	16	13
Los Alamos	8	4	4	12	2	10	42	20	22	40	17	23
Luna	26	18	8	61	42	19	73	43	30	65	35	30
McKinley	57	40	17	79	48	31	100	47	53	84	30	54
Mora	11	6	5	12	7	5	9	5	4	13	6	7
Otero	62	36	26	104	48	56	134	60	74	102	32	70
Quay	13	6	7	19	14	5	21	10	11	25	12	13
Rio Arriba	50	36	14	57	37	20	78	37	41	71	29	42
Roosevelt	12	8	4	26	12	14	39	19	20	55	21	34
San Juan	97	55	42	152	89	63	213	113	100	154	58	96
San Miguel	30	17	13	54	31	23	56	34	22	71	27	44
Sandoval	120	74	46	129	63	66	209	110	99	204	86	118
Santa Fe	122	75	47	147	85	62	217	105	112	286	109	177
Sierra	28	21	7	48	30	18	69	42	27	50	22	28
Socorro	18	11	7	19	11	8	41	21	20	36	13	23
Taos	38	19	19	41	23	18	55	24	31	72	29	43
Torrance	33	20	13	14	10	4	34	20	14	27	5	22
Union	1	1	0	6	3	3	10	3	7	9	4	5
Valencia	74	46	28	108	61	47	153	80	73	109	45	64
Health Region												
Northwest	372	227	145	524	299	225	723	372	351	594	240	354
Northeast	285	174	111	359	203	156	515	256	259	620	241	379
Bernalillo County	624	405	219	671	390	281	1,252	623	629	1,357	537	820
Southeast	246	139	107	363	215	148	591	286	305	620	217	403
Southwest	407	241	166	603	349	254	895	478	417	757	303	454

Unknown ages or sexes, if any, are included in the "All Ages" or "Both Sexes" categories. Numbers may not add to total due to unknowns.

For counties included in state health regions, please see Technical Appendix.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol. 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-3
Death Rates by Age, Sex, County and Health Region
New Mexico, 2007, and United States, 2006

Resident County	All Ages			Under 1 Year			1 to 4 Years			5 to 14 Years		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	776.5	924.8	657.8	690.7	756.3	622.0	28.4	30.5	26.3	15.2	17.6	12.8
New Mexico	902.2	1,014.6	798.2	712.4	825.7	592.6	47.4	43.8	51.1	15.0	16.9	13.1
Bernalillo	842.9	959.4	737.2	721.2	847.5	589.5	56.2	65.5	46.7	15.5	12.6	18.4
Catron	829.5	1,393.9	469.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Chaves	1,008.1	1,089.1	935.6	529.1	742.6	284.1	0.0	0.0	0.0	0.0	0.0	0.0
Cibola	994.3	1,203.1	809.5	1,866.7	1,595.7	2,139.0	68.7	130.7	0.0	26.1	0.0	51.3
Colfax	865.5	994.1	748.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Curry	927.3	1,043.8	831.4	935.8	763.4	1,126.8	0.0	0.0	0.0	15.2	0.0	30.8
De Baca	952.5	834.4	951.8	0.0	0.0	0.0	0.0	0.0	0.0	423.7	0.0	775.2
Doña Ana	838.7	933.7	755.0	302.8	354.0	248.9	34.8	17.3	52.5	17.7	20.9	14.3
Eddy	974.1	1,035.2	896.6	1,280.0	1,269.8	1,290.3	40.2	80.1	0.0	0.0	0.0	0.0
Grant	1,043.0	1,179.9	922.8	537.6	0.0	1,204.8	0.0	0.0	0.0	24.7	48.4	0.0
Guadalupe	1,723.0	1,375.2	2,047.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Harding	831.4	1,389.1	304.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	928.4	847.9	933.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Lea	941.2	1,019.9	860.0	870.4	1,125.7	598.8	52.7	51.6	53.8	25.3	49.6	0.0
Lincoln	719.0	918.1	544.5	1,395.3	980.4	1,769.9	257.1	260.4	253.8	0.0	0.0	0.0
Los Alamos	575.1	541.0	603.2	0.0	0.0	0.0	158.0	318.5	0.0	0.0	0.0	0.0
Luna	946.2	1,180.8	729.7	428.3	416.7	440.5	0.0	0.0	0.0	42.9	87.9	0.0
McKinley	1,263.4	1,422.8	1,131.0	1,054.5	1,712.3	361.0	86.4	85.5	87.4	24.0	36.1	12.0
Mora	943.4	951.6	934.5	2,127.7	3,846.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Otero	922.3	895.8	929.3	674.8	797.9	547.9	0.0	0.0	0.0	9.5	18.0	0.0
Quay	797.1	1,045.9	619.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Rio Arriba	1,122.0	1,363.5	895.3	571.4	355.9	819.7	47.7	0.0	91.6	15.2	30.7	0.0
Roosevelt	1,015.4	1,013.6	1,009.7	1,860.5	1,834.9	1,886.8	0.0	0.0	0.0	37.1	70.9	0.0
San Juan	820.6	969.1	693.1	1,124.3	1,411.8	798.9	84.1	34.1	132.9	28.8	23.0	34.8
San Miguel	1,072.1	1,276.1	889.1	657.9	1,250.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Sandoval	949.1	1,085.6	831.1	550.2	829.9	273.6	31.9	30.8	33.1	0.0	0.0	0.0
Santa Fe	902.6	960.1	845.2	804.5	840.3	766.3	51.1	33.6	69.1	11.5	11.4	11.7
Sierra	1,013.5	1,205.3	838.6	0.0	0.0	0.0	223.7	0.0	450.5	0.0	0.0	0.0
Socorro	1,166.9	1,242.8	1,082.2	1,851.9	2,702.7	952.4	0.0	0.0	0.0	0.0	0.0	0.0
Taos	979.2	1,120.8	852.8	549.5	565.0	534.8	0.0	0.0	0.0	24.8	48.5	0.0
Torrance	1,218.3	1,240.0	1,101.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Union	581.4	647.7	514.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Valencia	1,088.8	1,230.8	968.7	393.3	191.6	606.1	50.6	49.4	52.0	15.7	15.2	16.2
Health Region												
Northwest	964.7	1,117.1	829.9	877.4	1,116.1	625.5	62.9	53.1	73.2	17.3	17.1	17.6
Northeast	923.3	1,020.8	828.5	639.7	703.8	570.3	40.5	32.2	49.0	10.4	15.4	5.3
Bernalillo County	842.9	959.4	737.2	721.2	847.5	589.5	56.2	65.5	46.7	15.5	12.6	18.4
Southeast	958.9	1,042.0	876.2	917.4	997.2	831.8	22.7	29.5	15.6	14.1	16.8	11.3
Southwest	898.0	1,001.6	801.2	455.1	476.4	432.6	33.1	18.8	47.6	15.2	23.2	6.9

Age-specific death rates are the numbers of deaths per 100,000 population in specified sex and age groups.

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Unknown ages, sexes and counties, if any, are included in the "All Ages," "Both Sexes" and New Mexico categories.

For counties included in state health regions, please see Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Mortality Section

Table M-3 (continued)

Resident County	15 to 24 Years			25 to 34 Years			35 to 44 Years			45 to 54 Years		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	82.2	119.3	42.8	106.3	146.8	64.3	190.2	238.7	141.6	427.5	541.0	317.7
New Mexico	101.5	153.5	47.4	172.3	234.9	109.1	216.7	309.6	125.3	431.8	580.2	291.1
Bernalillo	87.6	145.0	29.1	135.7	180.3	90.0	211.8	292.2	131.7	425.8	596.1	266.9
Catron	445.4	778.2	0.0	0.0	0.0	0.0	257.1	632.9	0.0	399.5	520.8	272.5
Chaves	113.5	152.6	75.1	235.7	288.2	180.8	216.3	218.8	213.5	548.0	655.5	447.9
Cibola	132.9	260.8	0.0	217.3	388.9	59.6	209.0	292.5	133.0	611.2	796.6	424.5
Colfax	154.2	185.9	114.9	236.2	308.6	160.8	167.8	216.2	115.9	372.4	564.5	169.9
Curry	12.5	24.3	0.0	172.0	269.9	65.3	236.0	330.8	144.7	367.4	450.9	285.3
De Baca	0.0	0.0	0.0	0.0	0.0	0.0	993.4	2,054.8	0.0	339.0	595.2	0.0
Doña Ana	42.8	68.3	16.4	96.9	103.6	89.8	159.9	246.6	79.2	359.6	476.7	250.8
Eddy	72.2	139.8	0.0	142.1	215.8	76.7	193.9	288.1	106.7	415.1	585.6	236.2
Grant	151.6	300.8	0.0	221.0	198.2	242.0	179.3	218.7	144.6	500.1	717.5	311.4
Guadalupe	384.6	221.7	607.9	0.0	0.0	0.0	101.5	169.2	0.0	425.5	230.9	735.3
Harding	0.0	0.0	0.0	2,173.9	4,545.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	111.4	217.9	0.0	185.2	400.0	0.0	487.8	719.4	248.1	235.6	233.1	238.1
Lea	152.7	239.7	62.3	158.8	232.1	81.5	203.3	185.1	222.5	446.8	486.5	405.4
Lincoln	177.0	265.8	75.8	177.8	369.5	0.0	252.9	379.9	137.7	512.8	590.8	441.6
Los Alamos	136.7	254.0	0.0	0.0	0.0	0.0	91.9	188.9	0.0	164.8	55.2	273.1
Luna	137.7	140.2	135.4	140.7	229.9	65.1	110.5	118.0	103.8	766.8	893.4	639.8
McKinley	152.9	235.3	67.6	474.9	805.0	166.9	329.2	513.5	161.3	482.1	597.7	383.3
Mora	107.0	0.0	222.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Otero	75.0	94.3	56.0	74.3	150.0	0.0	176.7	260.8	92.9	443.5	562.4	333.4
Quay	149.5	145.3	153.8	130.7	272.5	0.0	228.0	160.3	289.0	875.0	1,282.1	487.8
Rio Arriba	309.7	510.2	96.0	321.4	447.3	191.9	417.2	618.7	211.0	676.2	932.1	426.4
Roosevelt	26.4	0.0	53.0	235.7	462.2	0.0	160.8	78.9	245.7	422.7	544.5	316.5
San Juan	168.5	246.6	93.3	300.6	410.8	207.7	248.1	389.3	102.8	489.6	649.6	329.8
San Miguel	189.3	367.8	0.0	228.1	264.2	192.9	190.4	393.5	0.0	526.0	734.4	328.9
Sandoval	50.1	61.5	37.9	114.0	167.7	60.7	186.3	292.7	83.7	407.0	532.0	285.3
Santa Fe	96.0	132.5	55.9	100.4	116.4	83.9	200.4	303.1	99.4	263.8	379.6	161.4
Sierra	60.4	0.0	129.4	0.0	0.0	0.0	178.6	365.9	0.0	1,052.0	1,348.3	764.2
Socorro	55.1	101.0	0.0	360.2	347.8	373.5	312.6	323.1	302.8	561.8	906.3	222.9
Taos	128.7	199.7	53.1	371.5	576.6	142.9	379.1	495.7	264.9	416.5	573.7	274.4
Torrance	35.2	67.1	0.0	50.0	89.4	0.0	209.7	268.8	145.7	395.1	557.1	222.6
Union	165.3	309.6	0.0	290.7	518.1	0.0	0.0	0.0	0.0	284.5	270.3	300.3
Valencia	141.0	144.2	137.4	311.7	342.5	280.2	237.4	339.4	132.6	475.1	675.6	274.3
Health Region												
Northwest	125.1	174.7	74.1	279.3	403.6	162.8	240.6	368.3	115.5	469.7	621.7	322.6
Northeast	155.9	235.1	68.6	174.5	231.5	114.9	231.5	354.9	108.8	354.6	483.5	234.9
Bernalillo County	87.6	145.0	29.1	135.7	180.3	90.0	211.8	292.2	131.7	425.8	596.1	266.9
Southeast	87.6	130.3	43.4	185.2	278.1	90.2	215.4	249.8	181.0	465.9	582.3	350.6
Southwest	70.7	107.4	32.4	115.4	141.2	88.9	182.4	264.5	104.9	451.2	592.9	318.1

Age-specific death rates are the numbers of deaths per 100,000 population in specified sex and age groups.
 Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
 Unknown ages, sexes and counties, if any, are included in the "All Ages," "Both Sexes" and New Mexico categories.
 For counties included in state health regions, please see Technical Appendix.
 Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.
 U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-3 (continued)

55 to 64 Years			65 to 74 Years			75 to 84 Years			85 Plus Years			Resident County
Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	
890.9	1,110.0	687.0	2,062.1	2,516.2	1,677.9	5,115.0	6,177.7	4,388.3	13,253.1	14,309.1	12,759.0	United States
916.6	1,165.8	684.6	1,703.2	2,119.5	1,342.7	3,896.7	4,671.9	3,329.1	25,117.7	23,563.7	26,221.3	New Mexico
942.3	1,267.7	639.0	1,433.7	1,848.4	1,093.3	3,359.3	4,021.9	2,888.0	24,353.9	23,007.7	25,324.3	Bernalillo
919.8	1,017.8	815.2	1,964.6	2,836.9	881.1	4,924.2	5,357.1	4,605.3	15,555.6	38,461.5	6,250.0	Catron
935.6	1,001.1	878.0	1,666.0	2,469.1	1,038.6	4,606.0	5,391.8	4,025.9	28,750.0	25,954.2	30,687.8	Chaves
811.6	879.1	753.8	2,274.6	3,198.7	1,412.9	3,712.3	3,793.1	3,646.6	26,380.4	30,434.8	23,404.3	Cibola
845.4	1,065.9	612.9	1,715.0	1,891.9	1,546.4	3,753.9	4,798.0	3,019.5	24,456.5	24,637.7	24,347.8	Colfax
1,192.5	1,399.4	1,015.2	1,925.4	2,290.6	1,605.5	3,505.5	4,207.9	3,088.2	26,486.5	26,865.7	26,271.2	Curry
1,102.9	757.6	1,428.6	619.2	632.9	606.1	1,619.4	4,301.1	0.0	30,555.6	7,692.3	43,478.3	De Baca
818.2	951.6	695.0	1,654.5	2,001.6	1,365.0	4,072.6	4,954.6	3,378.4	24,063.0	23,185.8	24,707.4	Doña Ana
1,013.6	1,275.9	762.8	2,213.3	2,644.6	1,851.0	4,654.8	5,177.5	4,263.6	25,411.3	20,673.1	28,318.6	Eddy
921.3	957.2	886.3	2,002.7	2,614.8	1,437.0	4,526.9	5,876.3	3,470.5	30,226.0	29,166.7	30,952.4	Grant
1,877.9	2,463.1	1,345.3	915.3	465.1	1,351.4	5,607.5	10,227.3	2,381.0	72,222.2	37,500.0	100,000.0	Guadalupe
1,851.9	3,448.3	0.0	602.4	1,176.5	0.0	3,409.1	3,030.3	3,636.4	11,764.7	16,666.7	9,090.9	Harding
763.4	955.4	586.5	2,788.8	3,018.9	2,531.6	3,978.8	4,268.3	3,755.9	22,413.8	9,090.9	30,555.6	Hidalgo
1,133.9	1,372.9	901.6	2,061.3	2,809.3	1,438.8	4,382.1	5,136.7	3,850.6	22,894.2	19,230.8	25,266.9	Lea
814.2	1,105.0	552.5	1,632.2	2,123.0	1,150.3	2,766.0	2,971.8	2,578.0	14,427.9	18,823.5	11,206.9	Lincoln
259.2	248.4	270.8	798.9	277.0	1,282.1	2,916.7	3,081.7	2,781.3	19,607.8	17,525.8	21,495.3	Los Alamos
966.2	1,408.5	566.2	2,117.3	2,849.4	1,350.4	4,495.1	5,429.3	3,605.8	22,491.3	26,923.1	18,867.9	Luna
980.4	1,479.3	546.6	2,229.7	3,312.6	1,480.4	6,349.2	7,219.7	5,735.9	33,466.1	26,548.7	39,130.4	McKinley
1,639.3	1,685.4	1,587.3	2,076.1	2,464.8	1,700.7	2,839.1	3,246.8	2,454.0	31,707.3	28,571.4	35,000.0	Mora
977.6	1,189.7	784.1	1,933.8	1,820.3	2,043.0	4,585.9	4,934.2	4,337.6	24,757.3	18,390.8	29,411.8	Otero
911.0	852.3	968.2	1,529.8	2,295.1	791.1	2,461.9	2,695.4	2,282.2	20,161.3	28,571.4	15,853.7	Quay
1,154.2	1,608.6	668.6	1,956.1	2,730.6	1,282.9	4,274.0	4,731.5	3,931.0	28,514.1	29,896.9	27,631.6	Rio Arriba
746.7	995.0	498.1	2,000.0	1,973.7	2,023.1	4,096.6	4,567.3	3,731.3	31,250.0	26,250.0	35,416.7	Roosevelt
704.0	823.0	592.0	1,787.2	2,087.2	1,485.5	4,093.8	5,624.7	3,130.9	16,941.7	15,591.4	17,877.1	San Juan
895.5	1,050.7	750.6	2,727.3	3,542.9	2,081.4	3,582.9	5,089.8	2,458.1	31,277.5	29,347.8	32,592.6	San Miguel
1,080.5	1,377.3	802.4	1,631.1	1,736.5	1,541.7	4,254.0	5,203.4	3,537.0	28,333.3	29,251.7	27,699.5	Sandoval
661.6	872.4	477.5	1,456.2	1,822.9	1,141.4	3,441.7	3,897.6	3,101.6	31,188.7	28,311.7	33,270.7	Santa Fe
1,601.8	2,638.2	735.3	2,519.7	2,912.6	2,057.1	4,174.2	5,343.5	3,114.2	21,186.4	19,469.0	22,764.2	Sierra
967.2	1,161.6	765.9	1,596.6	1,903.1	1,307.2	6,119.4	7,191.8	5,291.0	31,858.4	27,083.3	35,384.6	Socorro
858.8	907.4	815.1	1,430.1	1,648.7	1,222.8	3,620.8	4,137.9	3,301.4	29,032.3	30,208.3	28,289.5	Taos
1,917.5	2,339.2	1,501.2	1,523.4	2,145.9	883.0	8,252.4	10,638.3	6,250.0	31,034.5	17,857.1	37,288.1	Torrance
189.4	373.1	0.0	1,279.3	1,421.8	1,162.8	3,448.3	2,272.7	4,430.4	14,516.1	17,391.3	12,820.5	Union
1,075.6	1,352.9	804.6	2,424.8	2,990.2	1,947.0	6,356.5	8,016.0	5,181.0	24,168.5	21,634.6	26,337.4	Valencia
Health Region												
917.7	1,162.6	690.1	1,949.9	2,378.9	1,573.0	4,697.6	5,856.4	3,883.2	23,817.2	22,727.3	24,617.5	Northwest
764.7	965.8	576.5	1,605.7	1,940.7	1,311.1	3,568.5	4,167.3	3,124.6	28,837.2	27,139.6	30,031.7	Northeast
942.3	1,267.7	639.0	1,433.7	1,848.4	1,093.3	3,359.3	4,021.9	2,888.0	24,353.9	23,007.7	25,324.3	Bernalillo County
1,033.2	1,213.3	866.2	1,895.0	2,454.1	1,423.8	4,178.7	4,868.1	3,688.9	26,127.3	23,408.8	27,870.0	Southeast
943.1	1,159.4	742.1	1,836.8	2,207.7	1,492.4	4,303.3	5,156.4	3,617.3	24,193.0	22,919.8	25,124.5	Southwest

Age-specific death rates are the numbers of deaths per 100,000 population in specified sex and age groups.

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Unknown ages, sexes and counties, if any, are included in the "All Ages," "Both Sexes" and New Mexico categories.

For counties included in state health regions, please see Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Mortality Section

**Table M-4
Numbers and Rates Ranked by Leading Causes of Death by County and Health Region
New Mexico, 2007, and United States, 2006
Both Sexes**

County			1		2		3		4		5	
	All Causes		Diseases of heart		Malignant neoplasms		Accidents (Unintentional injuries)		Chronic lower respiratory diseases		Cerebrovascular disease	
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate
United States	2,426,264	776.5	631,636	200.2	559,888	180.7	121,599	39.8	124,583	40.5	137,119	43.6
New Mexico	15,400	902.2	3,263	209.2	3,210	167.0	1,277	69.5	897	52.1	649	41.8
Bernalillo	4,829	842.9	1,057	196.8	976	153.9	396	66.6	257	44.3	244	47.2
Catron	43	829.5	7	114.9	13	242.1	3	70.3	1	34.5	0	0.0
Chaves	594	1,008.1	138	246.0	127	192.5	32	52.3	39	61.1	28	51.1
Cibola	228	994.3	27	130.8	50	194.7	24	88.9	13	60.9	5	21.3
Colfax	142	865.5	33	213.9	30	176.7	8	54.9	8	39.8	6	31.2
Curry	344	927.3	89	268.8	66	150.9	28	73.9	20	51.8	11	27.0
De Baca	25	952.5	7	239.8	5	192.5	2	107.4	2	50.2	1	43.1
Doña Ana	1,287	838.7	245	176.1	278	157.8	89	54.2	86	55.1	58	40.9
Eddy	498	974.1	119	234.3	112	192.6	32	65.2	29	50.7	15	29.3
Grant	351	1,043.0	77	229.7	72	183.7	22	71.0	24	74.8	18	50.2
Guadalupe	44	1,723.0	9	316.2	10	328.2	3	52.0	2	41.4	3	187.4
Harding	9	831.4	2	172.0	3	152.9	3	425.7	0	0.0	0	0.0
Hidalgo	55	928.4	14	232.4	10	169.8	2	51.8	4	63.7	4	53.0
Lea	471	941.2	103	220.8	95	162.3	39	67.6	35	75.2	14	29.2
Lincoln	181	719.0	32	131.0	53	171.3	21	100.3	10	34.3	8	32.9
Los Alamos	115	575.1	23	130.7	29	135.2	7	32.6	7	35.0	3	13.8
Luna	267	946.2	97	348.0	59	180.3	19	69.4	13	52.5	6	29.6
McKinley	495	1,263.4	76	240.5	70	192.5	67	104.6	22	71.6	13	41.2
Mora	47	943.4	16	426.3	8	125.3	3	40.8	2	50.8	5	90.5
Otero	481	922.3	136	308.7	100	156.5	21	38.9	30	54.0	21	39.7
Quay	98	797.1	29	221.5	16	116.7	5	47.1	9	90.9	5	40.8
Rio Arriba	373	1,122.0	65	239.0	66	178.8	48	113.9	21	69.6	8	26.6
Roosevelt	158	1,015.4	37	271.2	38	216.2	12	68.6	13	78.1	8	52.4
San Juan	867	820.6	145	148.1	176	154.3	112	95.4	43	37.9	31	33.4
San Miguel	266	1,072.1	55	245.6	45	158.2	20	65.0	17	66.2	4	19.5
Sandoval	809	949.1	159	213.1	173	170.6	54	59.7	48	55.5	42	60.9
Santa Fe	943	902.6	168	192.3	231	190.5	85	70.6	49	51.5	36	35.0
Sierra	219	1,013.5	59	284.6	51	233.0	11	59.6	31	126.3	10	44.1
Socorro	151	1,166.9	38	355.8	25	187.0	10	61.5	8	56.3	6	48.8
Taos	269	979.2	44	171.2	58	168.6	31	106.8	7	20.0	10	52.0
Torrance	127	1,218.3	29	340.0	36	301.8	11	106.8	8	72.9	5	51.8
Union	30	581.4	6	99.2	6	112.8	1	22.9	1	25.0	0	0.0
Valencia	578	1,088.8	121	271.5	123	206.3	54	84.9	38	73.3	20	41.1
Health Region												
Northwest	2,977	964.7	528	196.9	592	173.6	311	84.3	164	53.8	111	42.6
Northeast	2,229	923.3	419	200.0	483	176.1	206	74.3	114	48.1	75	33.7
Bernalillo County	4,829	842.9	1,057	196.8	976	153.9	396	66.6	257	44.3	244	47.2
Southeast	2,197	958.9	524	242.0	462	176.8	153	64.6	147	62.2	82	37.1
Southwest	3,162	898.0	734	227.3	697	172.9	209	58.5	215	60.2	136	40.4

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Deaths with unknown county of residence, if any, are included in the New Mexico totals. Numbers may not add to totals.

* Table is ranked by number of deaths for New Mexico.

For counties included in state health regions, please see Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-4 (continued)

6		7		8		9		10		County
Diabetes mellitus		Intentional self-harm (Suicide)		Chronic liver disease & cirrhosis		Influenza and pneumonia		Alzheimer's disease		
No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
72,449	23.3	33,300	10.9	27,555	8.8	56,326	17.8	72,432	22.6	United States
642	35.2	400	19.5	383	17.8	349	23.0	324	25.3	New Mexico
153	25.0	124	18.7	113	16.0	112	21.4	103	22.5	Bernalillo
3	39.9	4	92.7	0	0.0	1	34.5	3	81.9	Catron
22	36.8	10	15.1	12	17.3	14	27.9	28	54.5	Chaves
14	56.4	6	20.4	6	19.1	3	22.8	5	15.0	Cibola
3	14.3	5	36.8	9	43.8	6	43.1	1	4.7	Colfax
13	31.6	5	11.2	5	10.5	8	25.5	19	58.4	Curry
1	32.1	1	53.8	0	0.0	1	43.1	0	0.0	De Baca
74	46.9	26	13.8	31	16.0	24	16.2	27	23.2	Doña Ana
15	28.9	9	18.3	6	11.6	5	12.4	11	21.5	Eddy
13	41.1	5	14.9	3	8.4	9	34.9	10	39.2	Grant
2	36.1	0	0.0	1	21.0	1	86.2	1	86.2	Guadalupe
0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	Harding
3	39.6	3	48.6	2	33.0	1	26.7	1	26.7	Hidalgo
14	29.8	9	16.9	12	21.5	18	41.5	7	19.8	Lea
8	32.6	3	16.4	2	6.0	2	11.2	3	23.1	Lincoln
0	0.0	4	18.2	0	0.0	2	7.5	4	22.7	Los Alamos
10	30.2	4	13.1	2	7.1	4	16.3	4	15.2	Luna
40	118.3	17	24.2	39	67.3	9	32.6	5	20.9	McKinley
5	62.6	1	13.0	1	14.1	0	0.0	0	0.0	Mora
22	42.7	16	24.9	5	6.7	12	22.3	5	18.8	Otero
2	12.2	3	19.9	5	39.8	0	0.0	0	0.0	Quay
25	70.7	15	34.4	25	53.9	9	36.2	4	24.9	Rio Arriba
8	49.0	2	10.4	0	0.0	0	0.0	5	32.1	Roosevelt
44	41.1	25	18.0	20	14.9	21	22.2	11	16.2	San Juan
18	73.0	9	29.5	4	10.8	9	38.6	4	27.3	San Miguel
33	31.4	26	21.8	29	23.7	27	31.9	19	35.8	Sandoval
31	21.1	25	15.6	21	13.1	20	20.1	25	35.4	Santa Fe
7	39.9	2	8.5	3	17.4	5	23.6	2	13.1	Sierra
9	60.5	6	32.9	3	17.5	3	18.0	4	39.7	Socorro
22	92.6	13	43.5	9	24.4	8	41.8	7	40.5	Taos
5	52.5	5	28.3	0	0.0	1	10.9	1	17.8	Torrance
1	15.5	1	39.4	0	0.0	4	90.5	0	0.0	Union
22	34.2	16	23.0	15	20.0	10	19.7	5	14.0	Valencia
										Health Region
153	46.6	90	21.2	109	26.1	70	25.3	45	22.2	Northwest
107	39.3	73	24.2	70	20.4	59	29.2	46	28.6	Northeast
153	25.0	124	18.7	113	16.0	112	21.4	103	22.5	Bernalillo County
75	31.9	39	15.7	40	15.1	46	23.3	70	34.7	Southeast
154	42.4	74	18.1	51	12.0	62	19.8	60	24.1	Southwest

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Deaths with unknown county of residence, if any, are included in the New Mexico totals. Numbers may not add to totals.

* Table is ranked by number of deaths for New Mexico.

For counties included in state health regions, please see Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Mortality Section

**Table M-5
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
All Ages**

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	902.2	1014.6	798.2
Diseases of heart	209.2	232.8	187.9
Malignant Neoplasms	167.0	199.2	142.0
Accidents (unintentional injuries)	69.5	94.0	46.4
Chronic lower respiratory disease	52.1	58.4	47.3
Cerebrovascular diseases	41.8	37.6	44.6
American Indian or Alaska Native			
All Causes	917.8	1074.7	781.9
Accidents (unintentional injuries)	81.4	121.0	47.3
Diseases of heart	133.6	185.4	95.5
Malignant Neoplasms	114.6	127.3	105.2
Diabetes mellitus	81.0	83.0	78.2
Chronic liver disease and cirrhosis	51.1	66.2	37.9
Asian or Pacific Islander			
All Causes	662.3	659.3	658.0
Malignant Neoplasms	119.6	128.3	111.8
Diseases of heart	170.6	188.4	156.8
Accidents (unintentional injuries)	58.3	70.8	49.3
Diabetes mellitus	43.7	10.3	65.9
Cerebrovascular diseases	54.3	107.0	17.0
Black or African American			
All Causes	794.3	1000.5	601.8
Diseases of heart	179.3	160.1	184.2
Malignant Neoplasms	104.7	143.1	69.3
Diabetes mellitus	80.5	126.1	42.6
Accidents (unintentional injuries)	46.6	86.1	15.4
Cerebrovascular diseases	34.4	47.4	22.4
Essential hypertension and hypertensive renal disease	36.0	40.1	32.2
Hispanic			
All Causes	914.0	1035.5	800.0
Malignant Neoplasms	171.0	201.8	146.4
Diseases of heart	204.1	224.7	184.9
Accidents (unintentional injuries)	66.5	97.3	36.9
Diabetes mellitus	47.8	44.8	49.7
Chronic lower respiratory disease	41.7	48.0	36.9
White			
All Causes	883.5	972.5	801.3
Diseases of heart	218.2	241.1	197.2
Malignant Neoplasms	173.9	208.2	147.3
Chronic lower respiratory disease	60.5	66.4	55.8
Accidents (unintentional injuries)	64.9	80.4	50.3
Cerebrovascular diseases	41.4	36.5	44.4

**Table M-6
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
Ages 1 to 4 Years**

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	47.4	43.8	51.1
Accidents (unintentional injuries)	17.1	11.9	22.5
Congenital malformations	6.0	6.0	6.1
Assault (homicide)	4.0	6.0	2.0
Influenza and pneumonia	3.0	4.0	2.0
Malignant Neoplasms	3.0	2.0	4.1
American Indian or Alaska Native			
All Causes	87.8	88.3	87.4
Accidents (unintentional injuries)	43.9	14.7	72.8
Asian or Pacific Islander			
All Causes	0.0	0.0	0.0
Black or African American			
All Causes	76.0	0.0	151.1
Hispanic			
All Causes	45.0	38.5	51.8
Accidents (unintentional injuries)	13.7	11.6	15.9
Congenital malformations	5.9	3.9	8.0
White			
All Causes	33.1	39.2	26.9
Accidents (unintentional injuries)	9.9	13.1	6.7

Age-adjusted rates (table M-5) are the numbers of deaths per 1000,000 U.S. standard population. Age-specific rates are the numbers of deaths per 100,000 population in specified age group.

Note: Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are three or more deaths. The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, ages or causes of death, if any, are included in the "All Races," "All Ages" and "All Causes" categories.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	15.0	16.9	13.1
Accidents (Unintentional injuries)	7.1	8.4	5.8
Intentional self harm (Suicide)	2.1	4.2	0.0
Malignant neoplasms (Cancer)	1.4	0.7	2.2
Assault (Homicide)	1.1	1.4	0.7
American Indian or Alaska Native			
All Causes	23.9	33.6	14.3
Accidents (Unintentional injuries)	9.6	9.6	9.5
Intentional self harm (Suicide)	7.2	14.4	0.0
Asian or Pacific Islander			
All Causes	0.0	0.0	0.0
Black or African American			
All Causes	10.2	20.3	0.0
Hispanic			
All Causes	15.3	15.8	14.8
Accidents (Unintentional injuries)	6.6	8.6	4.5
Assault (Homicide)	2.2	2.9	1.5
White			
All Causes	11.6	11.3	11.9
Accidents (Unintentional injuries)	8.1	9.0	7.1

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	101.5	153.5	47.4
Accidents (Unintentional injuries)	47.1	72.8	20.5
Intentional self harm (Suicide)	18.9	30.8	6.4
Assault (Homicide)	14.1	23.4	4.5
Malignant neoplasms (Cancer)	3.8	5.5	1.9
Congenital malformations	1.9	1.8	1.9
Diseases of heart	1.9	3.1	0.6
American Indian or Alaska Native			
All Causes	151.2	223.8	81.6
Accidents (Unintentional injuries)	63.6	94.0	34.4
Intentional self harm (Suicide)	30.7	58.2	4.3
Assault (Homicide)	19.7	31.3	8.6
Asian or Pacific Islander			
All Causes	111.5	138.0	80.5
Black or African American			
All Causes	123.7	201.5	39.6
Accidents (Unintentional injuries)	47.6	73.3	19.8
Assault (Homicide)	47.6	91.6	0.0
Hispanic			
All Causes	102.2	164.1	38.6
Accidents (Unintentional injuries)	49.0	82.1	15.2
Assault (Homicide)	17.0	30.9	2.8
Intentional self harm (Suicide)	15.7	26.9	4.1
Malignant neoplasms (Cancer)	4.1	5.4	2.8
Congenital malformations	2.0	1.3	2.8
White			
All Causes	76.4	106.8	43.6
Accidents (Unintentional injuries)	38.2	54.3	20.8
Intentional self harm (Suicide)	19.1	28.0	9.5
Assault (Homicide)	4.5	5.3	3.8
Diseases of heart	2.7	3.5	1.9
Malignant neoplasms (Cancer)	2.7	5.3	0.0

Age-specific rates are the numbers of deaths per 100,000 population in specified age group.

Note: Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are three or more deaths. The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, ages or causes of death, if any, are included in the "All Races," "All Ages" and "All Causes" categories.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

**Table M-9
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
Ages 25 to 44 Years**

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	197.1	276.3	118.2
Accidents (Unintentional injuries)	58.5	87.8	29.3
Intentional self harm (Suicide)	25.2	40.5	10.0
Malignant neoplasms (Cancer)	16.5	17.1	15.9
Chronic liver disease and cirrhosis	14.8	22.3	7.4
Assault (Homicide)	12.2	20.1	4.4
Diseases of heart	12.2	18.6	5.9
American Indian or Alaska Native			
All Causes	335.4	519.7	161.9
Accidents (Unintentional injuries)	104.3	162.2	49.8
Chronic liver disease and cirrhosis	46.5	62.9	31.1
Intentional self harm (Suicide)	44.9	86.1	6.2
Assault (Homicide)	19.3	36.4	3.1
Diseases of heart	14.4	23.2	6.2
Asian or Pacific Islander			
All Causes	34.0	35.8	32.3
Black or African American			
All Causes	124.2	186.7	53.1
Accidents (Unintentional injuries)	37.2	58.3	13.3
Assault (Homicide)	18.6	23.3	13.3
Hispanic			
All Causes	197.7	284.4	109.7
Accidents (Unintentional injuries)	61.1	98.1	23.5
Intentional self harm (Suicide)	23.7	39.6	7.5
Malignant neoplasms (Cancer)	15.8	14.8	16.8
Chronic liver disease and cirrhosis	15.0	25.6	4.2
Assault (Homicide)	14.1	23.9	4.2
White			
All Causes	167.9	212.2	124.2
Accidents (Unintentional injuries)	46.2	60.1	32.5
Intentional self harm (Suicide)	23.6	33.0	14.3
Malignant neoplasms (Cancer)	23.1	25.2	21.0
Diseases of heart	12.0	15.5	8.6
Assault (Homicide)	8.2	11.6	4.8

**Table M-10
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
Ages 45 to 64 Years**

Race/Ethnicity	Rate		
	Total	Male	Female
All Races			
All Causes	626.5	814.2	450.0
Malignant neoplasms (Cancer)	164.1	180.3	148.9
Diseases of heart	113.3	164.6	65.0
Accidents (Unintentional injuries)	69.3	103.7	36.9
Chronic liver disease and cirrhosis	37.9	57.7	19.2
Diabetes Mellitus	29.9	35.0	25.1
American Indian or Alaska Native			
All Causes	612.4	846.0	411.1
Chronic liver disease and cirrhosis	97.9	137.7	63.6
Malignant neoplasms (Cancer)	93.3	98.4	89.0
Accidents (Unintentional injuries)	79.7	118.0	46.6
Diseases of heart	77.4	127.9	33.9
Diabetes Mellitus	54.6	78.7	33.9
Asian or Pacific Islander			
All Causes	320.8	333.5	311.5
Malignant neoplasms (Cancer)	166.8	151.6	178.0
Diseases of heart	38.5	60.6	22.2
Black or African American			
All Causes	620.2	882.0	321.7
Diseases of heart	131.6	158.8	100.5
Diabetes Mellitus	122.2	194.0	40.2
Malignant neoplasms (Cancer)	94.0	141.1	40.2
Accidents (Unintentional injuries)	47.0	70.6	20.1
Hispanic			
All Causes	641.0	887.7	411.3
Malignant neoplasms (Cancer)	164.0	188.9	140.4
Diseases of heart	103.9	162.4	48.3
Accidents (Unintentional injuries)	79.8	124.4	37.3
Chronic liver disease and cirrhosis	49.4	80.6	19.7
Diabetes Mellitus	29.2	34.6	24.1
White			
All Causes	619.8	761.9	485.5
Malignant neoplasms (Cancer)	176.5	187.8	165.9
Diseases of heart	124.9	173.3	79.2
Accidents (Unintentional injuries)	61.1	86.7	36.9
Intentional self harm (Suicide)	30.9	47.7	15.0
Chronic lower respiratory disease	28.1	28.9	27.3

Age-specific rates are the numbers of deaths per 100,000 population in specified age group.
 Note: Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are three or more deaths. The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, ages or causes of death, if any, are included in the "All Races," "All Ages" and "All Causes" categories.
 Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.
<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Table M-11
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
Ages 65 to 84

Race/Ethnicity	Rate		
	Total	Male	Female
New Mexico - All Races			
All Causes	2,598.3	3,103.9	2,189.2
Malignant neoplasms	703.1	867.4	570.1
Diseases of heart	554.3	701.1	435.5
Chronic lower respiratory disease	221.6	261.1	189.5
Diabetes mellitus	134.8	157.4	116.5
Cerebrovascular diseases	126.0	132.3	120.8
American Indian or Alaska Native			
All Causes	2,457.7	2,782.1	2,219.1
Malignant neoplasms	432.9	471.5	404.5
Diseases of heart	426.3	565.9	323.6
Diabetes mellitus	319.7	298.6	335.2
Cerebrovascular diseases	119.9	157.2	92.5
Accidents (unintentional injuries)	106.6	141.5	80.9
Chronic liver disease and cirrhosis	106.6	110.0	104.0
Asian or Pacific Islander			
All Causes	1,462.8	1,039.5	1,734.5
Malignant neoplasms	528.2	311.9	667.1
Diseases of heart	243.8	104.0	333.6
Diabetes mellitus	162.5	104.0	200.1
Black or African American			
All Causes	2,514.3	3,822.7	1,446.7
Malignant neoplasms	572.6	941.8	271.2
Diseases of heart	497.9	720.2	316.5
Diabetes mellitus	224.0	277.0	180.8
Essential (Primary) hypertension	224.0	277.0	180.8
Cerebrovascular diseases	199.2	332.4	90.4
Nephritis, nephrotic syndrome and nephrosis	149.4	221.6	90.4
Hispanic			
All Causes	2,611.1	3,192.4	2,146.4
Malignant neoplasms	658.4	798.8	546.1
Diseases of heart	567.3	718.7	446.4
Diabetes mellitus	201.9	234.6	175.7
Chronic lower respiratory disease	174.2	201.9	152.0
Cerebrovascular diseases	112.1	118.8	106.8
White			
All Causes	2,619.7	3,085.9	2,235.4
Malignant neoplasms	757.4	941.7	605.4
Diseases of heart	566.2	711.3	446.6
Chronic lower respiratory disease	265.1	304.2	232.9
Cerebrovascular diseases	131.6	131.8	131.4
Accidents (unintentional injuries)	96.2	108.7	86.0

Table M-12
Death Rates by Race/Ethnicity and Sex for
Ranked Causes of Death
New Mexico, 2007
Ages 85 Plus Years

Race/Ethnicity	Rate		
	Total	Male	Female
New Mexico - All Races			
All Causes	25,117.7	23,563.7	26,221.3
Diseases of heart	7,679.1	7,170.2	8,040.5
Malignant neoplasms	3,053.8	3,615.7	2,654.8
Cerebrovascular diseases	1,546.0	1,195.0	1,795.2
Chronic lower respiratory disease	1,469.7	1,516.8	1,436.2
Alzheimer's disease	1,291.5	934.6	1,545.0
American Indian or Alaska Native			
All Causes	23,015.9	22,653.7	23,266.2
Diseases of heart	3,968.3	5,178.0	3,132.0
Malignant neoplasms	2,645.5	2,912.6	2,460.9
Diabetes mellitus	1,851.9	1,618.1	2,013.4
Influenza and pneumonia	1,719.6	1,294.5	2,013.4
Cerebrovascular diseases	1,322.8	970.9	1,566.0
Asian or Pacific Islander			
All Causes	25,352.1	27,586.2	23,809.5
Diseases of heart	8,450.7	10,344.8	7,142.9
Black or African American			
All Causes	17,910.4	13,157.9	20,800.0
Diseases of heart	5,472.6	1,315.8	8,000.0
Diabetes mellitus	1,492.5	2,631.6	800.0
Hispanic			
All Causes	24,987.9	22,586.1	26,800.2
Diseases of heart	7,236.5	6,324.1	7,925.0
Malignant neoplasms	3,496.8	4,065.5	3,067.7
Cerebrovascular diseases	1,627.0	1,242.2	1,917.3
Chronic lower respiratory disease	1,262.7	1,411.6	1,150.4
Alzheimer's disease	1,189.9	734.0	1,533.9
White			
All Causes	25,435.1	24,136.3	26,340.3
Diseases of heart	8,144.2	7,715.3	8,443.0
Malignant neoplasms	2,960.7	3,546.8	2,552.2
Chronic lower respiratory disease	1,598.6	1,612.2	1,589.1
Cerebrovascular diseases	1,541.8	1,174.6	1,797.8
Alzheimer's disease	1,381.0	1,059.4	1,605.1

Age-specific rates are the numbers of deaths per 100,000 population in specified age group.

Note: Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are three or more deaths. The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, ages or causes of death, if any, are included in the "All Races," "All Ages" and "All Causes" categories.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Table M-13
Numbers and Rates of Deaths by Race/Ethnicity and Sex
New Mexico, 2003-2007, and United States, 2003-2006

Race/Ethnicity and Sex	2003		2004		2005		2006		2007	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
New Mexico										
All Races										
Both Sexes	14,493	796.6	14,197	764.7	14,866	770.3	15,231	761.2	15,400	902.2
Male	7,590	945.5	7,486	924.8	7,824	919.8	7,960	900.0	8,233	1,014.6
Female	6,902	665.7	6,710	631.2	7,042	643.6	7,271	639.8	7,167	798.2
American Indian or Alaska Native										
Both Sexes	1,024	807.8	1,020	809.3	1,063	739.2	1,010	689.6	1,142	917.8
Male	594	1,023.2	591	1,095.3	619	966.9	563	825.9	657	1,074.7
Female	430	632.8	429	614.5	444	566.1	447	573.5	485	781.9
Asian or Pacific Islander										
Both Sexes	51	331.8	47	275.3	75	445.5	70	365.4	90	662.3
Male	28	448.2	19	302.9	38	591.7	29	379.7	36	659.3
Female	23	259.8	28	266.7	37	363.6	41	357.7	54	658.0
Black or African American										
Both Sexes	237	843.7	236	770.4	246	768.0	249	712.3	248	794.3
Male	142	1,082.0	130	919.8	121	836.2	140	846.7	163	1,000.5
Female	95	642.5	106	638.4	125	707.0	109	586.7	85	601.8
Hispanic										
Both Sexes	4,533	801.1	4,465	751.5	4,717	763.6	4,865	759.7	4,924	914.0
Male	2,520	970.2	2,589	951.7	2,620	917.6	2,647	884.7	2,788	1,035.5
Female	2,013	652.1	1,876	580.1	2,097	628.5	2,218	644.2	2,136	800.0
White										
Both Sexes	8,625	783.5	8,417	756.9	8,742	766.5	8,983	761.1	8,954	883.5
Male	4,291	904.7	4,151	880.3	4,412	905.5	4,546	901.2	4,553	972.5
Female	4,334	677.1	4,266	654.9	4,330	649.7	4,437	642.8	4,401	801.3
United States										
All Races										
All Deaths	2,448,288	832.7	2,397,615	800.8	2,448,017	798.8	2,426,264	776.5	N/A	N/A
Male	1,201,964	994.3	1,181,668	955.7	1,207,675	951.1	1,201,942	924.8	N/A	N/A
Female	1,246,324	706.2	1,215,947	679.2	1,240,342	677.6	1,224,322	657.8	N/A	N/A

The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown sexes and races, if any, are included in the "Both Sexes" and "All Races" categories.

U.S. Data Source: CDC, National Center for Health Statistics; NVSR, Vol. 57, No. 14, released April 2009, <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Table M-14
Rates for Leading Causes of Death
New Mexico, 1999-2007

Cause of Death		1999	2000	2001	2002	2003	2004	2005	2006	2007
All Causes		827.6	802.5	814.5	797.5	796.6	764.7	770.3	761.2	902.2
1	Diseases of heart	212.5	195.3	197.5	179.9	174.7	172.7	176.8	167.8	209.2
2	Malignant neoplasms	170.8	173.1	166.2	169.9	165.2	159.3	159.2	153.7	167.0
3	Accidents (Unintentional injuries)	51.4	53.6	53.7	57.0	60.8	60.3	61.3	64.0	69.5
4	Chronic lower respiratory diseases	53.0	47.0	47.4	50.0	53.5	41.6	45.5	46.2	52.1
5	Cerebrovascular diseases	53.2	49.0	49.0	39.7	40.5	33.9	33.1	32.2	41.8
6	Diabetes mellitus	31.8	29.8	32.1	32.8	34.9	33.9	32.0	27.5	35.2
7	Intentional self-harm (Suicide)	17.9	18.1	20.4	18.9	18.4	18.7	17.3	17.1	19.5
8	Chronic liver disease and cirrhosis	16.9	19.6	17.8	16.0	16.5	14.2	14.2	15.0	17.8

* Leading causes of death are ranked for the year 2007. Rank is based on the number of deaths per cause.

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Age-adjusting allows for comparison across time periods.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators. 1999 Population Source: U.S. Census Bureau in collaboration with the National Center for Health Statistics (NCHS).

Mortality Section

Table M-15
Number and Rates of Deaths for 39 Selected Causes by Age
New Mexico, 2007

Cause of Death - (ICD-10 Code)	All Ages		0-4 Years		5-24 Years		25-64 Years		65+ Years	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Tuberculosis (A16-A19)	3	0.1	0	0.0	0	0.0	0	0.0	3	1.1
Syphilis (A50-A53)	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.6	0	0.0	0	0.0	30	2.8	3	1.1
Malignant neoplasms (C00-C97)	3,210	167.0	3	2.4	16	2.7	951	89.4	2,238	842.1
Malignant neoplasms of stomach (C16)	99	5.3	0	0.0	0	0.0	32	3.0	67	25.2
Malignant neoplasms of colon, rectum and anus (C18-C21)	313	17.2	0	0.0	1	0.2	101	9.5	210	79.0
Malignant neoplasms of pancreas (C25)	193	9.9	0	0.0	0	0.0	68	6.4	125	47.0
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	772	38.8	0	0.0	1	0.2	180	16.9	590	222.0
Malignant neoplasms of breast (C50)	232	12.2	0	0.0	0	0.0	98	9.2	134	50.4
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	124	6.2	0	0.0	0	0.0	53	5.0	71	26.7
Malignant neoplasms of prostate (C61)	207	12.5	0	0.0	0	0.0	17	1.6	190	71.5
Malignant neoplasms of urinary tract (C64-C68)	187	10.0	0	0.0	0	0.0	48	4.5	139	52.3
Non-Hodgkin's lymphoma (C82-C85)	117	6.1	0	0.0	2	0.3	24	2.3	91	34.2
Leukemia (C91-C95)	129	6.9	1	0.8	7	1.2	27	2.5	94	35.4
Other malignant neoplasms (Residual)	837	41.9	2	1.6	5	0.8	303	28.5	527	198.3
Diabetes mellitus (E10-E14)	642	35.2	0	0.0	0	0.0	177	16.6	465	175.0
Alzheimer's disease (G30)	324	25.4	0	0.0	0	0.0	5	0.5	319	120.0
Major cardiovascular diseases (I00-I78)	4,334	278.7	5	4.0	8	1.3	807	75.8	3,514	1,322.2
Diseases of heart (I00-I09, I11, I13, I20-I51)	3,263	209.2	3	2.4	6	1.0	661	62.1	2,593	975.7
Hypertensive heart disease with or without renal diseases (I11, I13)	214	13.1	0	0.0	0	0.0	64	6.0	150	56.4
Ischemic heart diseases (I20-I25)	2,092	131.8	0	0.0	1	0.2	445	41.8	1,646	619.3
Other diseases of heart (I00-I09, I26-I51)	957	64.3	3	2.4	5	0.8	152	14.3	797	299.9
Essential (primary) hypertension & hypertensive renal disease (I10, I12)	200	13.0	0	0.0	0	0.0	31	2.9	169	63.6
Cerebrovascular diseases (I60-I69)	649	41.8	2	1.6	2	0.3	87	8.2	558	210.0
Atherosclerosis (I70)	95	6.7	0	0.0	0	0.0	7	0.7	88	33.1
Other diseases of circulatory system (I71-I78)	127	7.9	0	0.0	0	0.0	21	2.0	106	39.9
Influenza and pneumonia (J10-J18)	349	23.0	4	3.2	1	0.2	71	6.7	273	102.7
Chronic lower respiratory diseases (J40-J47)	897	52.1	1	0.8	1	0.2	110	10.3	785	295.4
Peptic ulcer (K25-K28)	20	1.0	0	0.0	0	0.0	8	0.8	12	4.5
Chronic liver disease and cirrhosis (K70, K73-K74)	383	17.8	0	0.0	0	0.0	279	26.2	104	39.1
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	249	14.6	1	0.8	1	0.2	47	4.4	200	75.3
Pregnancy, childbirth, and the puerperium (O00-O99)	4	0.2	0	0.0	0	0.0	4	0.4	0	0.0
Certain conditions originating in the perinatal period (P00-P96)	83	4.3	83	66.1	0	0.0	0	0.0	0	0.0
Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99)	76	3.9	46	36.6	8	1.3	13	1.2	9	3.4
Sudden infant death syndrome (R95)	15	0.8	15	11.9	0	0.0	0	0.0	0	0.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	68	4.1	2	1.6	6	1.0	28	2.6	32	12.0
All other diseases (Residual)	1,748	109.5	15	11.9	15	2.5	477	44.8	1,241	466.9
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	366	17.7	6	4.8	108	18.0	199	18.7	53	19.9
All other and unspecified accidents and adverse effects	910	51.7	15	11.9	62	10.4	479	45.0	354	133.2
Intentional self-harm (suicide) (X60-X84, Y87.0)	400	19.5	0	0.0	66	11.0	278	26.1	56	21.1
Assault (homicide) (X85-Y09, Y87.1)	164	8.0	10	8.0	48	8.0	97	9.1	9	3.4
All other external causes (Y10-Y36, Y87.2, Y89)	103	5.1	15	11.9	17	2.8	66	6.2	5	1.9

Other and unknown ages, if any, are included in the "All Ages" category. Numbers of deaths may not sum to totals, due to unknown ages. For the all ages category, age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. The other rates are age-specific rates. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Table M-16
Number and Rates of Deaths for 39 Selected Causes by Sex
New Mexico, 2007

Cause of Death - (ICD-10 Code)	Both Sexes		Female		Male	
	Number	Rate	Number	Rate	Number	Rate
Tuberculosis (A16-A19)	3	0.1	2	0.2	1	0.1
Syphilis (A50-A53)	0	0.0	0	0.0	0	0.0
Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.6	3	0.2	30	3.0
Malignant neoplasms (C00-C97)	3,210	167.0	1,487	142.0	1,723	199.2
Malignant neoplasms of stomach (C16)	99	5.3	35	3.5	64	7.5
Malignant neoplasms of colon, rectum and anus (C18-C21)	313	17.2	139	14.1	174	21.0
Malignant neoplasms of pancreas (C25)	193	9.9	102	9.9	91	9.9
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	772	38.8	347	32.7	425	46.2
Malignant neoplasms of breast (C50)	232	12.2	230	22.1	2	0.2
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	124	6.2	124	11.4	0	0.0
Malignant neoplasms of prostate (C61)	207	12.5	0	0.0	207	29.3
Malignant neoplasms of urinary tract (C64-C68)	187	10.0	66	6.6	121	14.4
Non-Hodgkin's lymphoma (C82-C85)	117	6.1	54	5.0	63	7.3
Leukemia (C91-C95)	129	6.9	54	5.7	75	8.4
Other malignant neoplasms (Residual)	837	41.9	336	31.0	501	55.0
Diabetes mellitus (E10-E14)	642	35.2	326	34.0	316	36.2
Alzheimer's disease (G30)	324	25.3	216	29.7	108	19.3
Major cardiovascular diseases (I00-I78)	4,334	278.7	2,124	258.2	2,210	300.2
Diseases of heart (I00-I09, I11, I13, I20-I51)	3,263	209.2	1,535	187.9	1,728	232.8
Hypertensive heart disease with or without renal diseases (I11, I13)	214	13.1	114	13.7	100	11.9
Ischemic heart diseases (I20-I25)	2,092	131.8	890	108.4	1,202	159.4
Other diseases of heart (I00-I09, I26-I51)	957	64.3	531	65.8	426	61.5
Essential (primary) hypertension & hypertensive renal disease (I10, I12)	200	13.1	110	13.4	90	12.5
Cerebrovascular diseases (I60-I69)	649	41.8	377	44.6	272	37.6
Atherosclerosis (I70)	95	6.7	39	4.8	56	9.2
Other diseases of circulatory system (I71-I78)	127	7.9	63	7.5	64	8.1
Influenza and pneumonia (J10-J18)	349	23.0	176	21.7	173	24.2
Chronic lower respiratory diseases (J40-J47)	897	52.1	447	47.3	450	58.4
Peptic ulcer (K25-K28)	20	1.0	6	0.6	14	1.4
Chronic liver disease and cirrhosis (K70, K73-K74)	383	17.8	119	10.6	264	25.4
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	249	14.6	114	11.7	135	18.3
Pregnancy, childbirth, and the puerperium (O00-O99)	4	0.2	4	0.5	0	0.0
Certain conditions originating in the perinatal period (P00-P96)	83	4.3	31	3.3	52	5.3
Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99)	76	3.9	36	3.7	40	4.0
Sudden infant death syndrome (R95)	15	0.8	4	0.4	11	1.1
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	68	4.1	36	4.2	32	3.8
All other diseases (Residual)	1,748	109.5	959	114.7	789	99.8
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	366	17.7	107	10.3	259	25.4
All other and unspecified accidents and adverse effects	910	51.7	322	36.1	588	68.5
Intentional self-harm (suicide) (X60-X84, Y87.0)	400	19.5	74	7.0	326	32.9
Assault (homicide) (X85-Y09, Y87.1)	164	8.0	38	3.7	126	12.5
All other external causes (Y10-Y36, Y87.2, Y89)	103	5.1	41	3.9	62	6.3

Other and unknown sexes, if any, are included in the "Both Sexes" categories. Numbers of deaths may not sum to totals, due to unknowns. Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Mortality Section

Table M-17
Number and Rate of Deaths for 39 Selected Causes by Health Region
New Mexico, 2007

Cause of Death - (ICD-10 Code)	All Regions		Northwest		Northeast		Bernalillo Co.		Southeast		Southwest	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Tuberculosis (A16-A19)	3	0.1	3	0.9	0	0.0	0	0.0	0	0.0	0	0.0
Syphilis (A50-A53)	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.6	10	2.4	3	0.8	12	1.7	2	0.8	6	1.7
Malignant neoplasms (C00-C97)	3,210	167.0	592	173.6	483	176.1	976	153.9	462	176.8	697	172.9
Malignant neoplasms of stomach (C16)	99	5.3	14	3.9	15	7.4	36	5.6	13	4.5	21	5.2
Malignant neoplasms of colon, rectum and anus (C18-C21)	313	17.2	64	18.9	63	25.2	75	12.8	46	19.0	65	16.6
Malignant neoplasms of pancreas (C25)	193	9.9	40	12.2	31	11.3	48	7.5	24	8.6	50	12.2
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	772	38.8	149	43.6	106	36.4	233	34.8	119	44.3	165	40.0
Malignant neoplasms of breast (C50)	232	12.2	32	9.0	36	11.0	72	12.0	42	17.5	50	12.6
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	124	6.2	28	8.5	24	8.0	35	5.0	16	6.2	21	5.2
Malignant neoplasms of prostate (C61)	207	12.5	36	12.4	31	15.0	67	12.0	25	10.0	48	13.5
Malignant neoplasms of urinary tract (C64-C68)	187	10.0	39	11.0	26	9.1	63	10.4	22	8.3	37	9.7
Non-Hodgkin's lymphoma (C82-C85)	117	6.1	19	4.4	18	6.6	37	6.0	14	5.4	29	6.7
Leukemia (C91-C95)	129	6.9	25	5.6	15	5.0	50	8.4	16	6.4	23	5.7
Other malignant neoplasms (Residual)	837	41.9	146	29.8	118	40.9	260	39.4	125	46.6	188	45.6
Diabetes mellitus (E10-E14)	642	35.2	153	46.6	107	39.3	153	25.0	75	31.9	154	42.4
Alzheimer's disease (G30)	324	25.3	45	22.2	46	28.6	103	22.5	70	34.7	60	24.1
Major cardiovascular diseases (I00-I78)	4,334	278.7	705	264.1	549	262.7	1,419	266.3	675	313.1	984	301.7
Diseases of heart (I00-I09, I11, I13, I20-I51)	3,263	209.2	528	196.9	419	200.0	1,057	196.8	524	242.0	734	227.3
Hypertensive heart disease with or without renal diseases (I11, I13)	214	13.1	37	13.6	34	17.6	76	13.2	27	11.8	40	11.1
Ischemic heart diseases (I20-I25)	2,092	131.8	313	113.1	263	118.7	685	125.8	356	163.5	474	145.1
Other diseases of heart (I00-I09, I26-I51)	957	64.3	178	70.2	122	63.7	296	57.7	141	66.7	220	71.1
Essential (primary) hypertension & hypertensive renal disease (I10, I12)	200	13.1	32	12.1	32	18.4	53	10.3	31	14.4	52	14.2
Cerebrovascular diseases (I60-I69)	649	41.8	111	42.6	75	33.7	244	47.2	82	37.1	136	40.4
Atherosclerosis (I70)	95	6.7	11	4.6	10	4.7	19	3.7	14	7.9	41	14.1
Other diseases of circulatory system (I71-I78)	127	7.9	23	7.8	13	6.0	46	8.4	24	11.6	21	5.7
Influenza and pneumonia (J10-J18)	349	23.0	70	25.3	59	29.2	112	21.4	46	23.3	62	19.8
Chronic lower respiratory diseases (J40-J47)	897	52.1	164	53.8	114	48.1	257	44.3	147	62.2	215	60.2
Peptic ulcer (K25-K28)	20	1.0	4	0.9	3	0.8	8	1.2	2	0.7	3	1.2
Chronic liver disease and cirrhosis (K70, K73-K74)	383	17.8	109	26.1	70	20.4	113	16.0	40	15.1	51	12.0
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	249	14.6	43	13.4	39	15.5	80	14.5	44	18.6	43	11.7
Pregnancy, childbirth, and the puerperium (O00-O99)	4	0.2	2	0.5	0	0.0	2	0.3	0	0.0	0	0.0
Certain conditions originating in the perinatal period (P00-P96)	83	4.3	21	5.2	6	2.5	33	5.5	11	4.4	12	2.9
Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99)	76	3.9	21	5.4	10	3.5	24	3.8	11	4.6	9	2.1
Sudden infant death syndrome (R95)	15	0.8	4	1.0	2	0.8	5	0.8	3	1.2	1	0.2
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	68	4.1	15	4.8	5	1.9	19	3.4	13	6.9	16	4.0
All other diseases (Residual)	1,748	109.5	325	117.0	268	118.9	593	108.7	250	117.5	312	94.2
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	366	17.7	109	25.7	64	21.3	88	13.8	47	17.6	58	13.6
All other and unspecified accidents and adverse effects	910	51.7	202	58.6	142	53.0	307	52.7	106	47.0	151	45.0
Intentional self-harm (suicide) (X60-X84, Y87.0)	400	19.5	90	21.2	73	24.2	124	18.7	39	15.7	74	18.1
Assault (homicide) (X85-Y09, Y87.1)	164	8.0	40	9.4	20	7.1	56	8.6	28	11.1	19	4.6
All other external causes (Y10-Y36, Y87.2, Y89)	103	5.1	29	6.9	17	6.0	25	4.0	7	2.7	25	6.0

Other and unknown regions, if any, are included in the "All Regions" category. Numbers of deaths may not sum to totals, due to unknowns. Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Table M-18
Number and Rate of Deaths for 39 Selected Causes by Race/Ethnicity
New Mexico, 2007

Cause of Death - (ICD-10 Code)	All Races		American Indian or Alaska Native		Asian or Pacific Islander		Black		Hispanic		White	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Tuberculosis (A16-A19)	3	0.1	2	1.7	0	0.0	0	0.0	0	0.0	1	0.1
Syphilis (A50-A53)	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.6	7	3.4	0	0.0	2	5.0	15	1.9	9	0.8
Malignant neoplasms (C00-C97)	3,210	167.0	131	114.6	28	119.6	36	104.7	983	171.0	2,028	173.9
Malignant neoplasms of stomach (C16)	99	5.3	8	5.1	2	6.2	0	0.0	58	11.2	31	2.8
Malignant neoplasms of colon, rectum and anus (C18-C21)	313	17.2	14	12.0	2	7.2	6	15.7	123	22.7	168	15.4
Malignant neoplasms of pancreas (C25)	193	9.9	12	11.8	1	4.1	0	0.0	61	11.2	119	9.8
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	772	38.8	14	13.1	6	24.8	13	40.0	177	30.2	559	46.0
Malignant neoplasms of breast (C50)	232	12.2	5	3.4	2	5.4	1	7.7	71	12.0	153	13.4
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	124	6.2	8	7.4	3	11.3	2	4.6	40	6.6	71	5.9
Malignant neoplasms of prostate (C61)	207	12.5	12	13.0	0	0.0	2	5.6	64	13.0	129	12.5
Malignant neoplasms of urinary tract (C64-C68)	187	10.0	14	12.2	1	5.3	2	6.0	61	10.9	109	9.5
Non-Hodgkin's lymphoma (C82-C85)	117	6.1	2	2.4	1	5.3	0	0.0	38	5.8	75	6.7
Leukemia (C91-C95)	129	6.9	4	1.8	1	4.1	2	5.0	28	3.6	94	7.7
Other malignant neoplasms (Residual)	837	41.9	38	32.2	9	45.9	8	20.0	262	42.9	520	42.9
Diabetes mellitus (E10-E14)	642	35.2	92	43.7	6	81.0	26	80.6	259	47.8	259	23.9
Alzheimer's disease (G30)	324	25.3	9	13.3	2	27.1	3	18.4	76	22.5	234	27.6
Major cardiovascular diseases (I00-I78)	4,334	278.7	197	200.6	19	224.8	75	275.2	1,269	276.8	2,765	285.4
Diseases of heart (I00-I09, I11, I13, I20-I51)	3,263	209.2	138	133.6	15	170.6	47	179.3	947	204.1	2,110	218.2
Hypertensive heart disease with or without renal diseases (I11, I13)	214	13.1	8	7.0	0	0.0	6	20.6	79	15.9	120	12.3
Ischemic heart diseases (I20-I25)	2,092	131.8	88	86.8	11	135.0	28	113.8	610	129.4	1,352	137.0
Other diseases of heart (I00-I09, I26-I51)	957	64.3	42	39.8	4	35.5	13	44.9	258	58.8	638	68.9
Essential (primary) hypertension & hypertensive renal disease (I10, I12)	200	13.1	15	18.3	0	0.0	11	36.0	67	15.3	107	11.1
Cerebrovascular diseases (I60-I69)	649	41.8	33	36.1	4	54.3	11	34.4	194	43.2	404	41.4
Atherosclerosis (I70)	95	6.7	5	7.5	0	0.0	1	2.6	28	7.1	61	6.6
Other diseases of circulatory system (I71-I78)	127	7.9	6	5.2	0	0.0	5	23.0	33	7.1	83	8.2
Influenza and pneumonia (J10-J18)	349	23.0	39	42.6	1	3.1	4	14.3	114	25.6	190	20.1
Chronic lower respiratory diseases (J40-J47)	897	52.1	19	24.6	2	27.1	7	23.7	204	41.7	661	60.5
Peptic ulcer (K25-K28)	20	1.0	1	0.5	0	0.0	1	2.4	5	0.7	13	1.2
Chronic liver disease and cirrhosis (K70, K73-K74)	383	17.8	90	51.1	1	4.1	3	7.4	176	23.6	111	9.5
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	249	14.6	23	19.4	0	0.0	8	21.4	88	16.9	130	12.6
Pregnancy, childbirth, and the puerperium (O00-O99)	4	0.2	1	0.5	0	0.0	0	0.0	3	0.4	0	0.0
Certain conditions originating in the perinatal period (P00-P96)	83	4.3	10	3.9	1	3.0	3	6.1	49	4.9	20	3.4
Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99)	76	3.9	12	4.7	0	0.0	4	8.4	34	3.9	25	3.4
Sudden infant death syndrome (R95)	15	0.8	4	1.6	0	0.0	1	2.0	5	0.5	5	0.9
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	68	4.1	5	3.8	1	2.7	1	2.6	16	2.2	45	5.2
All other diseases (Residual)	1,748	109.5	156	144.1	12	116.6	22	100.4	537	109.3	1,014	105.1
Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2)	366	17.7	72	31.4	3	7.9	9	17.8	141	16.8	137	14.6
All other and unspecified accidents and adverse effects	910	51.7	88	49.9	4	50.5	12	28.8	332	49.7	468	50.2
Intentional self-harm (suicide) (X60-X84, Y87.0)	400	19.5	52	22.3	3	8.1	4	6.5	138	16.9	202	20.4
Assault (homicide) (X85-Y09, Y87.1)	164	8.0	26	10.8	3	12.0	10	16.4	84	9.6	41	4.9
All other external causes (Y10-Y36, Y87.2, Y89)	103	5.1	23	11.4	0	0.0	3	6.1	40	4.5	36	4.0

Other and unknown races, if any, are included in the "All Races" category. Numbers of deaths may not sum to totals, due to unknowns. Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

New Mexico Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico. <http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

Mortality Section

Table M-19
Number of Injury Deaths by Sex, County and Health Region
New Mexico, 2007, and United States, 2006

Resident County	Total Injuries (Unintentional and Intentional)			Accidents (Unintentional Injuries)						Intentional Injuries						Other		
				Motor vehicle accidents			All other accidents			Intentional self harm (Suicide)			Assault (Homicide)			Other injury*		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	181,586	124,781	56,805	45,316	31,633	13,683	76,283	47,308	28,975	33,300	26,308	6,992	18,573	14,717	3,856	8,114	4,815	3,299
New Mexico	1,946	1,363	583	367	260	107	910	588	322	400	326	74	164	126	38	105	63	42
Bernalillo	601	407	194	89	67	22	307	186	121	124	94	30	56	43	13	25	17	8
Catron	8	7	1	0	0	0	3	3	0	4	4	0	0	0	0	1	0	1
Chaves	53	36	17	8	5	3	24	13	11	10	8	2	9	9	0	2	1	1
Cibola	32	27	5	9	9	0	15	11	4	6	5	1	2	2	0	0	0	0
Colfax	15	12	3	2	2	0	6	4	2	5	4	1	1	1	0	1	1	0
Curry	38	25	13	8	5	3	20	12	8	5	4	1	2	2	0	3	2	1
De Baca	3	2	1	1	0	1	1	1	0	1	1	0	0	0	0	0	0	0
Doña Ana	130	85	45	19	15	4	70	43	27	26	20	6	7	5	2	8	2	6
Eddy	46	32	14	7	6	1	25	16	9	9	7	2	4	3	1	1	0	1
Grant	35	24	11	6	6	0	16	9	7	5	5	0	4	4	0	4	0	4
Guadalupe	5	2	3	3	2	1	0	0	0	0	0	0	2	0	2	0	0	0
Harding	3	3	0	0	0	0	3	3	0	0	0	0	0	0	0	0	0	0
Hidalgo	6	5	1	1	1	0	1	0	1	3	3	0	0	0	0	1	1	0
Lea	57	40	17	19	13	6	20	14	6	9	9	0	7	3	4	2	1	1
Lincoln	29	18	11	5	3	2	16	9	7	3	2	1	1	1	0	4	3	1
Los Alamos	12	10	2	2	1	1	5	4	1	4	4	0	0	0	0	1	1	0
Luna	27	21	6	10	6	4	9	7	2	4	4	0	2	2	0	2	2	0
McKinley	102	80	22	25	16	9	42	31	11	17	17	0	12	11	1	6	5	1
Mora	5	4	1	1	0	1	2	2	0	1	1	0	0	0	0	1	1	0
Otero	44	34	10	5	5	0	16	14	2	16	11	5	3	2	1	4	2	2
Quay	12	6	6	2	0	2	3	2	1	3	3	0	4	1	3	0	0	0
Rio Arriba	73	55	18	19	13	6	29	21	8	15	14	1	6	5	1	4	2	2
Roosevelt	16	9	7	2	1	1	10	6	4	2	1	1	2	1	1	0	0	0
San Juan	170	113	57	45	31	14	67	38	29	25	21	4	18	12	6	15	11	4
San Miguel	37	30	7	6	5	1	14	10	4	9	8	1	7	6	1	1	1	0
Sandoval	87	59	28	10	7	3	44	25	19	26	22	4	5	4	1	2	1	1
Santa Fe	118	86	32	19	13	6	66	48	18	25	20	5	3	3	0	5	2	3
Sierra	15	10	5	2	1	1	9	5	4	2	2	0	1	1	0	1	1	0
Socorro	17	15	2	4	4	0	6	4	2	6	6	0	0	0	0	1	1	0
Taos	49	35	14	11	7	4	20	17	3	13	9	4	1	1	0	4	1	3
Torrance	17	10	7	6	3	3	5	1	4	5	5	0	1	1	0	0	0	0
Union	2	2	0	1	1	0	0	0	0	1	1	0	0	0	0	0	0	0
Valencia	79	57	22	20	12	8	34	27	7	16	11	5	3	3	0	6	4	2
Health Region																		
Northwest	470	336	134	109	75	34	202	132	70	90	76	14	40	32	8	29	21	8
Northeast	316	236	80	64	44	20	142	106	36	73	61	12	20	16	4	17	9	8
Bernalillo County	601	407	194	89	67	22	307	186	121	124	94	30	56	43	13	25	17	8
Southeast	228	153	75	47	30	17	106	67	39	39	33	6	28	19	9	8	4	4
Southwest	328	229	99	58	44	14	151	95	56	74	62	12	19	16	3	26	12	14

Deaths with unknown sex or county of residence, if any, are included in the "New Mexico Total" and "Both Sexes" categories. Numbers of deaths may not sum to total due to unknowns.

*Other Injury includes Legal intervention, Operations of war, Complications of medical or surgical care, and Events of undetermined intent.

The U.S. numbers for Total Injuries do not include Complications of medical or surgical care.

For counties included in the state health regions, please see the Technical Appendix.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 17, 2008. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-20
Death Rates for Injury by Sex, County and Health Region
New Mexico, 2007, and United States, 2006

Resident County	Total Injuries (Unintentional and Intentional)			Accidents (Unintentional Injuries)						Intentional Injuries						Other		
				Motor vehicle accidents			All other accidents			Intentional self harm (Suicide)			Assault (Homicide)			Other injury*		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
United States	58.8**	N/A	N/A	15.0	21.4	8.8	N/A	N/A	N/A	10.9	18.0	4.5	6.2	9.7	2.5	N/A	N/A	N/A
New Mexico	102.2	145.7	61.1	17.8	25.5	10.3	51.7	68.5	36.1	19.5	32.9	7.0	8.0	12.5	3.7	5.2	6.4	4.0
Bernalillo	98.0	133.8	63.6	13.9	21.3	6.9	52.7	63.9	41.7	18.7	29.4	8.6	8.6	13.5	3.8	4.0	5.7	2.6
Catron	174.4	331.6	23.7	0.0	0.0	0.0	70.3	165.2	0.0	92.7	166.4	0.0	0.0	0.0	0.0	11.5	0.0	23.7
Chaves	83.4	124.6	45.8	13.2	17.0	9.3	39.1	52.9	27.8	15.1	24.9	6.6	13.5	27.2	0.0	2.5	2.6	2.1
Cibola	117.2	216.1	29.4	29.1	60.3	0.0	59.8	103.6	23.1	20.3	35.5	6.4	8.0	16.7	0.0	0.0	0.0	0.0
Coffax	109.5	184.0	37.4	11.5	21.3	0.0	43.4	71.0	21.5	36.8	57.8	15.9	7.1	12.9	0.0	10.7	20.9	0.0
Curry	96.9	127.6	67.5	17.1	21.4	13.2	56.8	69.0	45.1	11.2	17.6	4.7	4.8	9.8	0.0	6.9	9.8	4.4
De Baca	161.2	191.6	112.8	61.7	0.0	112.8	45.7	80.3	0.0	53.8	111.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Doña Ana	75.6	108.3	47.1	8.7	14.2	3.4	45.5	64.4	30.0	13.8	22.7	6.1	3.5	4.7	2.2	4.0	2.3	5.5
Eddy	93.9	131.6	57.0	12.7	21.1	4.3	52.5	70.7	34.8	18.3	28.9	8.3	8.2	11.0	5.2	2.2	0.0	4.5
Grant	110.3	148.2	68.1	17.7	35.9	0.0	53.3	56.7	44.2	14.9	31.6	0.0	11.6	24.0	0.0	12.7	0.0	23.9
Guadalupe	87.6	61.5	133.9	52.0	61.5	49.6	0.0	0.0	0.0	0.0	0.0	0.0	35.6	0.0	84.3	0.0	0.0	0.0
Harding	425.7	952.4	0.0	0.0	0.0	0.0	425.7	952.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	120.3	190.2	43.1	25.1	54.2	0.0	26.7	0.0	43.1	48.6	97.0	0.0	0.0	0.0	0.0	19.8	39.0	0.0
Lea	99.2	142.8	56.9	30.0	40.8	18.6	37.6	54.3	18.4	16.9	35.5	0.0	11.6	9.0	14.4	3.1	3.3	2.9
Lincoln	139.7	188.8	92.7	27.1	38.3	15.9	73.2	81.5	67.2	16.4	29.1	4.8	4.9	9.2	0.0	18.1	30.8	4.7
Los Alamos	59.6	103.7	14.7	10.0	11.7	7.4	22.6	37.3	7.4	18.2	37.1	0.0	0.0	0.0	0.0	8.7	17.6	0.0
Luna	97.7	159.3	42.0	36.1	48.7	24.9	33.2	51.7	17.1	13.1	27.4	0.0	7.7	16.1	0.0	7.5	15.4	0.0
McKinley	151.2	249.7	62.8	34.0	46.4	22.8	70.6	111.6	34.8	24.2	51.0	0.0	14.8	27.7	2.8	7.6	13.1	2.4
Mora	83.2	126.7	30.9	14.8	0.0	30.9	26.0	49.0	0.0	13.0	24.5	0.0	0.0	0.0	0.0	29.4	53.1	0.0
Otero	74.2	115.3	36.5	8.4	16.8	0.0	30.5	49.7	13.0	24.9	37.4	13.8	4.8	5.9	3.8	5.6	5.5	5.8
Quay	109.6	131.7	94.5	13.7	0.0	25.7	33.4	54.2	18.9	19.9	40.5	0.0	42.6	36.9	49.8	0.0	0.0	0.0
Rio Arriba	170.8	241.6	93.8	43.1	58.3	27.0	70.8	89.4	47.1	34.4	63.4	5.2	13.8	22.5	4.9	8.7	8.1	9.5
Roosevelt	90.8	105.4	76.0	9.0	10.4	7.3	59.6	73.8	44.9	10.4	10.8	10.7	11.8	10.3	13.0	0.0	0.0	0.0
San Juan	139.6	184.9	95.2	35.6	48.8	22.6	59.8	66.7	52.1	18.0	29.9	6.3	14.7	22.1	8.5	11.5	17.4	5.7
San Miguel	122.7	199.8	49.7	20.3	34.8	6.0	44.7	64.0	26.3	29.5	52.2	8.7	23.7	40.1	8.7	4.5	8.6	0.0
Sandoval	87.1	120.8	56.3	7.5	10.8	4.3	52.2	63.2	42.5	21.8	38.5	6.6	3.8	6.2	1.4	1.8	2.1	1.4
Santa Fe	92.0	143.9	47.0	12.0	17.0	7.1	58.6	93.3	30.1	15.6	26.0	5.9	2.3	4.7	0.0	3.4	2.9	3.9
Sierra	80.5	92.4	70.0	7.7	11.0	5.2	51.9	38.0	64.9	8.5	17.4	0.0	7.5	15.1	0.0	5.0	11.0	0.0
Socorro	100.8	162.0	37.0	17.7	34.4	0.0	43.7	46.9	37.0	32.9	68.3	0.0	0.0	0.0	0.0	6.4	12.4	0.0
Taos	168.2	252.8	89.8	40.6	50.3	30.3	66.2	116.8	18.1	43.5	69.2	22.6	4.6	8.7	0.0	13.3	7.8	18.9
Torrance	141.9	120.4	132.6	42.8	44.2	41.9	64.0	9.4	90.7	28.3	54.7	0.0	6.8	12.1	0.0	0.0	0.0	0.0
Union	62.3	113.2	0.0	22.9	42.9	0.0	0.0	0.0	0.0	39.4	70.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Valencia	120.4	183.9	62.7	28.1	34.7	22.8	56.9	96.2	20.4	23.1	34.5	13.1	4.2	8.6	0.0	8.2	10.0	6.3
Health Region																		
Northwest	121.8	178.7	69.3	25.7	36.0	16.1	58.6	79.9	39.5	21.2	37.2	6.4	9.4	15.5	3.5	6.9	10.1	3.8
Northeast	111.6	171.2	55.8	21.3	29.5	12.8	53.0	82.2	26.7	24.2	41.5	8.0	7.1	11.3	3.0	6.0	6.7	5.3
Bernalillo County	98.0	133.8	63.6	13.9	21.3	6.9	52.7	63.9	41.7	18.7	29.4	8.6	8.6	13.5	3.8	4.0	5.7	2.6
Southeast	94.4	132.3	58.7	17.6	22.6	12.6	47.0	64.4	30.9	15.7	27.3	4.9	11.1	14.8	7.3	3.1	3.3	3.0
Southwest	87.4	123.5	53.2	13.6	21.0	6.2	45.0	57.5	33.3	18.1	31.5	5.7	4.6	7.7	1.5	6.2	5.8	6.5

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard populations.

Deaths with unknown sex or county of residence, if any, are included in the "New Mexico Total" and "Both Sexes" categories. Numbers of deaths may not sum to total due to unknowns.

*Other Injury includes Legal intervention, Operations of war, Complications of medical or surgical care, and Events of undetermined intent.

**The U.S. numbers for Total Injuries do not include complications of medical or surgical care.

For counties included in the state health regions, please see the Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol 57, Number 14, released April 17, 2009. <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-21
Number of Deaths due to Firearms, Alcohol, Drugs, and Injury at Work
by Sex, County and Health Region
New Mexico, 2007, and United States, 2006

Resident County	Firearms			Alcohol-induced			Drug-induced			Injury at work		
	Both Sexes	Males	Females	Both Sexes	Males	Females	Both Sexes	Males	Females	Both Sexes	Males	Females
United States	30,896	26,712	4,184	22,073	16,472	5,601	38,396	24,507	13,889	5,298	4,869	429
New Mexico	291	248	43	408	303	105	462	311	151	39	36	3
Bernalillo	95	77	18	136	102	34	181	120	61	9	8	1
Catron	3	3	0	1	0	1	2	2	0	0	0	0
Chaves	10	10	0	8	5	3	13	6	7	2	2	0
Cibola	2	2	0	8	5	3	6	5	1	1	1	0
Colfax	2	2	0	4	4	0	3	3	0	0	0	0
Curry	3	3	0	6	4	2	8	7	1	1	1	0
De Baca	1	1	0	0	0	0	1	1	0	0	0	0
Doña Ana	23	19	4	13	13	0	33	22	11	1	1	0
Eddy	9	8	1	7	7	0	14	9	5	3	3	0
Grant	7	6	1	8	6	2	6	5	1	0	0	0
Guadalupe	2	0	2	1	0	1	0	0	0	0	0	0
Harding	0	0	0	0	0	0	0	0	0	1	1	0
Hidalgo	3	3	0	2	2	0	1	1	0	0	0	0
Lea	10	8	2	5	4	1	14	10	4	5	4	1
Lincoln	3	2	1	5	3	2	6	4	2	1	1	0
Los Alamos	1	1	0	0	0	0	3	2	1	0	0	0
Luna	7	7	0	0	0	0	2	0	2	0	0	0
McKinley	4	4	0	41	31	10	8	6	2	0	0	0
Mora	0	0	0	0	0	0	1	1	0	0	0	0
Otero	10	9	1	5	2	3	10	7	3	2	2	0
Quay	5	4	1	6	4	2	1	0	1	0	0	0
Rio Arriba	9	9	0	26	20	6	20	13	7	2	1	1
Roosevelt	2	1	1	0	0	0	4	4	0	0	0	0
San Juan	16	14	2	28	17	11	21	9	12	7	7	0
San Miguel	6	5	1	8	4	4	9	6	3	0	0	0
Sandoval	18	15	3	25	20	5	21	13	8	1	1	0
Santa Fe	16	15	1	33	26	7	32	24	8	1	1	0
Sierra	1	1	0	3	3	0	2	2	0	0	0	0
Socorro	2	2	0	5	2	3	4	3	1	0	0	0
Taos	8	6	2	10	8	2	10	6	4	1	1	0
Torrance	2	2	0	1	1	0	2	1	1	0	0	0
Union	1	1	0	0	0	0	1	1	0	0	0	0
Valencia	10	8	2	13	10	3	23	18	5	1	1	0
Health Region												
Northwest	50	43	7	115	83	32	79	51	28	10	10	0
Northeast	45	39	6	82	62	20	78	55	23	4	3	1
Bernalillo County	95	77	18	136	102	34	181	120	61	9	8	1
Southeast	40	35	5	32	24	8	55	37	18	12	11	1
Southwest	61	54	7	43	32	11	68	47	21	4	4	0

The categories presented are not mutually exclusive.
 Injury at work deaths include all deaths (ages 15 and over) due to external causes that occurred while the decedent was working.
 Deaths with unknown county of residence, if any, are included in the total categories. Numbers may not add to totals due to unknowns.
 For counties included in state health regions, please see Technical Appendix.
 U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol. 57, No. 14, <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-22
Death Rates due to Firearms, Alcohol or Drugs, and Injury at Work
by Sex, County and Health Region
New Mexico, 2007, and United States, 2006

County	Firearms			Alcohol-induced			Drug-induced			Injury at work		
	Both Sexes	Males	Females	Both Sexes	Males	Females	Both Sexes	Males	Females	Both Sexes	Males	Females
United States	10.3	18.1	2.8	7.4	11.2	3.7	12.8	16.6	9.1	2.2	4.2	0.4
New Mexico	14.3	25.2	4.1	19.0	29.0	9.7	22.3	30.6	14.3	1.8	3.4	0.3
Bernalillo	14.8	24.7	5.5	19.3	29.7	9.6	26.7	35.9	17.8	1.3	2.4	0.3
Catron	74.7	131.3	0.0	11.5	0.0	23.7	53.3	125.1	0.0	0.0	0.0	0.0
Chaves	15.6	31.4	0.0	12.0	15.2	9.1	21.1	20.2	21.9	3.2	6.2	0.0
Cibola	7.2	15.2	0.0	25.6	34.8	17.3	21.2	35.4	7.2	3.0	6.4	0.0
Colfax	21.4	41.8	0.0	18.6	37.2	0.0	19.0	36.9	0.0	0.0	0.0	0.0
Curry	6.9	13.8	0.0	12.7	18.1	8.1	17.2	31.9	4.7	2.4	4.9	0.0
De Baca	53.8	111.4	0.0	0.0	0.0	0.0	45.7	80.3	0.0	0.0	0.0	0.0
Doña Ana	12.6	21.7	4.4	6.7	14.3	0.0	17.8	25.2	11.1	0.4	0.9	0.0
Eddy	18.8	33.3	5.2	14.0	29.3	0.0	29.4	37.3	21.3	4.7	9.1	0.0
Grant	20.1	35.7	5.2	22.4	34.7	11.5	20.0	33.1	8.2	0.0	0.0	0.0
Guadalupe	35.6	0.0	84.3	86.2	0.0	155.1	0.0	0.0	0.0	0.0	0.0	0.0
Harding	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	294.7	616.2	0.0
Hidalgo	48.6	97.0	0.0	35.7	70.4	0.0	19.8	39.0	0.0	0.0	0.0	0.0
Lea	16.9	26.0	7.9	8.3	13.6	3.1	24.1	34.6	12.9	7.8	12.4	3.4
Lincoln	12.4	18.4	4.8	16.9	23.1	11.4	24.6	34.1	17.1	3.5	7.2	0.0
Los Alamos	7.6	16.0	0.0	0.0	0.0	0.0	16.3	23.5	7.4	0.0	0.0	0.0
Luna	25.0	52.0	0.0	0.0	0.0	0.0	8.8	0.0	17.1	0.0	0.0	0.0
McKinley	6.2	13.5	0.0	60.8	98.6	28.2	9.8	15.4	4.3	0.0	0.0	0.0
Mora	0.0	0.0	0.0	0.0	0.0	0.0	13.0	24.5	0.0	0.0	0.0	0.0
Otero	16.4	31.4	2.8	6.6	5.5	7.6	15.0	21.6	8.6	2.9	5.9	0.0
Quay	47.9	77.4	21.3	45.2	56.2	33.9	12.5	0.0	18.9	0.0	0.0	0.0
Rio Arriba	20.8	41.2	0.0	56.1	87.0	25.8	43.6	56.3	30.8	4.4	3.9	4.9
Roosevelt	10.4	10.8	10.7	0.0	0.0	0.0	21.7	43.6	0.0	0.0	0.0	0.0
San Juan	11.7	21.9	2.4	20.6	25.0	15.6	17.0	12.4	20.1	5.3	10.8	0.0
San Miguel	18.8	30.1	8.7	24.0	29.4	21.0	28.8	40.1	18.6	0.0	0.0	0.0
Sandoval	14.8	25.6	4.9	19.8	32.6	7.5	17.4	22.8	12.5	0.7	1.5	0.0
Santa Fe	10.8	20.5	1.6	20.4	32.8	9.0	20.1	31.3	9.4	0.5	1.0	0.0
Sierra	5.0	11.0	0.0	13.4	28.3	0.0	10.0	20.1	0.0	0.0	0.0	0.0
Socorro	11.4	23.3	0.0	31.3	24.9	37.3	20.8	32.5	9.5	0.0	0.0	0.0
Taos	31.3	55.5	11.3	26.5	42.3	11.3	30.3	42.5	16.4	2.0	4.2	0.0
Torrance	12.1	23.5	0.0	5.1	10.2	0.0	11.4	10.9	11.8	0.0	0.0	0.0
Union	39.4	70.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Valencia	16.5	28.7	5.7	16.8	26.7	7.7	31.2	48.9	13.1	1.6	3.1	0.0
Health Region												
Northwest	12.1	22.1	3.1	26.8	39.5	14.7	18.8	24.7	13.1	2.3	4.7	0.0
Northeast	16.3	29.1	4.4	24.4	37.2	12.3	24.1	35.2	12.9	1.1	1.5	0.7
Bernalillo County	14.8	24.7	5.5	19.3	29.7	9.6	26.7	35.9	17.8	1.3	2.4	0.3
Southeast	15.7	27.7	4.0	12.2	18.9	6.0	22.5	30.0	14.8	4.6	8.4	0.7
Southwest	14.9	27.0	3.4	10.3	15.9	5.2	17.4	25.2	10.4	0.9	2.0	0.0

Age-adjusted rates are the numbers of deaths per 100,000 U.S. standard population.

The categories presented are not mutually exclusive.

Injury at work deaths include all deaths (ages 15 and over) due to external causes that occurred while the decedent was working.

For counties included in state health regions, please see Technical Appendix.

Population Source: Bureau of Business and Economic Research (BBER) Population Estimates, University of New Mexico.

<http://www.unm.edu/~bber/>. BBER updated its methodology beginning with the population estimates for 2007. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the changes in population estimates used in the rate denominators.

U.S. Data Source: CDC, National Center for Health Statistics. NVSR, Vol. 57, No. 14 released April 17, 2009. <http://www.cdc.gov/nchs/>.

Table M-23
Numbers and Rates of Maternal Mortality by Age
New Mexico, 2000-2003 and 2004-2007 Averages

Age Group	2000-2003		2004-2007	
	(4-year) Total	Average Rate	(4-year) Total	Average Rate
All Ages	24	21.9	11	9.3
10 to 14 Years	0	0.0	0	0.0
15 to 19 Years	4	22.0	1	5.5
20 to 24 Years	2	5.9	1	2.7
25 to 29 Years	3	11.0	4	13.1
30 to 34 Years	4	21.5	3	15.1
35 to 39 Years	6	65.7	2	20.5
40 to 49 Years	5	218.8	0	0.0

Maternal mortality rates are the numbers of maternal deaths per 100,000 live births. The rate numerators are total maternal deaths for the specified 4-year period divided by 4. Rate denominators are the total births for the specified 4 year period divided by 4.

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Other and unknown ages, if any, are included in the "All Ages" category.

A maternal death is defined as the death of a woman while pregnant or within 42 days of pregnancy, excluding external injury deaths. For more information on maternal death, please see Technical Appendix.

Table M-24
Numbers and Average Rates of Maternal Mortality by Cause
New Mexico, 2000-2003 and 2004-2007

Maternal Cause of Death (ICD-10 Code) <i>10th Revision International Classification of Diseases</i>	2000-2003		2004-2007	
	(4-year) Total	Average Rate	(4-year) Total	Average Rate
Total Maternal Causes (A34, O00-O95, O98-O99)	24	21.9	11	9.3
Pregnancy with abortive outcomes (O00-O08)	3	2.7	0	0.0
Ectopic pregnancy (O00)	1	0.9	0	0.0
Spontaneous abortion (O03)	0	0.0	0	0.0
Medical abortion (O04)	0	0.0	0	0.0
Other abortion (O05)	0	0.0	0	0.0
Other & unspecified pregnancy with abortive outcomes (O01-O02, O06-O07)	1	0.9	0	0.0
Complications following abortion and ectopic and molar pregnancy (O08)	1	0.9	0	0.0
Other direct obstetric causes (A34, O10-O92)	17	15.5	9	7.6
Eclampsia and pre-eclampsia (O11, O13-O16)	6	5.5	0	0.0
Hemorrhage of pregnancy and childbirth and placenta previa (O20, O44-O46, O67, O72)	3	2.7	2	1.7
Complications predominantly related to puerperium (A34, O85-O92)	2	1.8	4	3.4
Obstetrical tetanus (A34)	0	0.0	0	0.0
Obstetrical embolism (O88)	1	0.9	2	1.7
Other complications predominantly related to the puerperium (O85-O87, O89-O92)	1	0.9	2	1.7
All other direct obstetric causes (O10, O12, O21-O43, O47-O66, O68-O71, O73-O75)	6	5.5	3	2.5
Obstetric death of unspecified cause (O95)	0	0.0	1	0.8
Indirect obstetric causes (O98-O99)	4	3.6	1	0.8

Maternal mortality rates are the numbers of maternal deaths per 100,000 live births. The rate numerators are total maternal deaths for the specified 4-year period divided by 4. Rate denominators are the total births for the specified 4-year period divided by 4.

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

A maternal death is defined as the death of a woman while pregnant or within 42 days of pregnancy, excluding external injury deaths. For more information on maternal death, please see Technical Appendix.

Table M-25
Numbers and Rates of Infant Mortality by Sex and Race/Ethnicity
New Mexico, 2007

Race/Ethnicity	2007					
	Both Sexes		Male		Female	
	#	Rate	#	Rate	#	Rate
All Races						
Infant Deaths	188	6.1	112	7.1	76	5.1
Neonatal Deaths	116	3.8	73	4.6	43	2.9
Postneonatal Deaths	72	2.4	39	2.5	33	2.2
American Indian or Alaska Native						
Infant Deaths	30	7.6	18	9.2	12	6.1
Neonatal Deaths	13	3.3	8	4.1	5	2.6
Postneonatal Deaths	17	4.3	10	5.1	7	3.6
Asian or Pacific Islander						
Infant Deaths	1	2.0	1	3.6	0	0.0
Neonatal Deaths	1	2.0	1	3.6	0	0.0
Postneonatal Deaths	0	0.0	0	0.0	0	0.0
Black or African American						
Infant Deaths	9	14.1	6	17.1	3	10.4
Neonatal Deaths	6	9.4	4	11.4	2	6.9
Postneonatal Deaths	3	4.7	2	5.7	1	3.5
Hispanic						
Infant Deaths	105	6.3	59	6.9	46	5.7
Neonatal Deaths	67	4.0	39	4.6	28	3.4
Postneonatal Deaths	38	2.3	20	2.4	18	2.2
White						
Infant Deaths	42	4.8	28	6.1	14	3.3
Neonatal Deaths	28	3.2	21	4.6	7	1.7
Postneonatal Deaths	14	1.6	7	1.5	7	1.7

Infant mortality rates are the numbers of infant deaths under 1 year of age per 1,000 live births.

Infant deaths based on race/ethnicity of decedent, as reported on the death certificate. Rate denominator uses race/ethnicity of mother as reported on the birth certificate. Neonatal = Under 28 days. Postneonatal = 28 days to under 1 year. For further explanation of neonatal and postneonatal periods, please see the Technical Appendix.

Notes: The Hispanic category does not include American Indian, Asian or Pacific Islander or Black decedents. Other and unknown races, if any, are included in the "All Races" category.

Table M-26
Numbers of Infant Deaths by Age and Cause
New Mexico, 2007, and United States, 2006

Cause of Death	ICD-10 Code - 10th Revision International Classification of Diseases	United States 2006	New Mexico 2007	New Mexico			
				Under 1 Day	1 to 6 Days	7 to 27 Days	28 Days to Under 1 Year
All Causes		28,527	188	52	34	30	72
Certain infectious & parasitic diseases	A00-B99	479	8	1	0	1	6
-Whooping cough	A37	8	0	0	0	0	0
-Meningococcal infection	A39	11	0	0	0	0	0
-Septicemia	A40-A41	269	2	1	0	0	1
-Human immunodeficiency virus (HIV) disease	B20-B24	7	0	0	0	0	0
-Other viral diseases	A80-B19, B25-B34	113	3	0	0	1	2
-Remainder of certain infectious & parasitic diseases	A00-A36, A38, A42-A79, B35-B99	71	3	0	0	0	3
Neoplasms	C00-D48	141	0	0	0	0	0
-Malignant neoplasms	C00-C97	76	0	0	0	0	0
-Remainder of neoplasms	D00-D48	65	0	0	0	0	0
Diseases of the blood & blood-forming organs & certain disorders involving the immune mechanism	D50-D89	102	0	0	0	0	0
Endocrine, nutritional & metabolic diseases	E00-E88	207	1	0	1	0	0
Diseases of the nervous system	G00-G98	373	7	1	0	1	5
-Meningitis	G00, G03	61	2	0	0	0	2
-Remainder of diseases of the nervous system	G04-G98	312	5	1	0	1	3
Diseases of the ear & mastoid process	H60-H93	3	0	0	0	0	0
Diseases of the circulatory system	I00-I99	543	5	0	1	0	4
Diseases of the respiratory system	J00-J98	692	1	0	0	1	0
-Pneumonia	J12-J18	245	1	0	0	1	0
-Remainder of diseases of the respiratory system	J00-J11, J19-98	447	0	0	0	0	0
Diseases of the digestive system	K00-K92	582	4	0	0	1	3
Disease of the genitourinary system	N00-N95	180	2	0	0	1	1
Certain conditions originating in the perinatal period	P00-P96	14,321	82	38	20	17	7
-Maternal factors & complications of pregnancy, labor & delivery	P00-P04	3,150	21	16	3	1	1
---- Maternal complications of pregnancy	P01	1,683	9	8	1	0	0
----Complications of placenta, cord, & membranes	P02	1,140	7	4	2	1	0
----Other maternal factors & complications of pregnancy	P00-P09, P03-P04	327	5	4	0	0	1
-Disorders relating to length of gestation & fetal growth	P05-P08	4,940	40	16	12	8	4
----Disorders relating to short gestation & low birth weight	P07	4,841	39	16	12	7	4
-Intrauterine hypoxia & birth asphyxia	P20-P21	344	0	0	0	0	0
-Respiratory distress of newborn	P22	825	3	1	1	0	1
-Other respiratory conditions of newborn	P23-P28	1,207	5	3	1	1	0
-Bacterial sepsis of newborn	P36	807	2	0	0	2	0
-Hemorrhagic & hematological disorders of newborn	P50-P61	725	2	0	1	1	0
-Remainder of perinatal conditions	P10-P15, P29, P35, P37-P39, P70-P96	2,323	9	2	2	4	1
Congenital malformations, deformations & chromosomal abnormalities	Q00-Q99	5,819	40	10	11	5	14
-Congenital hydrocephalus	Q03	88	0	0	0	0	0
-Spina bifida	Q05	23	0	0	0	0	0
-Other congenital malformations of nervous system	Q00-Q02, Q04, Q06-Q07	390	6	5	0	0	1
-Congenital malformations of the heart	Q20-Q24	1,396	11	0	3	3	5
-Other congenital malformations of the circulatory system	Q25-Q28	236	0	0	0	0	0
-Congenital malformations of respiratory system	Q30-Q34	437	2	0	1	1	0
-Congenital malformation & deformations of musculoskeletal system, limbs & integument	Q65-Q85	619	7	3	2	1	1
-Down's syndrome & other chromosomal abnormalities	Q90-Q99	1,130	4	0	1	0	3
-Other congenital malformations	Q10-Q18, Q35-Q64, Q86-Q89	1,164	10	2	4	0	4
Symptoms, signs & abnormal clinical & lab findings	R00-R99	3,462	15	1	0	1	13
-Sudden infant death syndrome (SIDS)	R95	2,323	14	0	0	1	13
Accidents (Unintentional Injuries)	V01-X59	1,147	4	1	0	0	3
Assault (Homicide)	X85-Y09	336	6	0	0	0	6
Complications of medical and surgical care	Y40-Y84	23	0	0	0	0	0
Other external causes	Y10-Y36	92	12	0	1	2	9
Other causes not listed		25	1	0	0	0	1

Infant mortality refers to the numbers of infant deaths under 1 year of age.

U.S. Data Source: CDC, National Center for Health Statistics; NVSR, Vol. 57, No. 14, released 04/17/09, <http://www.cdc.gov/nchs/>. 2007 U.S. data not available at time of publication.

Table M-27
Method of Final Disposition for Deaths Occurring in New Mexico
New Mexico, 1998-2007

Year	Total	Burial		Cremation		Removal		Other and Unspecified	
	Number	Number	Percent	Number	Percent	Number	Percent	Number	Percent
2007	15,160	6,815	45.0	7,454	49.2	599	4.0	292	1.9
2006	15,044	7,021	46.7	7,169	47.7	597	4.0	257	1.7
2005	14,755	6,786	46.0	6,787	46.0	973	6.6	209	1.4
2004	14,046	6,556	44.4	6,193	42.0	1,073	7.3	224	1.5
2003	14,548	6,955	49.5	6,305	44.9	1,039	7.4	249	1.8
2002	14,175	7,038	48.4	5,867	40.3	1,070	7.4	200	1.4
2001	13,995	6,809	48.0	5,561	39.2	1,393	9.8	232	1.6
2000	13,250	6,518	46.6	5,072	36.2	1,461	10.4	199	1.4
1999	13,607	7,032	53.1	4,832	36.5	1,547	11.7	196	1.5
1998	12,804	6,688	49.2	4,386	32.2	1,511	11.1	219	1.6

Due to rounding, percent may not add to 100.

Note: "Removal" refers to removal of remains from New Mexico to another state or country. "Other and Unspecified" includes donation, entombment, and all other and unspecified means of disposition.

Technical Appendix



NEW MEXICO VITAL STATISTICS DATA SOURCES

DEATH DATA

Mortality statistics are based on items reported on the death certificate and include demographic, geographic, injury, medical and cause-of-death information on the decedent. In most cases, funeral directors work with an informant for the decedent (usually a relative or friend) to collect demographic and geographic information. The medical and cause-of-death section of the death certificate is completed by the attending physician or the Office of the Medical Investigator (OMI). Except for deaths occurring on tribal or military lands, the Office of the Medical Investigator has jurisdiction over all unexpected and unattended deaths in New Mexico.

In 2007 there were 15,400 deaths among New Mexico residents. To reflect the health status of New Mexicans, most data in this report are presented by New Mexico residence rather than occurrence. To facilitate reporting by residence, states, territories and other jurisdictions engage in an inter-jurisdictional exchange process. When deaths to New Mexico residents occur out of state, information from the death certificate is transmitted to NMVRHS for statistical reporting.

STATISTICAL FILE TIMELINE

To allow sufficient time to receive as many vital records as possible, including those that occur out of state, NMVRHS keeps the statistical files open for approximately 10 to 12 months after the end of the calendar year. At that time, the statistical file is closed so that statistics for that year can be consistently generated in subsequent years. Because some states experienced significant delays in providing death data for New Mexico residents that occurred in those states, the publication of New Mexico 2007 death data was later than usual.

STANDARD CERTIFICATES

Beginning in 2006, New Mexico adopted the 2003 revision of the United States Standard Death Certificate with the addition of some state-specific items.

OTHER DATA SOURCES

NATIONAL VITAL STATISTICS DATA

National vital statistics data are produced by the United States Centers for Disease Control's National Center for Health Statistics (NCHS). As part of the national vital statistics system, states, territories, and other jurisdictions provide birth, death, and fetal death data to NCHS. At the time of this report's publication, 2006 was the latest data year for U.S. birth and death information available from NCHS.

POPULATION DATA

Population estimates used as the denominators in calculating birth and death rates in this report were produced by the University of New Mexico's Bureau of Business and Economic Research (BBER) for years 2000 to 2007, and by the United States Census Bureau, in collaboration with NCHS, for previous years. National vital statistics data use population estimates and census counts produced by the Census Bureau.

Beginning with the population estimates for 2007, the University of New Mexico Bureau of Business and Economic Research (BBER) updated its estimates of age-specific migration with new data. This change caused a break in continuity between the 2006 and 2007 estimates. Older age groups were affected more than others. For rates calculated from BBER population estimates, any rate differences from 2006 to 2007 must be interpreted with caution, as they may be partly attributable to the change in population estimates used in the rate denominators. For additional information, see <http://ibis.health.state.nm.us/docs/Query/Pop/2007PopDataTechnicalNote.doc>.

The Demographic and Socioeconomic Section of this report contains additional population data from both BBER and the United States Census Bureau. Population counts are numbers from the decennial census, while population estimates (usually intercensal) are calculated using decennial counts as well as birth, death and migration data. Population projections use population growth trends to calculate population estimates for the future.

DEMOGRAPHIC AND SOCIOECONOMIC

RACE AND ETHNICITY

Following the United States standard birth and death certificates, race and Hispanic origin are collected as separate data items on New Mexico's birth and death certificates, reports of fetal death, and reports of induced terminations of pregnancy (abortions). Although not presented in this report, tribal affiliation is also collected. For birth certificates and fetal death reports, race, Hispanic origin, and tribal affiliation of the infant's mother and father (when available) are provided by the mother. For death certificates, decedent's race, Hispanic origin, and tribal affiliation are provided by an informant (usually a relative or friend) for the deceased.

Federal reports present race and ethnicity (Hispanic origin) as separate categories, where persons of Hispanic origin may be of any race. In the New Mexico Department of Health, race and Hispanic origin are presented using the following categories: American Indian or Alaska Native, Asian or Pacific Islander, Black or African American, Hispanic, and White. The Hispanic category includes persons who were indicated as being of Hispanic origin and whose reported race was White, Other or Unknown. While persons of Hispanic origin are included in the American Indian or Alaska Native, Asian or Pacific Islander, and Black or African American categories, they are not included in the White category. This report follows the New Mexico Department of Health race and ethnicity guidelines.

With the 2006 revision of New Mexico's death certificate, NMVRHS began collecting more than one race (multiple race). Collection of multiple race is in alignment with the U.S. Standard Death Certificate and recommendations made by the U.S. Office of Management and Budget (OMB). For comparability to previous years, other state and federal reports and population estimates used to calculate rates, mortality data in this report have been allocated to single race.

COMPONENTS OF CHANGE

Annual population increases or decreases are due to two components of change, termed Natural Increase and Net Migration.

Natural Increase

Natural Increase is the difference between births and deaths and is calculated by subtracting the number of deaths from the number of births in a population. In those cases where the number of deaths is greater than the number of births, Natural Increase is a negative number.

Net Migration

Net Migration is the difference between immigration (in-migration) and emigration (out-migration) and is calculated by subtracting the number of emigrants from the number of immigrants. Because the United States does not have a national residential registry, intercensal estimates of migration use a variety of data sources.

POVERTY

Poverty information in this report is produced by the United States Census Bureau, and uses income data reported on the American Community Survey (ACS). This measure of poverty is considered the official measure of poverty, as defined by the OMB, and is used by federal agencies for statistical reporting. Government aid programs use different poverty measures for eligibility criteria.

Poverty status is determined for all persons except people who are institutionalized, in college dormitories or in military group quarters. It is determined by comparing household income to 48 thresholds (income cutoffs). Thresholds vary by family size (1 to 9 or more persons) and number of family members under 18 years old (from no children to 8 or more children). Unrelated individuals and two-person families are further differentiated by the age of the reference person (under 65 years old and 65 years old and over).

To determine a person's poverty status, the person's total family income is compared to the poverty threshold appropriate for that person's family size and composition. If the total income of that person's family is less than the threshold appropriate for that family, then the person is considered poor, along with his or her family. If a person is not living with anyone related by birth, marriage, or adoption, then the person's own income is compared with his or her poverty threshold. The 2007 poverty thresholds are shown in the table on the following page.

PER CAPITA PERSONAL INCOME

Per capita personal income figures in this report are obtained from the United States Department of Commerce, Bureau of Economic Analysis. Per capita personal income is calculated by dividing the total personal income by the population. Total personal income is the income received by all persons from participation in production, from government and business transfer payments, and from government interest. Personal income is the sum of net earnings by place of residence, rental income of persons, personal dividend income, personal interest income, and transfer payments. Net earnings are earnings by place of work (the sum of wage and salary disbursements, other labor income, and proprietors' income) less personal contributions for social insurance, plus an adjustment to convert earnings by place of work to a place-of-residence basis. Personal income is measured before the deduction of personal income taxes and other personal taxes and is reported in current dollars (no adjustment is made for price changes).

STATE HEALTH REGIONS

In addition to county level data, this publication also reports data by New Mexico Health and Human Services Geographic Areas. The State Health Regions include the following counties:

- *Region 1 — (Northwest)* Cibola, McKinley, San Juan, Sandoval, and Valencia;
- *Region 2 — (Northeast)* Colfax, Guadalupe, Los Alamos, Mora, Rio Arriba, San Miguel, Santa Fe, Taos, and Union;
- *Region 3 —* Bernalillo;
- *Region 4 — (Southeast)* Chaves, Curry, De Baca, Eddy, Harding, Lea, Quay, and Roosevelt;
- *Region 5 — (Southwest)* Catron, Dona Ana, Grant, Hidalgo, Lincoln, Luna, Otero, Sierra, Socorro, and Torrance.

RESIDENCE DATA

Data are presented by the place where the person normally resided, regardless of where the event occurred.

OCCURRENCE DATA

Occurrence data refers to the place where the event occurred, regardless of the usual residence of the person involved.

Poverty Thresholds for 2007 by Size of Family and Number of Related Children Under 18 Years

Size of Family Unit	Weighted Average Thresholds	Related children under 18 years								
		None	One	Two	Three	Four	Five	Six	Seven	8 or more
One person (unrelated individual)	10,590									
Under 65 years	10,787	10,787								
65 years and over	9,944	9,944								
Two people	13,540									
Householder under 65 years	13,954	13,884	14,291							
Householder 65 years and over	12,550	12,533	14,237							
Three people	16,530	16,218	16,689	16,705						
Four people	21,203	21,386	21,736	21,027	21,100					
Five people	25,080	25,791	26,166	25,364	24,744	24,366				
Six people	28,323	29,664	29,782	29,168	28,579	27,705	27,187			
Seven people	32,233	34,132	34,345	33,610	33,098	32,144	31,031	29,810		
Eight people	35,816	38,174	38,511	37,818	37,210	36,348	35,255	34,116	33,827	
Nine people or more	42,739	45,921	46,143	45,529	45,014	44,168	43,004	41,952	41,691	40,085

Source: U.S. Census Bureau

MORTALITY

CAUSE OF DEATH

ICD Classification

Beginning with 1999 data, cause of death has been coded according to the tenth revision of the World Health Organization's International Classification of Diseases (ICD-10).¹ The International Classification of Diseases (ICD) is a system of classification developed in partnership with the World Health Organization (WHO) and WHO Collaborating Centers. (The North American Collaborating Center is housed at NCHS in Hyattsville, Maryland.) WHO member nations are required to use this classification system for comparability in the collection and classification of health statistics.

New Mexico began systematic record keeping of causes of death and disease morbidity in 1929 and became part of the United States Vital Statistics System in the same year. Revisions to the ICD have occurred almost every ten years since the first version went into effect in 1900. The tenth revision is the exception, going into effect 20 years after the implementation of the ninth revision. Revisions to the ICD are necessary to keep up with advances in medical science. Changes in classification due to revision may lead to discontinuities in cause of death trends. To account for differences between revisions, comparability ratios have been applied to statistics of deaths occurring from 1978-1998.²

In addition to serving as a classification system, the ICD also includes coding rules that allow the nosologist (person who codes cause of death) to select the underlying cause of death - the single condition on the death certificate that is considered most informative from a public health point of view. Also included are definitions (such as "maternal death"), regulations on the compilation and publication of statistics, a prescribed format of the medical certification of death (part of the death certificate), and tabulation lists that indicate cause-of-death groupings that should be used to present comparable mortality data.

Leading Causes of Death

Causes of death are ranked following procedures that are consistent with the recommendations of the 1951 Public Health Conference on Records and Statistics, where causes are ranked by the number of deaths in each rankable cause category.³ With each ICD revision, the list of rankable causes has been revised. The current rankable cause list is based on the "List of 113 selected causes," produced by NCHS.⁴ A separate list for infant deaths is based on the "List of 130 selected causes of infant death."

Alcohol-induced deaths

Causes of death and corresponding ICD-10 codes used by NMVRHS for alcohol-induced deaths are: Alcohol-induced pseudo-Cushing's syndrome, E24.4; Mental and behavioral disorders due to alcohol use, F10; Degeneration of nervous system due to alcohol, G31.2; Alcoholic polyneuropathy, G62.1; Alcoholic myopathy, G72.1; Alcoholic cardiomyopathy, I42.6; Alcoholic gastritis, K29.2; Alcoholic liver disease, K70; Alcohol-induced chronic pancreatitis, K86.0; Finding of alcohol in blood, R78.0; Accidental poisoning by and exposure by alcohol, X45; Intentional self poisoning by and exposure to alcohol, X65; Poisoning by and exposure to alcohol, undetermined intent, Y15. Alcohol-induced causes exclude other unintentional injuries, including motor vehicle accidents; homicides; and other causes not directly related to alcohol use. Infant deaths associated with maternal alcohol use are also excluded.

Drug-induced Deaths

This category includes Drug-induced folate deficiency anemia, D52.1; Drug-induced hemolytic anemia, D59.0; Drug-induced nonautoimmune hemolytic anemia, D59.2; Drug-induced aplastic anemia, D61.1; Secondary sideroblastic anemia due to drugs and toxins, D64.2; Drug-induced thyroiditis, E06.4; Drug-induced hypoglycemia without coma, E16.0; Drug-induced hypopituitarism, E23.1; Drug-induced Cushing's syndrome, E24.2; Drug-induced adrenocortical insufficiency, E27.3; Drug-induced obesity, E66.1; Mental and behavioral disorders due to psychoactive substance use (F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-

F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F17.0, F17.3-F17.5, F17.7-F17.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, and F19.7-F19.9); Other drug-induced secondary parkinsonism, G21.1; Drug-induced dystonia, G24.0; Drug-induced tremor, G25.1; Drug-induced chorea, G25.4; Drug-induced tics and other tics of organic origin, G25.6; Drug-induced headache, not elsewhere classified, G44.4; Drug-induced polyneuropathy, G62.0; Drug-induced myopathy, G72.0; Hypotension due to drugs, I95.2; Acute drug-induced interstitial lung disorders, J70.2; Chronic drug-induced interstitial lung disorders, J70.3; Drug-induced interstitial lung disorder, unspecified, J70.4; Drug-induced pemphigus, L10.5; Generalized skin eruption due to drugs and medicaments, L27.0; Localized skin eruption due to drugs and medicaments, L27.1; Drug-induced gout, M10.2; Drug-induced systemic lupus erythematosus, M32.0; Drug-induced osteoporosis with pathological fracture, M80.4; Drug-induced osteoporosis, M81.4; Other drug-induced osteomalacia in adults, M83.5; Osteonecrosis due to drugs, M87.1; Finding of opiate drug in blood, R78.1; Finding of cocaine in blood, R78.2; Finding of hallucinogen in blood, R78.3; Finding of other drugs of addictive potential in blood, R78.4; and Finding of psychotropic drug in blood, R78.5; Accidental poisoning by and exposure to drugs, medicaments and biological substances, X40-X44; Intentional self-poisoning (suicide) by and exposure to drugs, medicaments and biological substances, X60-X64; Assault (homicide) by drugs, medicaments and biological substances, X85; and Poisoning by and exposure to drugs, medicaments and biological substances, undetermined intent, Y10-Y14. Other unintentional injuries, homicides, and other causes indirectly related to drug use are excluded from this category. Infant deaths due to mother's drug use are also excluded.

Firearm Deaths

Mortality due to firearm use includes suicide, homicide, unintentional injury, legal intervention and undetermined deaths. This category excludes firearm injury deaths due to explosives and other causes indirectly related to firearms. The specific causes of death and ICD-10 codes included in this category are: Accidental discharge of firearms, W32-W34; Intentional self-harm (suicide) by discharge

of firearms, X72-X74; Assault (homicide) by discharge of firearms, X93-X95; Discharge of firearms, undetermined intent, Y22-Y24; and Legal intervention involving firearm discharge, Y35.0.

INJURY AT WORK

Information on deaths due to injuries sustained at work is from the injury at work check box on the death certificate. Except for deaths occurring on tribal lands or military facilities, the Office of the Medical Investigator investigates injury at work deaths that occur in New Mexico.

MATERNAL MORTALITY

Maternal deaths are defined by WHO as "the death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the duration and the site of the pregnancy, from any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes."¹ ICD-10 codes used to capture maternal deaths are: A34, O00-O95, and O98-O99. Causes of death due to maternal mortality are shown in Tables 3.32-3.34.

Maternal mortality rates are calculated by dividing the number of deaths due to maternal causes by the number of live births for the same period, and multiplying by 100,000. Maternal mortality rates are presented as the number of maternal deaths per 100,000 live births. Because only pregnant women are at risk of maternal mortality, live births are used as the denominator to approximate the population of pregnant women.

INFANT MORTALITY

Infant mortality is defined as the death of an infant under one year of age and is often separated into two age groups: neonatal and postneonatal. The neonatal period represents infants under 28 days. Postneonatal infants are 28 days to one year of age. The infant mortality rate is one of the most widely used health indicators and is computed by dividing the number of infant deaths by the number of live births in a period and multiplying by a constant (1,000 or 100,000). Neonatal and postneonatal mortality rates also use the total number of live births as the denominator. Another measure of infant mortality is the infant death rate, which uses the population of

infants as the denominator rather than the number of live births. In this report, only infant mortality rates are used.

Infant mortality rates by race/ethnicity are calculated by using the decedent's (the infant's) race/ethnicity reported on the death certificate and the mother's race/ethnicity reported on the birth certificate.

CALCULATING RATES FOR MORTALITY

In addition to maternal mortality rates and infant mortality rates, this publication uses crude death rates, age-specific death rates, and age-adjusted death rates to measure mortality. Except for maternal and infant mortality rates, which use live births as the denominator, death rates in this report are per 100,000 population.

Crude Death Rate

Crude death rates are the easiest to understand and are computed by dividing the number of deaths by the population, and multiplying by 100,000. Most of the rates in this report are calculated by residence. The New Mexico crude death rate is the number of deaths to New Mexico residents, divided by the population of New Mexico and multiplied by 100,000. Similar rates are calculated by county, region, or other specified area.

Age-specific Death Rate

Age-specific death rates limit the rate to a specific age category. For example, the age-specific death rate for 15-19 year-olds is calculated by dividing the number of deaths to people who were age 15-19 by the number of 15-19 year-olds in the population, and multiplying by 100,000.

Age-adjusted Death Rate

Because crude death rates are influenced by the age composition of the population, comparisons over time or between groups may be misleading. To account for differences in population age distributions, the age-adjusted death rate is used to compare relative mortality risks between groups and over time. This rate should be viewed as an index for comparison, rather than as a direct or actual measure of mortality risk. It is calculated by weighting the age-specific death rates and summing the products. The weights represent the proportion of the population in each age group. Beginning with

1999 data, NMVRHS joined NCHS and other agencies in using the 2000 United States standard population in determining the weights, which are shown in the following table.

Age	Weights
Under 1 Year	0.013818
1 to 4 Years	0.055317
5 to 14 Years	0.145565
15 to 24 Years	0.138646
25 to 34 Years	0.135573
35 to 44 Years	0.162613
45 to 54 Years	0.134834
55 to 64 Years	0.087247
65 to 74 Years	0.066037
75 to 84 Years	0.044842
Over 85 Years	0.015508

Weights for 75-84 and 85+ may be combined if the final category is 75+.

1. World Health Organization. International Statistical Classification of Diseases and Related Health Problems, Tenth Revision. Geneva: World Health Organization. 1992.
2. Anderson RN, Minino AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: Preliminary estimates. National vital statistics reports; vol 49 no 2. Hyattsville, Maryland: National Center for Health Statistics. 2001.
3. National Office of Vital Statistics. Summary report of the Third Annual Meeting of the Public Health Conference on Records and Statistics. 1951.
4. National Center for Health Statistics. Instruction Manual 9: ICD-10 Cause-of-Death Lists for Tabulating Mortality Statistics. Hyattsville, Maryland: National Center for Health Statistics. 2002.
5. Anderson RN, Rosenberg HM. Report of the second workshop on age adjustment. National Center for Health Statistics. Vital Health Stat 4(30). Hyattsville, Maryland: National Center for Health Statistics. 1998.

Bureau of Vital Records and Health Statistics

Located in the Public Health Nurse Memorial Building
1105 St. Francis Drive
Santa Fe, NM 87505

New Mexico Birth and Death Certificates can be obtained at the main office in Santa Fe, and counter customer service at various local public health offices around the state. You can call toll free 1-866-534-0051 to find the nearest location.



ADMINISTRATION

Larry Nielsen, Bureau Chief
Liz Padilla - Financial Specialist
Lisa Sandoval - Administrative Secretary

STATISTICS AND EPIDEMIOLOGY

Larry Nielsen, Director, State Center for Health Statistics

Rosemary Perez-Mendoza - Statistician
Camille Clifford - Epidemiologist
Patricia Drypolcher - Statistician
Irma Montoya - Management Analyst

Nancy Edwards - Training Coordinator
Tony Ortiz - Epidemiologist
Anne Worthington - Epidemiologist

BIRTH AND DEATH REGISTRATION

Renee Valencia, Deputy Registrar, Registration Manager and Nosologist

Evelyn Gaspar - Lex - Data Entry
Martha Lucero - Office Clerk
Jon Branch - Office Clerk

Melinda Roybal - Statistician
Melanie Silva - Office Clerk
Jessica Sosaya - Office Clerk
Peter Varela - Statistician

VITAL RECORDS

Mary Anne Escudero, Acting State Registrar and Vital Records Manager

Laura A. Gutierrez, Local Registrar and Vital Records Supervisor

Rita Encinas, Call Center Unit Supervisor

Percy Armijo - Legal Amendments
Debra Barnes - Legal Amendments
Louanna Gallegos - Order Processing
Genoveva Garcia - Legal Amendments
Ann Madrigal - Remote Office Liaison & Legal Amendments
Maxine Padilla - Order Processing

Monika Romero - Order Processing & Transaction Reports
Carolyn Serrano - Call Center Representative
Debbie Laemmle - Call Center Representative
Rebecca Ulibarri - Order Processing & Transaction Reports
Don Thompson, Issuance Clerk

