

APPENDIX 8: Food Handler and Daycare Worker Exclusion Criteria

Condition	Food Handler	Healthcare Worker	Daycare Worker	Daycare Attendee
Active diarrhea, including positive laboratory result for Campylobacteriosis, Cryptosporidiosis and Giardiasis	Exclude symptomatic persons until diarrhea has resolved, AND proper hygiene measures can be maintained	Exclude symptomatic persons until diarrhea has resolved, AND proper hygiene measures can be maintained	Exclude symptomatic persons until diarrhea has resolved, AND proper hygiene measures can be maintained	Exclude symptomatic persons until diarrhea has resolved, AND proper hygiene measures can be maintained
Salmonellosis	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*	Exclude symptomatic persons until diarrhea has resolved, AND proper hygiene measures can be maintained	Exclude symptomatic persons until diarrhea has resolved.
Shigellosis and STEC	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*	Exclude symptomatic persons until diarrhea has resolved, AND there are TWO consecutive negative stool cultures taken at least 24 hours apart, at least 48 hours after last dose of antimicrobials, if given*

*If a stool culture is positive, then it should be repeated until negative.