
Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.

- Are they breathing? Look, listen and feel. 2.
- If no response, call 911. 3.
- Check for clear airway. If blocked, roll on side and use 4. finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril).
- If not breathing, tilt head back, pinch nose and give 2 6. regular breaths.



- Look, listen and feel. If still not breathing give 1 breath every 5 seconds.
- If there is no response after 3 to 4 minutes give another dose of naloxone.
- Continue breathing until help arrives or they can breath on
- 10. Once breathing, put person in recovery position.





RESCUE BREATHING


- Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub. Are they breathing? Look, listen and feel.
- If no response, call 911. 3.
- Check for clear airway. If blocked, roll on side and use finger sweep to clear. 4
- Roll onto back and give naloxone (1/2 in each nostril). 5.
- If not breathing, tilt head back, pinch nose and give 2 6. regular breaths.
- Look, listen and feel. If still not breathing give 1 breath every 5 seconds.
- If there is no response after 3 to 4 minutes give another dose of naloxone. 8
- Continue breathing until help arrives or they can breath on
- 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch



For contact information and a list of programs and service tim please visit: www.nmhivguide.org

RESCUE BREATHING

- Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.
- Are they breathing? Look, listen and feel.
- 3 If no response, call 911.
- Check for clear airway. If blocked, roll on side and use 4. finger sweep to clear. 5
- Roll onto back and give naloxone (1/2 in each nostril). If not breathing, tilt head back, pinch nose and give 2
- regular breaths.



- If there is no response after 3 to 4 minutes give another dose of naloxone. 8. Continue breathing until help arrives or they can breath on 9.
- their own. 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch



For contact information and a list of programs and service times please visit: www.nmhivguide.org

- **RESCUE BREATHING**
- Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub. Are they breathing? Look, listen and feel.
- 3 If no response, call 911.
- 4. Check for clear airway. If blocked, roll on side and use finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril).
- 6. If not breathing, tilt head back, pinch nose and give 2 regular breaths.

Look, listen and feel. If still not breathing give 1 breath every 5 seconds.

- If there is no response after 3 to 4 minutes give another dose of naloxone. 8.
- 9. Continue breathing until help arrives or they can breath on their own.
- 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch



For contact information and a list of programs and service time please visit: www.nmhlvguide.org

RESCUE BREATHING

- Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.
- Are they breathing? Look, listen and feel.
- If no response, call 911. 3.
- 4. Check for clear airway. If blocked, roll on side and use finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril).
- If not breathing, tilt head back, pinch nose and give 2 regular breaths.
- Look, listen and feel. If still not breathing give 1 breath every 5 seconds. If there is no response after 3 to 4 minutes give another dose of naloxone.
- 8.
- 9 Continue breathing until help arrives or they can breath on their own.
- 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch





Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.

- Are they breathing? Look, listen and feel. 2.
- If no response, call 911. 3.
- 4. Check for clear airway. If blocked, roll on side and use finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril).
- 6. If not breathing, tilt head back, pinch nose and give 2
- regular breaths.
- Look, listen and feel. If still not breathing give 1 breath every 5 seconds. If there is no response after 3 to 4 minutes give another dose of naloxone.
- 9. Continue breathing until help arrives or they can breath on
- 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch 11.

ntact information and a list of programs and service time visit: www.nmhivguide.org



RESCUE BREATHING

Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub. Are they breathing? Look, listen and feel.

- 3.
- If no response, call 911.
 Check for clear airway. If blocked, roll on side and use finger sweep to clear. 4
- 5. Roll onto back and give naloxone (1/2 in each nostril).
- If not breathing, tilt head back, pinch nose and give 2 6.

regular breaths.

- Look, listen and feel. If still not breathing give 1 breath every 5 seconds. 7.
- If there is no response after 3 to 4 minutes give another dose of naloxone. 8.
- Continue breathing until help arrives or they can breath on 10.
- Once breathing, put person in recovery position. Remember, naloxone only works for 30-90 minutes. Watch 11.

For contact information and a list of programs and service tin please visit: <u>www.nmhivguide.org</u>



RESCUE BREATHING

Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.

- 2. Are they breathing? Look, listen and feel.
- 3 If no response, call 911.
- Check for clear airway. If blocked, roll on side and use 4. finger sweep to clear.
- 5 Roll onto back and give naloxone (1/2 in each nostril). If not breathing, tilt head back, pinch nose and give 2 6.
- regular breaths. Look, listen and feel. If still not breathing give 1 breath every 5 seconds.
- If there is no response after 3 to 4 minutes give another dose of naloxone. 8. Continue breathing until help arrives or they can breath on 9. their own.
- Once breathing, put person in recovery position.

Remember, naloxone only works for 30-90 minutes. Watch For contact information and a list of programs and service times, please visit: www.nmhivguide.org



RESCUE BREATHING Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub. Are they breathing? Look, listen and feel.

- 2.
- 3. If no response, call 911.
- 4. Check for clear airway. If blocked, roll on side and use finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril).
- 6. If not breathing, tilt head back, pinch nose and give 2 regular breaths.
- Look, listen and feel. If still not breathing give 1 breath every 5 seconds.
- If there is no response after 3 to 4 minutes give another dose of naloxone. 8.
- Continue breathing until help arrives or they can breath on 9. their own.
- Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch



For contact information and a list of programs and service times please visit: www.nmhlvguide.org

RESCUE BREATHING

- Check responsiveness. Ask, "Are you okay?" Shake foot. Use sternum rub.
- Are they breathing? Look, listen and feel.
- If no response, call 911.
- 4. Check for clear airway. If blocked, roll on side and use finger sweep to clear.
- Roll onto back and give naloxone (1/2 in each nostril). 6. If not breathing, tilt head back, pinch nose and give 2 regular breaths.
- Look, listen and feel. If still not breathing give 1 breath every 5 seconds.
- 8. If there is no response after 3 to 4 minutes give another dose of naloxone. 9
- Continue breathing until help arrives or they can breath on their own.
- 10. Once breathing, put person in recovery position.
- Remember, naloxone only works for 30-90 minutes. Watch



For contact information and a list of programs and service times, please visit: www.nmhivguide.org

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate - Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16.

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril. **Evaluate and support –** Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16.

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate - Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth

(Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds. **Evaluate** – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now? Do they need another dose of naloxone?

Nelevere leets 20.00 minutes. Cook help on

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16.

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate - Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16.

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing - Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

<u>Naloxone lasts 30-90 minutes</u>. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

Medication-Spray ½ dose (vial) of naloxone into each nostril.

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!

Thanks to Chicago Recovery Alliance for the original SCARE-ME material in 2001, changes and updates by D. Zurlo 10-28-16.

Things to Do With an Opiate Overdose

Stimulation - Are they responsive?

Shout their name, shake their foot, use a sternum rub.

Call 911 - Tell 911 someone is unresponsive.

Airway – Look-Listen-Feel. Open airway and clear mouth (Mouth sweep with curved fingers if there is a blockage).

Rescue Breathing – Give 1 breath every 5 seconds.

Evaluate – Are they any better? (Look-Listen-Feel).

 $\begin{tabular}{ll} \bf Medication- Spray~\%~dose~(vial)~of~naloxone~into~each~nostril. \end{tabular}$

Evaluate and support – Is the person breathing now?

Do they need another dose of naloxone?

Naloxone lasts 30-90 minutes. Seek help and support in case the person overdoses after the naloxone wears-off!