



# STATE OF NEW MEXICO VACCINATION CARD

CHILD'S NAME: FIRST \_\_\_\_\_ MIDDLE \_\_\_\_\_ LAST \_\_\_\_\_ DATE OF BIRTH: MONTH \_\_\_\_\_ DAY \_\_\_\_\_ YEAR \_\_\_\_\_

## RECOMMENDED INTERVALS – CHILD'S AGE IN MONTHS (mark colored squares for each completed vaccine)

VACCINE	DOSE	0	1	2	3	4	5	6	7	8	9	10	11	12	18	24	36	48	Catch up	DATE	CLINIC
DIPHTHERIA, TETANUS & PERTUSSIS	1ST																				
	2ND																				
	3RD																				
	4TH																				
	5TH																				
HAEMOPHILUS INFLUENZAE B	1ST																				
	2ND																				
	3RD																				
	4TH																				
HEPATITIS A	1ST																				
	2ND																				
HEPATITIS B	1ST																				
	2ND																				
	3RD																				
INFLUENZA	1ST																				
	2ND																				
	3RD																				
	4TH																				
	5TH																				
	6TH																				
ROTAVIRUS	1ST																				
	2ND																				
	3RD																				

Allow at least 6-mo. interval between doses 3 & 4

3rd dose not needed if Comvax or Pedvax-Hib vaccine used for dose 1 & 2

(Record any additional doses on the back side)

2 shots between 6 & 12 months, given min. 1 month apart

Rotateq requires 3-doses; Rotarix requires 2-doses

## Is your child up-to-date?

To check, place a ruler at the current age of your child. Are there X's in all the colored boxes to the left of the ruler? If not, ask your health care provider to check your card and give any needed shots. **Keep in a safe place and ALWAYS bring with you to any type of child health activity.**

## RECOMMENDED INTERVALS – CHILD'S AGE IN MONTHS (mark colored squares for each completed vaccine)

VACCINE	DOSE	0	1	2	3	4	5	6	7	8	9	10	11	12	48	Catch up	DATE	CLINIC
MEASLES, MUMPS & RUBELLA	1ST																	
	2ND																	
PNEUMO-COCCAL CONJUGATE (PCV-13)	1ST																	
	2ND																	
	3RD																	
	4TH																	
POLIO	1ST																	
	2ND																	
	3RD																	
	4TH																	
VARICELLA	1ST																	
	2ND																	

		ADOLESCENT SERIES			Age in years: 11-12 13-15 16-18			DATE	CLINIC
HUMAN PAPILLOMA-VIRUS	1ST								
	2ND								
	3RD								
MENINGO-COCCAL ACWY	1ST								
	2ND								
MENINGO-COCCAL B	1ST								
	2ND								
	3RD								
Tdap <sup>†</sup>	1ST								

Allow 6-12 month interval between dose 1 & 2  
3rd dose needed only if <5 months between dose 1 & 2, OR if starting series ≥15 yrs., then 1-2 mo. interval between dose 1 & 2, and 6 mo. interval between dose 1 & 3

Allow 5-year interval between dose 1 & 2

High-risk or upon request—ask your provider. 2 or 3 doses needed depending on vaccine.\*

<sup>†</sup>TETANUS, DIPHTHERIA & PERTUSSIS

\*Trumenba has 2-dose schedule, 0 and 6 mos.; and a 3-dose schedule, 0, 1 to 2, and 6 mos. Bexsero has 2-dose schedule, 0 and 1 mo.